Robert E. Gregg

.... 1

12:

A STUDY OF ANT POPULATIONS AT THE PLAINS-FOOTHILL BORDER

by

August Conklin

B.A. University of Colorado, 1942 M.A. Claremont Graduate School, 1951 M.A. University of Colorado, 1962

A thesis submitted to the Faculty of the Graduate School of the University of Colorado in partial fulfillment of the requirements for the degree of Doctor of Philosophy Department of Biology

This Thesis for the Doctor of Philosophy Degree by

August Conklin

has been approved for the

Department of

Biology

by

Robert E. Gregg

Erik K. Bonde

Listella listere was units and

torn did not

Date_

10

÷

Conklin, August (Ph.D., Zoology)

A Study of Ant Populations at the Plains-Foothill Border Thesis directed by Professor Robert E. Gregg

Three quadrats, one hundred meters to the side, were located at the plains-foothill border region southwest of Boulder, Colorado, near Bluebell Canyon. One quadrat was in a wooded area, another in a meadow, and the third in an ecotone, half-meadow, half-wooded area. Ants were intensively collected in these quadrats for three summers, from 1965 through 1967. The summer of 1968 was utilized for soil moisture and temperature determinations, as well as for actual counts of ant colonies.

A total of 25 species and subspecies was discovered. Of these 25 species, nine were species found in two quadrats, which provide evidence of a distinct boundary between the lower edge of the foothill zone and the plains, since eight of these nine were found in the ponderosa pine forest and the ecotone, but did not penetrate the meadow. The other species was found in the meadow and ecotone, but did not penetrate the ponderosa pine forest. There were two additional species in the meadow and five in the forest which did not penetrate the ecotone.

The Lincoln Index was utilized in this study to determine its efficacy in estimation of the size of selected ant colonies. It was found to be inaccurate, since use of this index results in great underestimation of ant colony size.

Individual nest stability and nest number constancy were also examined. The majority of the 25 species and subspecies are quite unstable in their nesting habits, and nest number was somewhat inconstant.

Total counts of ant nests were made, and these, plus the number of nests of the various species, made it possible to determine the ecological dominants in each quadrat for each year. It was also found that there were more species of ants in this study in the wooded quadrat, which had a slope facing north by northwest, than in the ecotone or in the meadow. Observations were also made with reference to compound nests, polydomous nests, and species which have muptial flights in late summer.

Soil moisture and temperature determinations were made in the various quadrats, and the probable causes for nest mortality or exodus were examined.

This abstract is approved as to form and content. I ... a recommend its publication.

Signed

. .

. 1

Faculty member in charge of dissertation

ACKNOWLEDGMENTS

As the writer of this dissertation, I wish to express my heartfelt thanks and deep appreciation to all those who not only offered helpful suggestions and criticisms, but also provided considerable encouragement.

Special thanks must be given to the following:

To my wonderful family, especially to my wife, Jane, who not only gave me much needed encouragement, but spent many hours in typing the manuscript, and to my sons Tom and Jim. Without Tom, who was my invaluable good right arm in the field, this study would probably not have been accomplished. Jim also gave me encouragement when I needed it.

To Doctor Gregg, who not only made many beneficial suggestions, but also spent considerable time in assisting me with my ant classifications. Doctor Gregg's advice and contributions were invaluable to this project.

To Doctors Erik K. Bonde, John W. Marr, Charles H. Norris, Robert W. Pennak, Sam Shushan, and Hobart Smith with whom I had many pleasant and worthwhile associations, and who provided assistance as well as inspiration to my program.

TABLE OF CONTENTS

PAGE

INTRODUCTION
DESCRIPTION OF THE RESEARCH AREA
METHODS AND MATERIALS 28
RESULTS
DISCUSSION
Distribution of the various species
Stability and frequency of colonies
Lincoln Index
Climatic conditions
Foraging activities
Nest evacuation
Compound nests
Polydomous nests 101
Census of colonies 102
Nuptial flights
SUMMARY 109
LITERATURE CITED
APPENDIX

•

LIST OF TABLES

TABLE		PAGE
I	LIST OF ANT SPECIES FOUND IN 1965 IN QUADRAT 1	116
II	LIST OF ANT SPECIES FOUND IN 1965 IN QUADRAT 2	118
III	LIST OF ANT SPECIES FOUND IN 1965 IN QUADRAT 3	124
IV	LIST OF ANT SPECIES FOUND IN 1966 IN QUADRAT 1	125
v	LIST OF ANT SPECIES FOUND IN 1966 IN QUADRAT 2	128
VI	LIST OF ANT SPECIES FOUND IN 1966 IN QUADRAT 3	135
VII	LIST OF ANT SPECIES FOUND IN 1967 IN QUADRAT 1	136
VIII	LIST OF ANT SPECIES FOUND IN 1967 IN QUADRAT 2	139
IX	LIST OF ANT SPECIES FOUND IN 1967 IN QUADRAT 3	147
X	RANGES AND ZONATION OF SPECIES	149
XI	SPECIES COMPOSITION OF QL	151
XII	SPECIES COMPOSITION OF Q2.	152
XIII	SPECIES COMPOSITION OF Q3	153
XIV	ANT SUBFAMILY COMPOSITION.	154
XV	ESTIMATION OF ANT COLONY SIZE BY THE LINCOLN INDEX	155
XVI	MORNING TEMPERATURE READINGS	156
XVII	NOON TEMPERATURE READINGS.	159
XVIII	AFTERNOON TEMPERATURE READINGS	162
XIX	WEEKLY MORNING TEMPERATURE AVERAGES IN ALL QUADRATS	165
XX	WEEKLY NOON TEMPERATURE AVERAGES IN ALL QUADRATS	166

...

TABLE

 I = 1 = 1 = 1 		
IXX	WEEKLY AFTERNOON TEMPERATURE AVERAGES IN ALL QUADRATS	16 7
XXII	WEEKLY MORNING TEMPERATURE EXTREMES IN ALL QUADRATS	168
XXIII	WEEKLY NOON TEMPERATURE EXTREMES IN ALL QUADRATS	169
XXIV	WEEKLY AFTERNOON TEMPERATURE EXTREMES IN ALL QUADRATS	170
XXA	U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS FOR SUMMER 1965	171
XXVI	U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS FOR SUMMER 1966	173
XXVII	U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS FOR SUMMER 1967	174
XXVIII	U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS FOR SUMMER 1968	175
XXIX	SOIL MOISTURE IN QUADRAT 1 IN 1968	176
XXX	SOIL MOISTURE IN QUADRAT 2 IN 1968	181
XXXI	SOIL MOISTURE IN QUADRAT 3 IN 1968	186
- XXXII	AVERAGE PER CENT MOISTURE IN ALL QUADRATS IN 1968	191
· XXXIII	ANT COLONY SIZE BY ACTUAL COUNT	192

PAGE

LIST OF FIGURES

• (

;

FIGURE		PAGE
l	Q3-The meadow. View from the southeast	. 12
2	Q3-The meadow. View from the northeast	. 14
3	Q1-The ecotone. View from the southeast	. 16
4	Q1The ecotone. View from the east	. 18
5	Q2-The wooded area. View from the north	20
6	Q2-The wooded area. View from the east	22
7	Ant nests in quadrat 1	30
8	Ant nests in quadrat 2	31
9	Ant nests in quadrat 3	32
10	Aerial view of Plains-Foothill border southwest of Boulder, Colorado	33
n	Location of quadrats.	35
	L	

INTRODUCTION

This investigation is a study of the ant faunal distribution of the plains-foothill area near Boulder, Colorado, and is concerned with the distribution of ants over a three-year period in three types of environments, namely, a forest, a meadow, and an ecotone of forest and meadow. All three of these environmental units are located at the plains-foothill border where there is a peripheral convergence of the grassland formation of the plains and the montane coniferous formation. The foothills consist of rounded hills, upturned sedimentary beds, and mesas, and near Boulder, the land forms are expressed as Mesa and Dakota ridge forms to which the border is confined. Gregg (1963) writes that the plains-foothill region is ecologically significant in that climatic conditions are typical neither of the plains nor of the foothills. Also, the topography is distinct, and a variety of soils are present.

In a review of the literature, no study strictly similar to mine was found. However, there are a few studies which embrace some of the objectives of my study in the areas of ant species distribution, nest stability, the use of the Lincoln Index as a tool for population census, and the effects of climate on ant distribution.

Many studies are concerned with distribution of ant species. Talbot (1934), for example, studied the distribution of ant species in the Chicago region with reference to ecological factors and physiological tolerances, and Barrett and Felton (1965) examined the

distribution of the wood ant, Formica rufa Linnaeus, in southeast England. Brian, Hibble, and Stradling (1965) published a paper on ant pattern and density in a southern English heath, and Francour (1966) established the distribution of certain species of ants in the Quebec region. Wheeler and Wheeler (1963) wrote a volume on the distribution of ants in North Dakota. Wheeler and Wheeler (1965) reported distribution of Veromessor lobognathus in South Dakota, whereas Gregg (1963), in a large volume, described the distribution of the ants of Colorado. Creighton examined habits and distribution of many ant species in numerous articles and also wrote a comprehensive volume on the ants of North America (1950). In his book on social insect populations, Brian (1965) discussed the critical factors in ant distribution, which are food and climate. Pickles (1937 and 1938) described changes in ant nest distribution, and Brian (1952) worked on the replacement of nests under stones. Yasuno (1965) made a study of the stability of the ant populations in grassland at Mt. Makkôda, Japan. Golley and Centry (1964) in a study on bioenergetics of the southern harvester ant, Pogonomyrmex badius, comment on their findings about movements of the ant hills for that particular species. They learned that movement of ant hills was a common phenomenon, and these movements appear to be most common in the fall. Wheeler (1910) states that there are a few species of ants besides the nomadic Dorylinae that seem to "enjoy" an occasional change of residence. Hə writes of Wasmann, who has shown that Formica sanguinea often has summer and winter residences analogous to the city and country homes of wealthy people. Brian (1965) discusses the hazards which are present in helping to cause instability in ant nests and cites work

done by Gosswald and Kloft (1961) in connection with damage done to ant nests by field mice. Brian (1965) delineates the modifying factors which affect the food supply, which, in turn, affects the stability of ant nests, and discusses interspecific and intraspecific competition as factors in stability of nests. Such competition introduces ant territorialism, which has been widely studied since Elton (1932) showed its existence in Formica rufa; Brian (1965) and Yasuno (1965) among others in the more recent literature, and Forel (1928) in the older literature, write on this subject. Pleometrosis, or the occurrence of more than one queen in a single nest of ants, is another cause for instability in ant nesting, and Schneirla (1949) and others have observed that pleometrosis has a considerable influence in the increase of population size. Schneirla (1957) found that the reasons for restlessness in the behavior of the Dorylinae in their characteristic nest instability are the need for food and their response to an internal brood change.

Stability of ant nests and the time of year for the nuptial flight, two objectives of my study, relate to ant movements for habitat selection. Wilson and Hunt (1966) examined habitat selection by the queens of two field-dwelling species of ants. These investigators concluded that major habitat selection is performed primarily during the nuptial flight but that the microhabitat (nest site) is selected only after searching on the ground following the flight. Talbot (1966) discussed flights of the ant <u>Aphaenogaster</u> <u>treatae</u> and found that this field ant has unusual flights, in that they take place at high temperatures (78° to 88° F), but only if light is reduced, either by continuous gray skies, or, more frequently, by moving clouds. Most flights take place when temperatures have been too high for workers to forage until clouds come by. Numerous studies have been made in ant behavior, and Sudd's book, <u>An Introduction to the Behavior of Ants</u>, describes many aspects of behavior in ants and provides an extensive list of references as well.

Numerous ecological studies have been performed on ants in the Boulder area or elsewhere in the State of Colorado, and beginning in 1910, W. W. Robbins of the University of Colorado Department of Biology made a survey of the ants of northern Colorado which was the first attempt to summarize the Colorado ant fauna. Periodically, since then, many species have been described from the state. L. F. Byars made an ecological study of ants in Boulder County in 1936 for his Master's thesis, and R. E. Gregg (1947) made a survey of ants in Colorado in a paper entitled "Altitudinal Indicators Among the Formicidae". E. D. Delfin (1954) conducted a general entomological survey in the Gregory Canyon area near Boulder, Colorado, and H. F. Borchert (1956) examined the distribution of ants of five canyon bottoms in the Boulder area. J. T. Browne (1958) did a Master's thesis study on ant distribution in Gregory Canyon, whereas M. E. Smith (1962) made a distributional study of formicids in the vicinity of Valmont Butte, Valmont, Colorado, near Boulder for a Master's thesis. W. H. Taussig (1962) also performed a Master's thesis study on ants in an ecological work on Formica neorufibarbis gelida Wheeler in the alpine tundra of Colorado, and, as mentioned previously, R. E. Gregg (1963) wrote a very detailed volume entitled The Ants of Colorado.

The Lincoln Index, which is a mark-recapture method of population census, has been used very extensively in fish and wildlife work, and according to Osburn (1953) the Lincoln Index was first developed by F. C. Lincoln (1930) to estimate North American duck populations. Jackson (1936), while studying tse-tse fly populations in Africa, derived, independently, a method identical to that of Lincoln's. The principle of the Lincoln Index is simple. A random sample of individuals is marked, and these are then released. Later, after a dispersal period, another random sample is collected and examined. The second catch may include a proportion of individuals recognized by their marks as having been caught in the marking period. The second catch should have the same proportion of recaptured insects to the total taken as the proportion initially marked to the total population.

population size = total marked x total caught when recapturing marked recaptures

Jackson's series of papers on tse-tse flies, reviewed by Andrewartha and Birch (1954), appears to present the first application of the method to insect populations, and in recent years this approach has been applied to many other insects such as butterflies, grasshoppers, locusts, and ground beetles; however, many investigators found the Lincoln Index was too crude to use without modification. Jackson (1939) and others recognized the need for consideration of birth, death, and migration rates, and Allee (1949) discussed limitations and advantages of the Lincoln Index. Odum and Pontin (1961) estimated the population density of the underground ant

Lasius flavus by tagging with radioactive phosphorus, and Golley and Gentry (1964) employed a similar tagging procedure with the southern harvester ant, <u>Pogonomyrmex badius</u>. The mark-recapture technique produced contradictory results. Chew (1960), who worked with <u>Pogonomyrmex occidentalis</u> in Arizona, used similar methods and did not achieve satisfactory results.

Odum (1959) states, "Temperature and moisture are so generally important in terrestrial environments and so closely interacting that they are usually conceded to be the most important part of climate." Knight (1965) states that the interaction of a number of climatic or microclimatic factors is most important ecologically, because it will often indicate why a particular organism is present or absent. In The World of an Insect by Chauvin (1967) the author writes that the German Rudolf Geiger once pointed out that what is taking place at a height of six feet has little bearing on what is happening by a man's feet, and therefore, the microclimate is extremely important to the insect. According to Gregg (1963) the two cardinal factors of temperature and moisture are correlated with the distribution of ants regardless of what else may be operating also, since it seems to be true that the number of species shows trends that parallel meteorological changes. Where conditions of temperature and moisture are both optimal, the greatest concentration of forms or taxonomic complexity are seen. Since so many ants are fossorial, they can obtain moisture deep in the soil, and therefore, temperature may appear to be more limiting for these, particularly a thermophilous group such as the Formicidae.

The literature on the subject of climate and insects is voluminous, as can be seen by examination of most texts of animal ecology, for example, Andrewartha and Birch's large volume (1954). However, very little information is available on the influence or importance of continuously varying humidities (Messenger, 1959). Messenger writes that rainfall is a major factor controlling the moisture content of soil, and therefore can be a major factor limiting the development and survival of some soil-inhabiting forms.

Wheeler (1910) says that the length of embryonic, larval, and pupal life in ants appears to be highly variable and to depend very intimately on temperature. A rise in temperature induces both females and workers to lay eggs and accelerates the growth of the larvae. Other factors being equal, development of eggs within the ovaries, the deposition of eggs, the feeding and growth of the larvae, pupation, and hatching all appear to be determined by temperature. The degree of heat suitable to the species probably varies for the different stages of development. Wheeler writes that the optimum temperature for northern ants lies between 70° and 80° F. At low temperatures, the exact level being different for each species, the insect comes to rest and shows no spontaneous activity; however, as temperature is raised, the ant becomes normally active, then excessively active, and ultimately passes into a state of heat stupor, followed by death.

Temperature, therefore, has a marked effect upon growth, and it also has an important effect upon metabolism. Wigglesworth (1965) writes that whereas the metabolism of warm-blooded animals is depressed as the external temperature rises, the metabolism of

cold-blooded animals increases. The increased activity which the insect shows with rising temperature is an indication of the increase in metabolism. What happens to the extra energy produced in the resting, or narcotized, animal as the result of raising the temperature is not known, but much of it must be expended by the augmented movements of the internal organs. In the developmental stages of insects, the energy is expended on growth, which is correspondingly accelerated. Of course, the optimum temperature may vary with the humidity. Odum (1959) states that temperature and moisture interact upon one another but this interaction depends upon the relative as well as the absolute values of each factor. Thus. temperature exerts a more severe limiting effect on organisms when moisture conditions are extreme, that is, either very high or very low, than when such conditions are moderate. Likewise, moisture plays a more critical role in the extremes of temperature. Diminution in the water content usually depresses metabolism and retards development, and a lower relative humidity results in a longer period of larval development. Sometimes the effect of desiccation is purely mechanical; the chorion of the egg may become too hard for the embryo to break through, or the fully developed insect may lack sufficient volume of water in its blood to rupture its pupal sheath. Sometimes the rate of development is retarded at very high humidities. High temperatures cause greater evaporation through the integument, and may increase the permeability of the cuticle to water, which will result in water loss in insect eggs with a resultant decrease in population.

Ants have very definite responses to both temperature and humidity; for instance, Sudd (1967) states that early experiments with Aphaenogaster fulva, Camponotus pennsylvanicus, and Acanthomyops latives showed that these ants moved sluggishly below 15° C. and that heat stupor began to set in at about 35°. They were rapidly killed by exposure to 50°, but inside this range they rest in a temperature of 24° - 27°. Sudd believes that ants are strongly affected by humidity and refers to their humidity sense. Low humidities cause death from excessive loss of water, especially when combined with a high temperature. At low humidities, earthnesting ants simply retreat deeper into the soil. Larvae lose water to the air even faster than adults, and queens lose more water than workers, but pupae are far more resistant to dessication than the other stages. The workers transport the ant young from place to place, thus utilizing to the advantage of the developing young the ever-varying temperature and humidity of the soil.

The objectives of my study are as follows:

1. To present data to indicate that ants may offer significant evidence that the plains-foothill border is an important biological boundary.

2. To determine the stability and frequency of ant colonies from one year to the next.

3. To determine the value of the Lincoln Index in making a census of an ant colony.

4. To relate climate to ant distribution.

5. To determine foraging activities of some ants.

6. To examine the probable causes for exodus from an ant nest.

7. To note the various species of ants which live in compound nests (plesiobiosis).

8. To locate nests of ants which appear to have branched from the original colony (polydomous nests).

9. To census populations of ant nests and discover the probable dominants among ants in the area.

10. To discover which ant species in the research area have nuptial flights in late summer.

۰.

10

2.2

DESCRIPTION OF THE RESEARCH AREA

In the summer of 1965 three sites were selected for study. One of these three areas is in a meadow near, and northeast of, Bluebell Canyon, and the other two areas are on the slopes of a ridge east of Bluebell Canyon proper; these two are actually at the approaches to the canyon less than one-fourth mile from the canyon itself. All three sites are very near to each other, as evidenced by the fact that temperature readings for all three stations could be completed in 20 to 25 minutes. Each of the three locations was set up as a quadrat with 100 meters to a side. The meadow was designated as Q3 and had a gentle slope, which faced north by northwest (See Figures 1 and 2). Q2 was the symbol applied to a rather heavily wooded area, which had a sharper slope than that of the meadow, but it also sloped north by northwest. The wooded area was located in the ponderosa pine-grama grass community. Q1 was the , ecotone, an area half wooded and half meadow, and it had a rather abrupt slope which faced east by southeast, but mostly east (See Figures 3 to 6).

Of the three research sites selected in this study, the forest, the meadow, and the ecotone, the forest and ecotone are in the Transition or Submontane Zone of Merriam (1898), which has an elevation of 6,000 - 8,000 feet (Gregg, 1963). The meadow is in the upper Plains or Upper Sonoran Zone, which has an elevation of 4500 - 6000 feet. Merriam also used other life zones for the division of North America into vegetational and forest units, but my

L.

--

3

. 1

11



...

;

. (

Figure 2. Q3-The meadow. View from the northeast.

۰.



ł

-:

:

...

Figure 3. Q1-The ecotons. View from the southeast.

40

i



٤

2:

3

. 1

Figure 4. Cl-The ecotone. View from the east.

۰.

10

...7



Figure 5. Q2-The wooded area. View from the north.

1

1 3

.....

-:

3

• 1

3

L



Ł

-:

3

.(

Figure 6. Q2-The wooded area. View from the east.

•



research area is not concerned with these other zones. Merriam's life zones were established on a climatic basis and set forth what he believed were temperature laws governing terrestrial distribution.

The area where I worked is at the southwest border of Boulder, Colorado, a city located 24 miles northwest of Denver at 40° north latitude, and 105° 16' west longitude. The front range of the Colorado Rockies rises steeply from the base of the foothills, which are approximately one mile high, to the Continental Divide 30 miles west of Boulder, which has mountains 11,000 to 14,000 feet in altitude. The mountain front can be divided into the foothills themselves, and a broad benchland, or mesa terrace, the benchland really being an upper extension of the plains.

Bluebell Canyon, which is found in the Transition Zone, has an approximate elevation of 6,050 feet, and has its mouth opening to the plains area. Its headwaters lie between the Flatirons, which are erect slabs of red sandstone on the east face of Green Mountain, and belong to the Fountain Formation. This formation was produced by deposition during the Pennsylvanian Period from 320 million to 280 million years before the present (Rodeck, 1964). The transition zone has many small intermittent streams, which are dry in the fall and winter, and the soils here are generally coarse-textured and shallow, with a soil water content which is usually low.

According to Byars (1936), the prevailing wind for Boulder County is westerly, having passed over the mountains; consequently, as the wind passes over the warmer plains, it takes up moisture and has a drying effect. The plains near the foothills are protected from the full force of the wind.

The climate of the plains near the front range of the Rocky Mountains, according to Armin (1963), is characterized by a relatively low humidity, a large amount of sunshine, light precipitation, moderate winds, a large daily range in temperature, high day temperatures in summer, and occasions when the temperature is subzero in the winter. As the distance increases from the foothills, conditions become moderately xerophytic. The nitrogen content of the soil and its organic nature increases with elevation, up to a certain limit, and there is a tendency for the soils to change from an acid to an alkaline condition from the alpine to the plains zone.

The climate of the foothills differs from that of the plains in general by having a narrower range of fluctuation of temperatures; higher minimum temperatures are found in the foothills region. According to Gregg (1963), in the Rocky Mountains the drop in average temperature with increasing elevation, or the lapse rate, is approximately 3° E per 1000 feet, lower than this in the winter, and slightly above in the spring and early summer. Air drainage has a greater influence than elevation, for cool winds of the montane zone (elevation 8,000 to 10,000 feet) bring mountain temperatures to narrow canyons, but the slopes facing the plains have almost the same temperatures as those of the plains. Wide valleys are warmer than narrow ones, and slope direction is important, since south-facing slopes obviously are warmer and drier than the north-facing ones. Wind velocities are low in the foothills, increasing from there both up the mountains and down onto the plains, and with an increase in altitude there is a substantial increase in precipitation. Moderate temperatures and moisture supplies combine to make conditions in the

foothill area very suitable for many organisms. The terraces, or small mesas, upon which two of my quadrats are located appear to be excellent environments for ants, and attract species from both the mountains and plains.

Armin (1963) states that the plant life of the plains zone is characterized by dry grassland with few or no trees. <u>Pinus</u> <u>ponderosa</u>, or ponderosa pine, sometimes occurs on dry bluffs where rock is exposed. The more distinctive grasses are the bunch grasses <u>Andropogon</u> and <u>Sorghastrum</u>, the wire grass <u>Aristida</u>, porcupine grass <u>Stipa</u>, the very common buffalo grass (<u>Buchloe</u>) and grama grass <u>Stipa</u>, the very common buffalo grass (<u>Buchloe</u>) and grama grass <u>Bouteloua</u>). Often herts are present, such as sunflower (<u>Helianthus</u> <u>annuus</u>), goldenrod (<u>Solidago</u> spp.) catnip (<u>Mentha spicata</u>), wild cucumber (<u>Gucurbita foetidissima</u>), ground mallow (<u>Malva neglecta</u>), horse thistle (<u>Cirsium spp.</u>), burdock (<u>Anthemis cotula</u>), ragweed (<u>Ambrosia trifida</u>), sage (<u>Artemisia glauca</u>), yucca (<u>Yucca glauca</u>), sand bur (<u>Cenchrus pauciflorus</u>), milkweed (<u>Asclepias speciosa</u>), prickly pear cactus (<u>Opuntia polyacentha</u>), mullein (<u>Verbascum</u> <u>thapsus</u>), and others. Many of these are not limited to the plains zone.

Ponderosa pine (<u>Pinus ponderosa</u>) is the dominant tree of the foothills zone. This tree occurs as a climax stand on gently rolling surfaces. Occasionally found between these trees are Rocky Mountain: juniper (<u>Juniperis communis</u>), squaw currant (<u>Ribes cereum</u>), and grasses such as buffalo grass (<u>Buchloe</u>). Other dominant herbs may be <u>Carex</u> spp. (sedges), wild geranium (<u>Geranium fremontii</u>), goldaster (<u>Chrysopsis foliosa</u>), and mountain parsley (<u>Harbouria</u> <u>trachypleura</u>). The valley floors also have grasses and many species

of herbs and sedges. Poison ivy (<u>Rhus radicans</u>), and sumac (<u>Rhus</u> <u>glabra</u>) are quite common in a small clearing north of the wooded quadrat. Of the grasses in my quadrats, blue grass (<u>Poa pratensis</u>). brome grass (<u>Bromus brizaeformis</u>), timothy (<u>Phleum pratense</u>), and cheat grass (<u>Bromus tectorum</u>) appeared quite prevalent.

10

~:
METHODS AND MATERIALS

The boundaries of the three quadrats were marked by using wooden stakes painted at the top with red paint which were driven into the ground at ten-meter intervals on each side of the quadrats. The writer and his son then crisscrossed back and forth through each quadrat laboriously turning over rocks and logs (of which there were only a few), and, in short, did everything possible in order to locate any ant colonies, which were to be found in these quadrats. Once the ants were exposed, they were collected into 2 and 4-dram vials (depending upon the size of the ant) containing 85% ethyl alcohol, where they were killed and preserved, and a minimum of 12 ants was collected as a sample from each colony. An aspirator was tried for removing them from the nest, but it was found to be too cumbersome. Sampling of the ants from each nest was accomplished as carefully as possible so as not to unduly disturb the colony.

Into each vial was inserted a slip of paper with the date and number of colony designated, as well as the quadrat number. Later, after the vial was emptied for removal of debris, and identification of the species had been made, the ants, and a slip with the species name, were included in the vial. The vial's cap was then labelled with the quadrat and colony number. A number was painted on the stake with black paint, and the stake was driven into the ground beside the colony. Subsequent stakes were also numbered and inserted beside the nests in the sequence in which these nests were found.

In the summers of 1966 and 1967, the same procedure was followed in carefully combing each quadrat for all possible colonies located within them. In 1966, when new nests were discovered, numbered stakes in a continued sequence of numbers from those used in 1965 were also inserted beside them. Since 1967 was to be the last summer for locating colonies, stakes were not driven beside the newly discovered nests that summer. A detailed map was prepared for recording the locations of the various ant colonies for each quadrat in 1965, and in the two subsequent summers the newly discovered colonies were also included in the map (Figures 7 to 9). Figure 10 is an aerial view of the general area in which the quadrats are located, and Figure 11 shows the specific location of the three quadrats. A binocular microscope was used for identification of the various ants, and in just a very few cases a monocular compound microscope with magnification of 100x was used for a very small. precise taxonomic characteristic. Robert E. Gregg's book, The Ants of Colorado (Gregg, 1963), was used as the reference for taxonomic keys in the identification of the various species, and William S. Creighton's book, The Ants of North America (Creighton, 1950), was also utilized, especially for his outline drawings of ants.

A segment of this study was concerned with an attempt to estimate population size of the ant colonies by using the Lincoln Index which, as previously mentioned, is a mark and recapture method of census. The ants which were selected to test this method consisted of two larger species, so that marking and recognition would be facilitated. Colonies of the species <u>Camponotus</u> (<u>T.</u>) <u>vicinus</u> Mayr and <u>Formica</u> (<u>N.</u>) <u>pallidefulva</u> Latreille were utilized. Various







ł.

~:·

1

. 1

Figure 10. Aerial view of plains-foothill border, southwest of Boulder, Colorado.

. . 1



Figure 11. Location of Quadrata

• 1

10

-:

•••

.1



substances were tried, but it was found that fingernail polish was quite satisfactory for the purpose of marking. The ants which were selected for census were chosen from a location outside of the study quadrats, inasmuch as the quadrats were to be carefully examined over a period of several summers. The procedure followed in the use of the Lincoln Index method of estimation of ant colony size was very similar to that used by Chew (1959). First, 50 to 100 worker ants were collected from a colony; next, these individuals were etherized, and each ant was marked on the dorsum of the thorax with a spot of red fingernail polish; third, the marked ants were counted and then released at the colony entrance after they had revived (the few ants whose legs became bound up by the application of too much polish were removed); fourth, a second sample of ants was collected 24 hours later, and the marked and unmarked individuals were counted; finally, the total number of workers in the colony was calculated by the Lincoln Index formula. In order to determine the value of the Lincoln Index with the colonies studied, a shovel was utilized in excavating each of these nests, and every ant discovered was collected and etherized for a subsequent total count of each colony.

In 1968, the fourth summer of this project, temperatures were recorded, and soil moisture analyses were made as well. Temperatures were taken in the various quadrats three times daily, from July 29 through August 28, with a Fahrenheit telethermometer and thermistor. The latter was pushed into the soil as a probe to a depth of approximately two to three inches for subsurface temperature. The telethermometer was Model YSI, Serial No. 430, 3 scale, manufactured by the Yallow Springs, Ohio, Instrument Company. Temperatures were

also taken at the surface, six inches above the ground, and about five feet above the surface. The last temperature reading is equivalent to standard meteorological temperature.

I consulted Dr. J. W. Marr, of the University of Colorado Arctic and Alpine Institute, for procedures to follow in soil moisture analysis. The materials used were a trowel, one-half-pint standard soil-sample cans, a shovel, a double-beam trip balance, and an oven. Samples of soil were placed in the one-half-pint cans, and three samples were obtained daily from each of the three quadrats. A central representative area in each quadrat was selected for soil moisture samples, as well as for temperatures. The soil samples were all taken under rocks of medium size, and included topsoil from just beneath the rock to a depth of approximately five inches. The soil was weighed to one-tenth of a gram. The cans were oven dried at 110° C. for 24 hours, and the dry weight recorded. The results were expressed as per cent moisture per dry weight soil, and were computed by the following formula:

"Soil moisture as per cent = wet weight of soil - dry weight x 100 of dry weight of soil dry weight of soil

At the end of the third summer (summer of 1967), after all three quadrats had been meticulously examined for new as well as old colonies (previously discovered ones), a census was begun to determine the number of adult individuals in the various nests of ants, according to the different species. It was hoped that this procedure might help indicate which of the various ant species apparently were dominants in their respective quadrat areas. This census was largely conducted in the fourth summer of research, since there was only time for the census of a few nests in the latter part of August, 1967. The census entailed excavation of each nest, and the destruction of all ants in the nest in order to make a total count. For the census of the various ant colonies, the nest was excavated and emptied into a plastic bag, and a piece of cotton saturated with ethyl acetate was inserted into the bag, which was tied with string. After the field work, the contents were removed a small portion at a time and carefully examined for ants with the use of a magnifying glass for tiny formicids. The specimens were meticulously removed one at a time with forceps, counted, and inserted into 6-dram vials filled with alcohol.

۰.

RESULTS

The results of the collection of ants from nest sites for all three years, 1965 through 1967, in all quadrats, can be seen in Tables I - IX. The research revealed 25 species and subspecies of ants over the period of these three years. Table X presents a list of these species and subspecies with information on their ranges and zonation, which has been obtained from Gregg (1963). The known range of the formicids in North America is presented to permit comparisons between the ant's overall dispersal as a species and its distribution in my research area. The altitudinal ranges are only the known limits in Colorado. The zones in which the ants occur refer only to Colorado, and these were determined by Gregg from the elevational records, the habitat types, and the geographic localities. The altitudes of life-zones in northeastern Colorado with their elevations in feet are:

Zone	Elevation in feet
Alpine	11,500 - 14,000
Subalpine	10,000 - 11,500
Montane	8,000 - 10,000
Submontane	6,000 - 8,000
Upper Sonoran	4,500 - 6,000

Distribution of the various species and the numbers of their colonies in the three quadrats can be examined in Tables XI to XIII. Table XIV provides information relative to the composition of three ant subfamilies in the three quadrats for the years 1965 to 1967. This study attempts to focus closely on detailed aspects of ant distribution and endeavors to discover if there is any discernible boundary for ants of a sharp and narrow nature between the lower edge of the foothill zone and the plains, as there appears to be for plants. Ant species which are in the meadow and stop there, but do not penetrate the forest, and those which are at home in the ponderosa pines, but do not enter the meadow, should support the idea of an important faunal boundary. An examination of Tables XI to XIII show formicid species which do provide evidence of this boundary.

An examination of Tables XI - XIII, which are concerned with species composition for the three quadrats for 1965 through 1967, shows considerable differences in numbers of colonies for the various species for each of the three years during which ants were collected from nest sites. Since another objective of this study was to determine frequency and stability of ant colonies, examination of the tables reveals short duration and lack of permanence for numerous colonies.

The results of colony size estimation by the Lincoln Index are shown in Table XV. Four separate nests of the carpenter ant, <u>Camponotus (T.) vicinus</u>, and two nests of <u>Formica (N.) pallidefulva</u> were used.

Another purpose of the study is to relate climate to ant distribution, and Tables XVI - XVIII present the raw data for daily temperature readings from July 29 through August 28, 1968, for morning, noon, and afternoon temperatures in each of the quadrats. To facilitate analysis of the data, Tables XIX - XXI were prepared

to demonstrate weekly temperature averages in all quadrats for morning, noon, and afternoon temperatures. Tables XXII - XXIV illustrate weekly temperature maximums and minimums in all quadrats for morning, noon, and afternoon temperatures. Tables XXV - XXVIII present United States Department of Commerce, Weather Eureau records of climatological observations at Boulder. These tables include daily maximums, minimums, and temperatures taken at observation time, and point out precipitation for Boulder at the South Side Fire Station, 2225 Baseline Road, which is the Weather Bureau station nearest to my research area. Insofar as soil moisture is concerned, raw data are presented in Tables XXIX - XXXI for all three quadrats, and Table XXXII depicts the average daily and weekly per cent of moisture in all quadrats.

Another purpose of my study was to observe foraging activities of some ants. Since my son and I were more directly concerned with collection and analysis of distribution of the various species, this objective was, of necessity, a limited one; nevertheless, foraging activities were observed for four species of ants. Three of these formicids are members of the subfamily Formicinae, and one of these formicids belongs to the genus <u>Camponotus</u>; the other two are members of the genus <u>Formica</u>. The other species observed is a member of the subfamily Dolichoderinae and the genus <u>Iridomyrmex</u>.

I base my ideas for nest evacuation upon conditions noted in the various ant nests which might be responsible for the abandonment of the nests. Other factors which do not lend themselves to observation must also be considered, and these are noted in the discussion of this thesis. Plesiobiosis is a name given by Wheeler (1910) to cases in which two or more colonies of different species of ants establish their nests in contiguity, or very close proximity, as often happens under the same stone. This condition was noted in 12 species, almost half of the total number of formicid species discovered in my research sites. All three subfamilies are represented by species which had compound nests, and at least one compound nest was found for each species. Of all nests examined over the entire period of the study, there were a total of 36 compound nests. Usually two species were found under the same rock in plesiobiosis, but twice there were three species discovered in the compound nest.

An attempt was made in my research to observe nests which apparently branch from the original colony. Nests of six species of ants were observed which had probably formed in this way. The subfamilies Myrmicinae and Formicinae are represented, but no Dolichoderinae ants appeared to have polydomous nests in my research area.

One of the objectives of this study was to determine the dominants among ants in the area. Like most animals in the terrestrial area, these insects are not true ecological dominants, since they do not exert a major controlling influence on the community. Therefore, they are influents at best, since formicids do not, by themselves, modify the major factoral complex of a community in such a way that other organisms can live at the same place. However, some ant species by sheer numbers or by their aggressive nature may "dominate" others.

Tables XI - XIII give the number of colonies per species in the various quadrats for all three years, and Table XIV shows composition of species by subfamily. It becomes apparent at once that the Formicinae have the greatest number of species for every quadrat and in every year; the Myrmicinae are next, and the Dolichoderinae are third. Table XXXIII represents the ant colony sizes by actual total count, which were made in 1968, a year after sampling had been completed.

One of the objectives of this study was to examine and note those ant species in the research area which apparently have nuptial flights in late summer. In 1966 and 1967 the ants were collected in August, and in both years, numerous colonies with winged specimens were observed. In 1965, the first summer of this study, collecting was accomplished in June and early July, and winged specimens were not observed. Twelve species of ants which had colonies with winged specimens were seen during the observation period. All three subfamilies of ants are represented among these 12 species.

10

DISCUSSION

Distribution of the Various Species

There were ten species of ants discovered in the meadow in the three years encompassed by this study, and of these ten, three species, or 30% of the total, do not penetrate the ponderosa pine forest. In fact, three other forms which do penetrate the forest do so at a very considerable reduction in the number of colonies from those found in the meadow.

The three species which do not penetrate the forest are

Lasius alienus americanus Emery Formica obscuripes Forel <u>F. fusca argentea</u> Wheeler

The two Formica species do not penetrate the ecotone. Of the three species, <u>Lasius americanus</u> had six colonies in the meadow in 1965, two in 1966, and one in 1967. There were five in the ecotone in 1965, three in 1966, and two in 1967. <u>Formica obscuripes</u>, found only in the meadow, had one colony in 1965, and two each in 1966 and 1967. <u>Formica fusca argentea</u> had one colony in the meadow in 1965 and none in the ecotone. From these data it would appear that the lower edge of the Transition is an actual boundary, since distributional limits are encountered for the species of ants which are in the meadow and do not penetrate the forest. For <u>Lasius americanus</u>, the termination of its distribution is sharply outlined at the lower edge of the Transition. Furthermore, the two <u>Formica</u> species help to support the idea of a terminal at the Transition Zone's lower border since they are in the meadow and do not enter the forest.

There were 20 species of ants found in the ponderosa pine forest, and of these 20, 13 species, or 65%, do not enter the meadow. Three additional species which do penetrate the meadow undergo a very considerable reduction in colony number from forest to meadow. In fact, for two of these three species only one colony was found in the meadow during the three years of this study, as compared to 24 colonies in the forest for one species and 144 in the forest for the other.

The 13 species in the ponderosa pine forest which do not enter the meadow are

> Lasius niger neoniger Emery L. (A.) claviger coloradensis Wheeler L. (C.) brevicornis microps Wheeler L. niger sitkaensis Pergande Formica (N.) pallidefulva Latreille F. (P.) limata Wheeler F. (P.) lasioides Emery Camponotus (T.) vicinus Mayr Crematogaster lineolata (Say) Myrmica schencki emeryana Forel Leptothorax rugatulus Emery Iridomyrmex pruinosus analis (E. André) Liometopum occidentale luctuosum Wheeler

Five of the above species do not penetrate the ecotone. These are

11

Lasius (C.) brevicornis microps Wheeler L. niger sitkaensis Pergande Formica (P.) limata Wheeler Leptothorax rugatulus Emery Liometopum occidentale luctuosum Wheeler

In an analysis of the eight species in the forest which penetrate the ecotone but do not enter the meadow, it may be seen that <u>Lasius niger meoniger</u> had 44 colonies in the forest in 1965, 30 in

1966, and 29 in 1967, but in the ecotone there were only two of this species in 1965, one in 1966, and none in 1967. Lasius claviger coloradensis had one colony in the ecotone in 1965 and none in the other two years; however, there was one colony in the forest in 1967, but none had moved into the meadow. The formicid, Formica (N.) pallidefulva, in the ecotone had 32 colonies in 1965, 22 in 1966, and 21 in 1967; and there were six in the forest in 1965, four in 1966, and six in 1967. In spite of the considerable number of nests of this species in the ecotone, none were ever discovered in the meadow. This would appear to be further evidence for a faunal distributional border between the lower edge of the foothill zone and the plains. Insofar as Formica (P.) lasioides is concerned, the forest had one colony in each of the three years, and the ecotone had one in 1966 and 1967, but two in 1965; however, no colonies were found in the meadow. <u>Camponotus</u> (\underline{T}_{\cdot}) <u>vicinus</u> was represented in the forest by 15 colonies in 1965, 16 in 1966, and nine in 1967, but in the ecotone there were eight in 1965, and nine each in 1966 and 1967. Further evidence can be seen here of a distributional limit between the lower edge of the foothill zone and the plains, since no vicinus colonies whatsoever were located in the meadow. In 1965, six colonies of Crematogaster lineolata were found in the forest. One of these was located in 1966, and five in 1967; however, only one was discovered in the ecotone, and this was in 1967, but none of these ants moved into the meadow. Myrmica schencki emeryana had one colony in the forest in 1967, one in the ecotone in 1965, and three colonies in 1966, but the meadow did not have any nests of emeryana at any time. There was one colony of Iridomyrmex pruinosus analis in the

forest in 1967, and in the ecotone there were two colonies each in 1965 and 1966, and one in 1967; however, none was found in the meadow.

In the above descriptions, since in some cases the nests were located at the same places in all of the three years, we can probably assume that the same colonies were present. However, there appeared to be considerable instability of the nests in the study areas, and in spite of this lack of permanence, no ants of the above species migrated into the meadow from the lower edge of the foothill zone. Although of no importance insofar as evidence of a faunal boundary is concerned, it is of interest to note that there were two species confined to the ecotone. These illustrate one of the principles of ecology, whereby organisms are often found which are characteristic of, and often restricted to, the ecotone. In any event, these ants increase the faunal wealth of the region, and these two species are

Formica obscuriventris clivia Creighton Pheidole pilifera coloradensis Emery

From the information gleaned from the three quadrats relative to the ant species which do not enter the meadow from the forest and those which do not penetrate the forest from the meadow, it would appear that an important boundary exists for ants between the plains and foothills, which thus parallels the more conspicuous vegetational changes.

Of the three subfamilies represented by the species in my research areas, the Myrmicinae had six, or 24% of the total number of species, and these six species were also in six different genera.

<u>Crematogaster</u> <u>Myrmica</u> <u>Aphaenogaster</u> <u>Solenopsis</u> <u>Leptothorax</u> Pheidole

Of these six Myrmicine species, two were in only one quadrat (22.2% of the total number of single-quadrat species represented), two were in two quadrats (18.2% of all two-quadrat species), and two were in all quadrats (40% of the three-quadrat species).

The Formicinae were represented by three genera with 16 species which represents 64% of the total number of species discovered. One of the genera, Formica, had four species confined to one quadrat (44.4% of the total number of single-quadrat species), and four species found in two quadrats (36.4% of all the two-quadrat species). Another genus of the Formicinae, <u>Lasius</u>, had two species found in only one quadrat (22.2% of all single-quadrat species), three two-quadrat species (27.3% of all two-quadrat species), and two three-quadrat species (40% of the total number of three-quadrat species). The third genus of the Formicinae, <u>Camponotus</u>, had one two-quadrat species (9.1% of the total number of two-quadrat species).

The Dolichoderinae, or third sub-family, had three genera with three species (one species per genus). These three genera are <u>Iridomyrmex</u> with one two-quadrat species (9.1% of all two-quadrat species), <u>Liometopum</u> with one single-quadrat species (11.1% of all single-quadrat species), and <u>Tapinoma</u> with one three-quadrat species (20% of all three-quadrat species). Obviously, species found in all quadrats would be very tolerant ones and therefore widely distributed, with single-quadrat and species found in two quadrats progressively less adaptive, as well as having a much narrower distribution. In this study, two-quadrat species form the largest group with a total of 11; however, this group is followed very closely by the single-quadrat group with nine species.

It is of interest, further, to note that of the Myrmicinae and Formicinae, the former are twice as numerous in genera as the latter. In the three-quadrat group, there are the same number of myrmicine species as formicine species, but in the two-quadrat group, there are eight formicine species to only two of the myrmicine species, or four times as many two-quadrat formicine species. These findings confirm Gregg's similar observations in his <u>The Ants of Colorado</u> (1963) for the Formicinae and Myrmicinae, and perhaps, as Gregg suggests, The Formicinae are more advanced and geographically and ecologically more adjustable. In any case, this situation does appear to parallel evolutionary advance, as suggested by Gregg.

Gregg (1963) draws up lists of ants which, he writes, demonstrate the existence of a faunal boundary in the area of the Transition Zone. Only those ants which penetrate the Transition from the Sonoran or the Canadian, and whose upper and lower limits are determined by that zone, are included in the lists. He found 45 ants with their upper distributional limits somewhere in the Transition Zone, and 17 ants with their lower limits in the Transition, and from this information he was able to establish that the submontane is a boundary, or meeting place, between northern types of ants and those from southern sources and thus coincides with the vegetation pattern.

My study does not contradict the conclusion reached by Gregg relative to the existence of the boundary in the area of the

Transition Zone. Gregg's study was made on a state-wide basis with records from all over Colorado, whereas my research is concerned with the lower edge of the Transition Zone in a very limited area in which the quadrats were selected near to each other in different kinds of vegetation. My study thus is more closely focused on the minutiae of distribution in an attempt to discover if there is actually a discernible boundary of a sharper and narrower nature than the distributional terminal described in Gregg's book.

Stability and Frequency of Colonies

The classification of the 25 species and subspecies of ants found in my research is as follows:

Family Formicidae

Subfamily Myrmicinae Genera and Species Crematogaster Lund Crematogaster lineolata (Say) Myrmica Latreille -: Myrmica schencki emeryana Forel Aphaenogaster Mayr Aphaenogaster (A.) subterranea valida Wheeler Solenopsis Westwood Solenopsis (D.) molesta validiuscula Emery Leptothorax Mayr Leptothorax rugatulus Emery Pheidole Westwood Pheidole pilifera coloradensis Emery

Subfamily Formicinae Genera and Species <u>Camponotus</u> Mayr <u>Lasius</u> Fabricius <u>Lasius alienus americanus</u> Emery <u>Lasius niger neoniger Emery</u> <u>Lasius niger sitkaensis</u> Pergande Subgenus <u>Chthonolasius</u> <u>Lasius</u> (<u>C.</u>) <u>brevicornis microps</u> Wheeler <u>Lasius</u> (C.) <u>umbratus aphidicola</u> (Walsh) Subgenus Acanthamyops

Lasius (A.) claviger coloradensis Wheeler

Lasius (A.) latipes (Walsh)

Formica Linnaeus

Subgenus Proformica

Formica (P.) lasioides Emery

Formica (P.) limata Wheeler

Subgenus Neoformica

Formica (N.) pallidefulva Latreille

Formica (N.) pallidefulva nitidiventris Emery Subgenus Formica

Fusca group

Formica fusca Linnaeus

Formica fusca argentea Wheeler

Rufa group

Formica obscuripes Forel

Formica obscuriventris clivia Creighton

Subfamily Dolichoderinae Genera and Species <u>Iridomyrmex Mayr</u> <u>Iridomyrmex pruinosus analis</u> (E. André) <u>Liometopum Mayr</u> <u>Liometopum occidentals luctuosum</u> Wheeler <u>Tapinoma Förster</u> <u>Tapinoma sessile</u> (Say)

In the following descriptions and characteristics of the various species which were discovered in my quadrats, much of the material used has been obtained from Gregg (1963) and Creighton (1950). In describing the frequency of ants, I used Gregg's (1963) classification of abundance of ants in Colorado. This classification is based upon four categories, and placement in these categories depends upon the number of records. These categories are

Abundant Common	(over 100) (50-100)	Frequent
Uncommon Rare	(10-50) (1-10)	Infrequent

The subfamily Myrmicinae surpasses all other groups of ants in the extent of variation in morphology and habits. As has been noted, six species of this subfamily were represented in my quadrats, and their descriptions follow.

<u>Crematogaster lineolata</u> (Say) has a range which extends from eastern Canada through the North Atlantic States and North Central States to eastern Colorado, a southern extension follows the Appalachian Highlands to northern Georgia. This ant is very widespread in the northern and eastern United States stretching as far west as Colorado. Its subspecies <u>emeryana</u> replaces it here, and the two probably intergrade. Most species of <u>Crematogaster</u> nest under stones, in logs, or in standing timber, and many tend aphis and build carton sheds over them. Similar carton containers are often made by <u>lineolata</u> and used as brood chambers, and such incubators may be several yards away from the main nest; however, these carton brood chambers are abandoned at the beginning of the fall.

This species is considered to be rare in Colorado, for Gregg lists two records as of 1963. In my research, <u>Crematogaster lineolata</u> was not found in the meadow at any time, nor, for that matter, in the ecotone during 1965 and 1966; however, one nest was located there in 1967. In the wooded region, in 1965, this investigator discovered six colonies, one of which was present all three years, and was the only one found in 1966. In 1967, five nests were observed and sampled. From the above data, it can be concluded that <u>lineolata</u> is unstable in its nesting habits, and can also certainly be considered an infrequent species.

Myrmica schencki emeryana Forel has its range in Newfoundland to Georgia and west to the Rocky Mountains. As of 1963 there were 38 records of this species, which classified it as uncommon. This myrmicid is a subspecies under the European species schencki. It and other members of the genus Myrmica prefer to nest in the soil, and often use a covering object above the nest. As a rule, they are inoffensive ants. The distribution of Myrmica is most interesting, as it is the only large Holarctic genus which lacks xerophilous or subtropical representatives on this continent. A map showing the distribution of Myrmica in North America would show a widespread occurrence in Canada, with northern limits reaching Labrador in the east and Alaska in the west. Proceeding southward, one does not find it restricted to areas of moderate to considerable elevation in both eastern and western United States as it is now known from sea level. In the western mountains, the genus is abundant in subalpine and Canadian zones, in decreasing numbers in the Transition Zone and absent in Sonoran areas.

In my study none was found in the meadow, but some were discovered in the wooded area and in the ecotone. Although my research revealed few colonies of <u>Myrmica schencki emeryana</u>, more were found in the ecotone than in the other two quadrats. In the ecotone in 1965, only one was recorded, but in 1966 there were three, one of which was probably the same nest as in 1965, since it was at the same site. Only one nest was discovered in the forest, and that was in 1967, but no colonies were ever found in the meadow. This ant must obviously be considered infrequent. Although the data for this

insect are meager, it would appear that <u>Myrmica emeryana</u> is inherently unstable in its nest habits, since it apparently changes nest sites frequently.

The range of <u>Aphaenogaster (A.)</u> <u>subterranea valida</u> wheeler is the Rocky Mountain region from southern Colorado north to British. Columbia and west to the mountains of Utah. Gregg (1963) lists 72 records, and therefore, classifies <u>valida</u> as common. This subspecies is common in northern and central Colorado, but appears to diminish in southern Colorado. It holds sharply to the Transition Zone; it is absent from every zone both above and below this one. <u>Aphaenogaster valida</u> is a Rocky Mountain form of the species <u>subterranea</u>, and appears to nest under stones in moist, shady foothill canyons, although Creighton (1950) writes that it nest in dry and fully exposed situations. This investigator found, by far, the vast majority of records in the woodland, a fact which supports an observation of Gregg's, in that more ant records are found in forests.

In my research area, <u>Aphaenogaster valida</u> was an abundant ant, especially in the wooded unit, for in that quadrat, 52 nests were discovered in 1965, 48 in 1966, and 44 in 1967. Of those 52 nests located in 1965, only ten continued throughout the three years (18% stability); however, 12 others persisted into 1966 but no longer. It is perhaps of interest that in 1966 there were 23 new nests, of which 13 were also present in 1967 (57% stability for the period 1966-1967), and, in addition, there were 13 new nests in 1967. In 1965, I discovered eight colonies in the ecotone, two of which persevered throughout the study period, and two others extended as far as 1966 but not into 1967. In 1966, the ecotone yielded six nests,

two of which were new, and the same number (six) was present in 1967, one of which was new; however, in the meadow, only one colony was discovered, and that was in 1966. From the above information, it appears that <u>Aphaenogaster valida</u> is quite stable.

The range of <u>Solenopsis</u> (D.) <u>molesta validiuscula</u> Emery is Pacific Coast states eastward to Idaho, Colorado, and New Mexico. There were 56 records in Colorado in 1963, and occurrence is, therefore, common. This tiny, brownish-yellow ant is the common western form of <u>Solenopsis molesta</u>. It is quite common in the lower foothill zone, although it is rather abundant at various places on the plains and other Sonoran habitats. This species reaches higher elevations than <u>molesta</u>, the Eastern species, and Gregg (1963) believes it replaces <u>molesta</u> in mountainous western United States.

These insects, as well as some of their relatives, are known to be thief ants (lestobiosis). Their colonies are usually founded in close proximity to the nest passages of some larger species with which the tiny passages from the nest of the thief ant communicate. A steady pilfering of brood or other food from the nest of the larger species is carried on in such obscurity that the larger species rarely seems aware of its loss. These thief ants only occasionally forage above ground, and are almost impossible to see when they do so because of their minute size. Their stings are so small (few individuals exceed 2 millimeters in length) they have no effect on human skin, although they are very bad-tempered and pugnacious.

Solenopsis (D.) molesta validiuscula Emery was ubiquitous, insofar as habitations in all three quadrats is concerned, although

not as abundant as some species such as <u>Aphaenogaster</u> (<u>A.</u>) <u>sub-</u> <u>terranea valida</u> Wheeler. In the ecotone in 1965 seven sites were recorded, none of which was present in 1966; however, three of the same sites were occupied by <u>Solenopsis validiuscula</u> in 1967. The year 1966 yielded only one nest in the ecotone; 1967, on the other hand, yielded ten colonies in that quadrat. In the wooded region in 1965 two sites were occupied by <u>Solenopsis validiuscula</u>; in 1966 none were present; but in 1967 five were established. In the meadow 1965 and 1966 yielded eight colonies each, one of which persisted through to 1967; three others were existent in 1965 and 1966, but did not continue into 1967, and three colonies were found in the meadow in 1967. It would seem that <u>Solenopsis validiuscula</u>'s nest stability rates are low.

Leptothorax rugatulus Emery has the following range: Rocky Mountains, Sierra Nevada and Cascade Ranges, mountains of Arizona and Utah, and an eastern extension into the Black Hills of South Dakota. Gregg (1963) classified its occurrence as uncommon with 31 records. Greighton (1950) writes that <u>Leptothorax rugatulus</u> is widely distributed in the Transition Zone of the Rocky Mountains. Gregg (1963) found one site as high as 8700 feet in elevation in southwestern Colorado. The population in the California Sierras averages darker than that in the Rockies, but the full color range is present in both areas. This species shows many slight variations in sculpture, but thoracic rugae are always present.

Leptothorax rugatulus Emery was never located in the ecotone and meadow, but in 1965 the wooded quadrat yielded six nests, none of which were present in 1966; however, the latter year saw five new

records established in the forest. 1967 was the most abundant year, for 12 colonies were discovered, and two of these 12 colonies had persisted from 1966. From the above, it appears that <u>rugatulus</u> has very unstable nesting habits.

The last one of the six species of the subfamily Myrmicinae discovered in my study area is <u>Pheidole pilifera coloradensis</u> Emery which has the following range: northern New Mexico through Colorado to the Dakotas. Gregg (1963) indicates 53 records with a consequent common occurrence for this insect; however, it is not considered a truly common ant in the total fauna of Colorado. It is found in both the Upper Sonoran and the Transition Zones, and is, therefore, a generally more tolerant ant than most of the other species of its gerus. <u>Pheidole pilifera coloradensis</u> Emery is host to the parasitic ant <u>Epipheidole inquilina</u> wheeler, and Gregg, at the writing of his book, <u>The Ants of Colorado</u>, had not discovered any individuals of the parasite. I am sorry to say that I had no better fortune than Dr. Gregg in finding <u>inquilina</u>.

Seeds have been found in nests of <u>Pheidola coloradensis</u>, and two species of plants have been recognized; <u>Frysimum</u> sp. (wallflower) and <u>Chenopodium leptophyllum</u> (narrowleaf goosefoot). Most of the species garner seeds, and it is believed that the large-headed major workers function as seed huskers. Since the enlarged head of the major is mainly filled with mandibular muscles, this enables the jaws to exert considerable pressure, which should be useful in cracking off the husks of seeds; however, the ant will accept other food as well as seeds, such as animal tissue.

The colonies are usually small; however, they are generally larger than those of the typical eastern <u>pilifera</u>. <u>Pheidole coloradensis</u> nests in the soil, and prefers to nest in canyon bottoms and along the banks of streams. Browne (1958) and Byars (1936) found this thermophilic ant mainly on warm dry slopes, and Smith (1962) found it on the sunny slopes of both sides of Valmont Butte. Its nest may be built under a stone or in open soil without a covering object, but when no cover is present, there is often a mound or crater of excavated soil around the nest entrance. In Colorado <u>Pheidole coloradensis</u> is most often found at elevations between 5000 and 6000 feet, and is more abundant on the eastern slopes of the Rockies than to the west of them.

This insect was rare in my study, and was never discovered in the forest or meadow. In the ecotone one nest was located in 1965 which disappeared by 1966; however, another colony was recorded in 1966 which persisted until 1967. In the latter year only that one colony which continued from 1966 was present. On the basis of this meager evidence, it is difficult to assess this formicid's stability.

In the subfamily Formicinae the structure of the genera is not strongly variable, but the habits of the ants in the subfamily are highly diverse. As outlined in the classification, three genera with a total of 16 species were found in the quadrats.

<u>Camponotus</u> (<u>I.</u>) <u>vicinus</u> Mayr has its range as follows: South Dakota to Oklahoma and west through the Rocky Mountains to the Pacific Coast from British Columbia south into the highlands of Mexico, and Gregg lists 166 records as of 1963 with its occurrence

as abundant. This frequent formicid is represented in a broad selection of plant communities, and occurs in a great variety of geographic localities. It inhabits numerous vegetation types, both open and wooded, but it is almost never found in dense forest. It prefers clay soil, and in almost all instances, its nests are found baneath rocks, but occasionally the nests are found under bouldars, and it is exceptional to find <u>Camponotus vicinus</u> in decaying logs. It is usually an inhabitant of high plains, mesas, and mountain foothills, and appears to avoid the hottest, lowest, and most arid of the desert regions.

This formicid was not located in the meadow at any time. In the ecotone there was great stability of colonies of this ant, since of the eight sites at which <u>Camponotus vicinus</u> was discovered, in 1965, all were perpetuated throughout the three years, and in 1966 and 1967, a new colony was added for each year providing the ecotone, therefore, with nine colonies in each of those two years. The forest held 15 nests in 1965, of which only two were maintained throughout the research period, and the same quadrat provided 16 colonies in 1966; seven of these continued into 1967. There were nine colonies in the wooded region in 1967, only two of these were new nests previously not established. It would seem that, in general, although stability of this ant is not at a consistently high rate, in comparison with some other species, <u>Camponotus vicinus</u> is apparently quite stable.

Of the genus Lasius Fabricius, Creighton (1950) writes

"As far as is known, ell our species of <u>Lasius</u> tend root coccids and aphids. In the case of the strongly hypogaeic species of <u>Chthonolasius</u> it is thought that these insects subsist mainly

on the secretions of the coccids and aphids. Other species, particularly our two representatives of the subgenus <u>Lasius</u>, are more active in foraging above ground and supplement this diet with various foods. Most of the species of <u>Lasius</u> are remarkably flexible as to the types of nest sites which they will utilize, although most of them appear to prefer well drained soil that is not too dry. The nests may be free in the soil, under stones or other covering objects or in and under rotten logs and stumps."

The range of <u>Lasius alienus americanus</u> Emery is southern Canada and all of the United States except southern Florida, Texas, and arid sections of the Southwest. Wilson (1955) regarded it as holarctic in distribution, and Creighton writes of its higher incidence in the East and Central States than in the West. There were, up to 1963, 148 records of this ant in Colorado, which classifies it as abundant. The distribution of this formicid is wide, since it is found in all the life zones in Colorado except the Alpine; it does, however, extend to over 10,000 feet in elevation. Gregg's records from the Upper Sonoran Zone predominate, with those from the Transition a very close second.

Ecologically, <u>Lasius americanus</u> tolerates a wide variety of habitats and microhabitats, but Gregg never found any in rotting logs, although this type of nest site is often used in the Eastern and Midwestern states. None of my records were found in rotting logs either. The moisture requirements of this ant probably force it to depend on supplies of water in the soil under moisture-conserving objects, such as stones. Smith (1962) writes that though this ant is often found building small craters in grassy areas, colonies were found only under rocks in the Valmont area. I also found this ant only under rocks in my region. Insofar as my research study is concerned, <u>Lasius alienus</u> <u>americanus</u> Emery was not present at all in the forest in any of the three years of this study. In 1965 in the ecotone there were five colonies, none of which extended through 1966 and into 1967; however, two of these five colonies persisted into 1966, and three nests were located in 1966. In 1967, there were two nests, one of which was also at the same location in 1965, but no ants were discovered there in 1966. Six <u>Lasius americanus</u> colonies were found in the meadow in 1965, and none of these persevered into 1966 and 1967; however, one did continue into 1966. In the year 1966 the meadow yielded two colonies, whereas in 1967 the meadow offered only one. These records would seem, therefore, to indicate that <u>Lasius</u> <u>americanus</u> shows little nest stability.

Lasius niger neoniger Emery has the following range: southern Canada and eastern United States, except Florida and the Gulf Coast, west to the Rocky Mountains. Gregg classifies this ant as abundant in Colorado with 133 records by 1963. Mrs. R. Gregg, according to Gregg (1963), showed that <u>neoniger's</u> habitat preferences are in open and especially sandy areas, and it nests in the soil or under covering objects (sticks, rocks, and so forth). Its common competitor is <u>Lasius alienus americanus</u> Emery. An open substratum free of the accumulation of organic debris is also believed to favor the occurrence of <u>neoniger</u>, and is a limiting factor in its distribution. Wilson (1955) emphasizes that this species has most of its nests in exposed locations such as fields, meadows, grassy roadsides, trails, and sandy blowouts. Gregg writes that this is especially true of the East, but that in the western states, <u>neoniger</u>

is often found in forests and areas other than those cited by Wilson. This is certainly borne out by my study, since I have by far a large majority of species in the wooded region.

Gregg found one case in which neoniger was about to become host to Lasius (A.) murphyi Forel. A dealated female of the latter species was seen entering a <u>neoniger</u> nest under a piece of dung. Solenopsis (D.) salina Wheeler was taken on a different occasion in a lestobiotic relationship (presumably) with neoniger. Wheeler (1917) gives a very complete account of the manner in which the female of subumbratus behaves as a temporary social parasite when founding its colony in the host nest of neoniger. The intruding female at first attempts to make "friends" with the neoniger workers, but is usually repulsed by them. The neoniger workers may seize the appendages of the intruder, but they do not press the attack and soon release her. The subumbratus female then hides in the neoniger nest, and appropriates a part of the brood, over which she crouches until she has secured the neoniger nest odor. She is, thereafter, accepted without any further trouble by the neoniger workers. Nothing definite is known as to what becomes of the meoniger queen, but it seems certain that she is eliminated, probably by the intruding female.

Lasius niger neoniger Emery was common in my research area, especially in the wooded area, but it was never found in the meadow. In 1965 at the ecotone two sites revealed this species of ant, and the year 1966 produced only one colony, and this one was at a new nest site, but no colony was discovered in 1967 at the ecotone. The wooded quadrat was a very favorable area for <u>neoniger</u>, since 44 nests
were located in 1965, and of this total, ten persevered throughout the duration of this project (23% stability). Another five were on hand both in 1965 and 1966, but had disappeared in 1967. In the second year of this study (1966) 30 colonies were located, ten, as previously mentioned, were stable throughout the research, nine others which were newly discovered in 1966 were also located at the same sites in 1967. In 1967 29 nests were recorded, six of these were entirely new. This formicid appears to have considerable nest stability, at least more so than some of the other species discovered in this study.

The range of my next species, Lasius niger sitkaensis Pergande, is transcontinental; it extends from Nova Scotia across the northern United States and southern Canada to the Pacific Coast, north to southern Alaska, and south through the Rocky Mountains. Great Basin, the mountains of California, and an isolated population in the Black Mountains, North Carolina. It is classified as abundant in Colorado with 164 records by 1963. This formicid is a geographical race of Lasius niger Mayr, and is primarily a western and moderately boreal form; however, the other race-neoniger is an eastern form which extends quite far south. Colorado, in the midst of their overlap, has an abundance of each subspecies and a wealth of intergrades as well. This ant is much more abundant in higher elevations and in boreal vegetation types than is neoniger. This investigator collected a much larger number of species of meoniger than sitkaensis. Gregg has one record of it serving as host to Lasius (C.) subumbratus Viereck, one record of a nest containing larvae of the syrphid fly Microdon, and one case in which the thief

ant <u>Solenopsis</u> (<u>D.</u>) <u>truncorum</u> Forel was found inhabiting the walls of a <u>Lasius</u> nest, but in no case did I find any association between <u>Lasius sitkaensis</u> and any other ant. Creighton does not believe <u>sitkaensis</u> is a northern race and <u>neoniger</u> a southern one, but Gregg's data certainly support the contention that both are geographical races. They are connected by innumerable intermediate stages or intergrades and their ranges overlap widely.

There were no colonies of <u>Lasius niger sitkaensis</u> Pergands in the ecotone and meadow for any of the three years spanned by this research. In 1965 the forest relinquished four colonies, one of which was present all three years (25% stability). In 1966 four nests were found, none of which, with exception of the station recorded in 1965, persisted into 1967. The forest relinquished two colonies in 1967, one of which was at a completely new site. This formicid apparently shows considerable instability in its nesting habits.

My next two species of <u>Lasius</u> ants belong to the subgenus <u>Cthonolasius</u>. The first one, <u>Lasius</u> (<u>C.</u>) <u>brevicornis microps</u> Wheeler had ll records by 1963, and was, of course, considered uncommon. Its range is Alberta south to Colorado and westward into California. <u>Lasius</u> (<u>C.</u>) <u>brevicornis</u> Emery is primarily an eastern species, but it does reach into the Rocky Mountains. Some western species, but it depart somewhat from the typical eastern form, and perhaps are intergrades between <u>brevicornis</u> and its subspecies <u>microps</u>, and Colorado is probably an area of extensive interbreeding. This formicid is represented in moist forests, but can also penetrate into warm and dry situations, of which it is more tolerant than the

typical <u>brevicornis</u>. However, in my research, all of the records were established in the wooded area. It regularly takes advantage of nesting sites which save moisture and enable the ant to escape the direct effects of hot, dry weather.

Lasius (C.) brevicornis microps Wheeler had no stations in the ecotone or meadow in any of the three years encompassed by this study; however, in 1965 the forest provided this investigator with samples from 25 colonies. Nine of these colonies continued on through 1967 for a stability of 32%, and there were 21 colonies in 1966 including those discovered in 1965. Four of the 21 were also present in 1965, but were not existent in 1967. In 1967 a total of 22 nests were located, which figure, of course, includes the original nine from 1965. Five of the colonies located in 1967 had been discovered in 1966 and persisted to 1967. Although this species does appear to carry through from one year to the next in a number of cases, in general, it is a valid conclusion that the majority of the nests of <u>Lasius microps</u> lacked stability.

The other species of <u>Lasius</u> belonging to the subgenus <u>Chthonolasius</u> is <u>Lasius</u> (C.) <u>umbratus aphidicola</u> (Walsh). This formicid has the following range Nova Scotia and New Brunswick south to the Gulf States, and westward through the United States to the Rocky Mountains. Its occurrence is classified as uncommon in Colorado, since there were 16 records as of 1963. This ant generally prefers fairly moist and cool situations, nesting along the borders of woods, near streams, in moist meadow soil, and very often under stones where moisture is conserved. In eastern woodlands, it can be seen in rotting logs, but Gregg (1963) writes that although in

Colorado it has not been seen in rotting logs, this may be due to the rather small number of records, and this investigator did not find any in rotting logs either. Gregg, on one occasion, found <u>Leptothorax rugatulus</u> Emery associated with this species in what was at least a plesiobiotic relationship.

In the ecotone during 1965 six nests of <u>Lasius</u> (C.) <u>umbratus</u> <u>aphidicola</u> (Walsh) were discovered, only one of which was also occupied in 1966 and 1967 (17% stability); 1966 offered only that one nest, and 1967 only one other in addition. In the wooded quadrat three nests were located in 1965, one of which was at the same site all three summers (33% stability). In 1966 two additional nests were found, and in 1967 six sites were occupied, one of which was the nest present all three years. The meadow had the largest number, since in 1965 there were 12 records; however, only two remained all three years (17% stability); two others persisted until 1966. There were six nests in 1966 and eight in 1967. From the above, it would appear that <u>Lasius aphidicola</u> is weak in nest stability.

The last two species of <u>Lasius</u> in my research area are both in the subgenus <u>Acanthomyops</u>. Without exception, the ants in this subgenus lead a largely subterranean existence, since the nests are often built under stones or logs, or at the base of old stumps, and infrequently, they are built in the soil with no covering object. During the marriage flight the workers come out of the nest in large numbers, but this is the only time when they are easy to find above ground, for most of the time they are underground tending root aphids and coccids. All <u>Acanthomoyops</u> species have a characteristic

odor of lemon verbena, which Wheeler considers to be nest odor, but Creighton believes it is only produced when the ants are disturbed, and is more of a repugnatorial device. Because of the peculiar structure of the females of some of the species, it is generally assumed that these species, and probably others as well, are temporary social parasites.

The first of the two <u>Acanthomyops</u> species in my quadrats is <u>Lasius (A.) claviger coloradensis</u> Wheeler. Its range is Colorado and northern New Mexico, and since there were ten Colorado records by 1963, it was classified as rare in occurrence. This ant is a subspecies of <u>Lasius (A.) claviger</u> (Roger), which is an eastern and midwestern species, and <u>Lasius claviger</u> is seldom found here, probably due to insufficient moisture. It overlaps with <u>Lasius colora-</u> <u>densis</u> in Colorado, but <u>coloradensis</u> is more tolerant of drier habitats, although it will nest in moist sites when these are available. Creighton (1950) writes that the nests of <u>coloradensis</u> are often located on open intermountain plateaus where the only cover is that furnished by sage-brush bushes.

<u>Iasius (A.) claviger coloradensis</u> Wheeler was found in the ecotone in 1965 at one site, but not again in 1966 or 1967. One nest was discovered in the forest, but only in 1967, and no nest was located in the meadow. There is too little information available here to comment on nest stability for this species.

The other <u>Acanthomyops</u> species in my study is <u>Lasius</u> (<u>A.</u>) <u>latipes</u> (Walsh). The range of this insect is coast to coast in the northern United States with southern extensions to South Carolina, in the Rocky Hountains to New Mexico, to the Sierras in Celifornia,

and north to Alaska. There were 41 records by 1963, and it, therefore, is considered uncommon. This species is the most successful species of the Acanthomyops group in Colorado, judging by the abundance of locations, from which Gregg has reported it. It is tolerant of widely ecological conditions which probably accounts for its broad dispersal, and it often invades rather dry situations, where it is usually found under large rocks or boulders. This formicid is a temporary social parasite on Lasius (subgenus Lasius), and Gregg found one case in which a mixed colony consisted of Lasius latipes and its host Lasius alienus americanus Emery. A case of probable plesiobiosis was discovered between latipes and Myrmica sabuleti americana Weber, and Solenopsis (D.) molesta validiuscula Emery was also present under the same boulder. Gregg observed a nuptial flight of latipes after a shower at 5 P.M. on August 6, 1955; it is noteworthy that most of Gregg's records for reproductives of this species are also in August.

Lasius (A.) latipes (Walsh) was not located in the ecotone until 1967, and then, only at one site which had been occupied by Lasius alienus americanus Emery in 1966. The wooded region yielded four colonies in 1965, only one of which persevered for all three years. In 1966 this particular colony was the only one present, and in 1967 there was one other besides the nest which persisted for the duration of this study. Twenty-five per cent stability was apparent here for the wooded quadrat. In the meadow in 1965 three nests were discovered none of which were located again, but two were noted in 1966, one of which was also present in 1967. In fact, in 1967 that single nest was the only one recorded. This species was obviously rare in my study, and also had poor nest stability.

In the genus <u>Formica</u> eight species of ants were discovered. This genus has many outstanding characteristics. It is the largest genus of ants in America north of Mexico, and its members make up about one-sixth of the entire ant fauna. A very large number of forms in this genus are endemic to the United States and Canada. Behavior is also quite variable in this genus, for there is slavemaking, various kinds of temporary social parasitism, and several distinct types of nest construction. Also, this genus affords excellent opportunity for zoogeographical studies.

The eight species of ants which were recorded from my study plots are in three subgenera, one of which is the subgenus <u>Proformica</u>. Two of the three species in this subgenus were discovered. Each of the three has nesting habits identical to the other two. The colonies are generally small, and the nests are usually constructed in soil beneath stones or other covering objects. Often these ants are regarded as timid, because they are preyed upon by species of the <u>sanguinea</u> group, but they do not lack in pugnacity and will defend their nest if disturbed. It is probably because of their small size, rather than their lack of courage, which makes it possible for <u>sanguinea</u> species to enslave them.

One of the two species of <u>Proformica</u> discovered in my research is <u>Formica</u> (P.) <u>lasioides</u> Emery. Its range is as follows: coast to coast in southern Canada and northern United States with a southern extension into the mountains of California, in the Rocky Mountains to New Mexico, and in the Appalachian Highlands. The occurrence is common in Colorado, since 79 records were established by 1963. This ant is a slave ant for <u>Formica</u> (<u>R.</u>) <u>wheeleri</u> Creighton.

Formica (P.) lasioides Emery was never found in the meadow, but in the ecotone there was one colony which persevered throughout the study, and this was the only colony discovered in both 1966 and 1967; in the ecotone, however, in 1965 there was an additional colony present. The wooded quadrat had one colony for each of the three years, and this nest was apparently the same for 1965 and 1966, but had disappeared by 1967; however, in that year another nest of Formica lasioides was located. Even on the basis of this meager evidence, it appears that nest stability in this species is quite high.

The other species of the subgenus <u>Proformica</u> which was recorded ed is <u>Formica</u> (P.) <u>limata</u> wheeler. This is an ant which has as its range the southern Rocky Mountains in Colorado and New Mexico, and the mountains of eastern Utah; it is also found in North Dakota and northern Minnesota. It is classified as uncommon on the basis of 20 records by 1963. Although Colorado is in or near the probable center of its range, this ant is not at all abundant. It seems to be fairly tolerant of dry conditions. This ant has a strongly shining surface and an almost hairless thorax.

Formica (P.) limata Wheeler was extremely rare in my research area; it was never found in the ecotone or meadow; it was located in the wooded quadrat in 1967, and then in a compound nest with <u>Tapi-</u> <u>noma sessile</u> (Say).

The second of the three subgenera of the genus Formica is <u>Neoformica</u>. The members of this group build rather obscure nests, and the colonies are relatively small in size. The nests are usually under stones or at the base of tufts of grass, and since these ants are very timid, they will generally make no effort to defend the nest. They often abandon the brood, although they often sneak back and try to rescue it later. Because they do not battle with other ants, they are easy victims of slave-making species. There were two species of this subgenus discovered in my study.

Formica (N.) pallidefulva Latreille has the following range: Virginia to Florida and west to Texas and Kansas, with sporadic occurrences northward along the east coast, in the middle west, and in Colorado. Gregg classified it as rare in Colorado based upon 4 records as of 1963. Creighton describes this insect as austral with a broad distribution in the southern and eastern United States. The stations from which Gregg (1963) secured pallidefulva were at the base of the foothills along the eastern mountain front, and here summer temperatures are high, and winters comparatively less severe. In my research area, with one exception, this ant's nests were beneath stones. The exception was a flattened can which apparently served its purpose well, since that particular site (ecotone nest number 15) was occupied by probably the same colony for the entire three years encompassed by this study. Incidentally, insofar as collection of samples of this species was concerned, I had to work very quickly in order to collect enough for an adequate sample. This insect is victimized by thief ants, as I discovered it in four lestobiotic associations with Solenopsis (D.) molesta validiuscula

Emery. In one of the four stations, three colonies were under the same rock, and the third species was <u>Tapinoma</u> <u>sessile</u> (Say).

Formica (N.) pallidefulva Latreille was common in my study area, for in 1965 in the ecotome 32 colonies were located, and of these, five (16%) persisted for all three years. Four more were present also in 1966, but were no longer there in 1967; however, in 1966 in the ecotome 22 nests were recorded, and in 1967 there were 21. In the forest six nests were discovered in 1965, but only four in 1966 and six in 1967. Two colonies were at the same site from 1965 to 1966, but all others were new locations. No colonies were ever found in the meadow. It is interesting that in three cases, colonies discovered in 1965 had no ants in 1966, but had the same sites in 1967. In view of the above, it can be seen that this formicid is quite unstable in its nesting habits.

The second species of the subgenus <u>Neoformica</u> discovered in my study is <u>Formica</u> (N.) <u>pallidefulva nitidiventris</u> Emery. Its range is southern Quebec and Ontario to the mountains of northern Georgia, and west to the Dakotas, Wyoming, Colorado, and northern New Mexico. Its occurrence is uncommon, based on the collection records as of 1963 which were 47. This subspecies replaces the typical <u>pallidefulva</u> over the northern half of the United States east of the Rocky Mountains, and in the Rocky Mountain states themselves, it is by far the most common member of the subgenus <u>Neoformica</u> (Gregg, 1963), but this was not the case in my study area, since <u>Formica pallidefulva</u> was much more numerous. It was formerly believed to occupy only the eastern foothills, but there is now abundant evidence to show that it occupies a broad territory in Colorado, which includes low elevations on the western slope. The majority of its records are from the lower foothills. As mentioned previously for <u>Formica pallidefulva</u>, it was also true for <u>Formica nitidiventris</u> that, since these ants are so timid, in collecting them, I had to move quickly in order to obtain a minimum sample.

Formica (N.) pallidefulva nitidiventris Emery was not located in the ecotone; however, in the wooded quadrat two nests were found in 1965 which did not have any ants in 1966. A third colony was discovered in 1967 at a different nest site. In the meadow one colony was brought to light in 1965, and this one endured all three years at that site. In 1966 another colony was found, but this one was no longer present in 1967; however, in the latter year, two new colonies were located. From these results, one can but conclude that considerable instability existed, insofar as this species is concerned, for only one colony was present all three years.

The subgenus <u>Formica</u> was represented by four species of ants, two of which are in the <u>fusca</u> group, and two in the <u>rufa</u> group, and the latter group includes mound-building species, which embrace many variants. The nests are very often thatched in the <u>rufa</u> group, and many members of that group are temporary social parasites. In the <u>fusca</u> group the species show a preference for nesting in the soil. They are ubiquitous and also very timid (ideal for slave-making).

Of the Formica fusca group, two species were located. One of these is Formica fusca Linnaeus, which is the very common "black ant" met in varying environments in prairies, foothills, and mountains. Its range is Holarctic; in North America from Newfoundland west to

Alaska and the entire northern half of the United States with southern extensions along the major mountain ranges into North Carolina and Tennessee, New Mexico and Arizona, and into California. It is classified as abundant on the basis of a total of 277 Colorado records as of 1963. This ant is probably the best known ant in the Northern Hemisphere, and it is absent from few major habitats, but prefers to nest in the soil. It is certainly among the most common of Northern ants, but in semi-arid regions, its occurrence tapers off sharply, and it is also limited by the upper timberline; for Gregg (1963) never found any established nests in the alpine tundra. Also, he never found any <u>Formica fusca</u> in the plains grassland habitat.

In the meadow area I only found one record of this species and only in 1967, and none were found in the ecotome. The nests, except for the one in the meadow, were all located in the wooded region, and these were often found near the base of trees.

Formica fusca has broad elevational tolerances and appears to be abundant at high altitudes and extends somewhat higher than argentea. The members of this species lack pugnacity, for they are usually regarded as cowardly and timid, but docility may be a better term than cowardly, since the individuals which have been brought into the nests of slave-making species acquire all the pugnacity of their captors. In any case, they appear to be ideal subjects for slavemaking. They are also ubiquitous, which is another advantage for the slave makers.

As stated above, Formica fusca Linnaeus was not discovered in the ecotone at any time; in fact, it was not located in the meadow

in 1965 nor 1966; however, one nest was observed in 1967. The forest in 1965 held nine colonies, one of which was extended throughout the three years (11% stability), and another continued into 1966, but was terminated by 1967. In 1966, seven nests were present, and in 1967 eight. From the above information, it is probably safe to conclude that <u>Formica fusca</u> is very unstable, insofar as nesting habits are concerned.

The other species of the fusca group found in my study area was Formica fusca argentea Wheeler. Its range is New England and the midwestern United States, and all of western United States from New Mexico, Arizona, and California to Montana and Washington with an extension into British Columbia, but is not found over the entire Holarctic or even Nearctic range of the typical fusca. According to Gregg (1963) geographically argentea is much commoner in western United States, and this area would seen to be its distributional center. Fragmentation of the range of fusca by the late glaciation may have led to isolation of a portion of it in the western and southwestern United States, giving a chance for accumulation of traits found in argentea. Now there is extensive hybridization between the two. Although argentea appears to extend far to the east, it is much more abundant in the west, and although its range is now engulfed by fusca, it would appear to have originated in the western states. It is classified as common in Colorado, since the number of records as of 1963 was 77. This species has broad elevational tolerances, and its altitudinal range is also like fusca. The former is more abundant at lower elevations and appears to extend a little lower than fusca, as it seems to be at home on the plains,

especially the short-grass prairie. Gregg once found this species living in close association under the same rock with a colony of <u>Lasius (A.) murphyi</u> Forel. Like <u>fusca</u>, these ants also are timid, and are brought into the nests of slave-making species. <u>Formica</u> <u>fusca argentea</u> Wheeler was rare in my study. It was only discovered at one site in 1965 in the meadow.

As in the case of the fusca group of the subgenus Formica, the rufa group also produced two species in my research area. One of these is Formica obscurices Forel, and the range of this ant is northern Indiana and Michigan westward across the northern United States and southern Canada to Oregon and British Columbia, with a southern extension through Utah and Colorado to northern New Maxico; Lake Tahoe, California. Its classification is abundant based on 119 Colorado records as of 1963. This species is probably the commonest thatching ant of the western states, and is most prevalent between 5000 and 8000 feet. It builds large, dome-shaped nests composed of thick masses of coarse plant debris (most often twigs), and would appear to inhabit a variety of vegetation types; however, Smith (1962) found one small nest beneath a rock. The finished mound does not depend upon any support for the detritus, but it is believed to be begun around the base of a small plant. Gregg's data corresponds with these descriptions, and the colony I found, which was present all three years, also gives support to the above information. The ant appears to prefer clay soil, but a wide variety of general habitats is used by this species, and it is one of our most eurokous formicids. Cole (1932) states that the colony always contains two or more queens, and that winged males and females are

present in large numbers through June and July. Weber (1935) showed that a typical marriage flight does not occur in <u>Formica obscuripes</u>; instead, reproductives come from the nest singly, or in small groups during a period of several weeks. <u>Formica obscuripes</u> Forel workers also appear to attend aphids on various flowering herbs. I found this ant to be very aggressive. The best known mound maker of the <u>rufa</u> group is <u>Formica obscuripes</u>, and its nests often form a conspicuous feature of the landscape in the west. There appears to be controversy over whether <u>obscuripes</u> females are temporary social parasites. Wheeler thought most, if not all, species of the <u>rufa</u> group were probably temporary social parasites, but Creighton does not believe this applies to <u>obscuripes</u> as well as some other species of <u>rufa</u>.

Two colonies of <u>Formica obscuripes</u> were recorded, and both of these were found in the meadow. One colony persisted all three years, but the other was found in 1966, (<u>Lasius</u> (<u>A.</u>) <u>latipes</u> (Walsh) occupied the nest site in 1965) and persisted in 1967. The latter nest was smaller than the former; however, the nest which had been discovered in 1965 was much smaller by 1967, for it appeared to have been damaged by some object, and was much reduced in size. The stability of the nests of this insect must be classified, on the basis of these two colonies, as excellent.

The other species of the <u>rufa</u> group of the subgenus <u>Formica</u> is <u>Formica</u> obscuriventris <u>clivia</u> Creighton. Its range is Iowa, Wisconsin, and Manitoba west to British Columbia, with a southern extension in the Rocky Mountains to Colorado, Utah, and New Mexico. It is classified as uncommon based on 47 records by 1963. The

typical subspecies of this ant is an eastern and midwestern insect, and is extremely rare and erratic in this region. The great majority of records Gregg has seen belong to the western subspecies <u>clivia</u>. Intergradation and hybridization of the latter with the former occurs from Minnesota to Illinois, but the region may be larger, and hybridization probably occurs in Colorado as well, due to the range overlap here. This formicid is most abundant in Transition Zone communities, with a sharp drop above and below these levels, but it is known from four zones.

Gregg found the greater number under rocks and most frequently in rich, loamy soil. I discovered one nest in the ecotone every year at the same station, which presumably was the same colony. The nest stability for this species based on this one nest each year is 100%, but, of course, if more than one colony had been found, this stability might have been less than 100%. This evidence is much too meager to establish a generalization on nest stability.

The third subfamily represented in my quadrats is the Dolichoderinae which has representatives which are rather uniform in both habits and structure. They prefer to nest in soil and show little evidence of distary specialization, and because of the latter, they have become serious household pests. Three genera with one species each were discovered in my collecting sites.

One of the three species located is <u>Iridomyrmex pruinosus</u> <u>analis</u> (E. André), which has the following range: California east to Texas, Oklahoma, and Kansas; north to southern Idaho. Its occurrence is uncommon based on 36 records established by 1963. This ant appears to enjoy hot, dry localities; however, it is not absent from certain rather moist situations also, such as canyon bottom forests. It will nest under covering objects, but is often found with nest openings exposed, surrounded by craters of discharged soil. The ant is difficult to collect, since it moves very quickly, especially in bright sunlight, which markedly increases the temperature of the soil. These insects are entomophagous, and all nest in soil.

<u>Iridomyrmex pruinosus analis</u> (E. André) was discovered in my research area at three different stations. Two of these were in the ecotone, and were found in 1965, and one of these colonies persisted through all three years, but the other was not found in 1967. The third colony was discovered in the wooded quadrat. Therefore, only one third of the nests had stability throughout the study period.

The second of the three Dolichoderinae species is <u>liometopum</u> occidentale <u>luctuosum</u> wheeler. Its range is southern Wyoming to New Mexico and Arizona, and the mountains of California; it is rare in Utah and Nevada. Based on 29 records established by 1963, it is considered uncommon. Wheeler (1917) considered <u>Liometopum luctuosum</u> to be quite rare, more so than <u>Liometopum apiculatum</u>, but as far as its presence in Colorado is concerned, just the opposite is true. Wheeler also wrote that <u>luctuosum</u> is associated with pine trees (or at least conifers), and has its nests under the roots of these trees. Gregg (1963) writes that it is frequently associated with evergreen vegetation, but it is by no means limited to that type. All the records I have of this species are in the wooded area which is, of course, Ponderosa pine.

Creighton points out that the ants in this genus are more agressive than those of most of the Dolichoderinae genera. These

are very pugnacious insects, which forage in files and attack fiercely if disturbed. They have a secretion with a powerful and disagreeable odor like that of butyric acid and spray this on intruders. The nests are usually built under stones or in hollow trees, and the nest chambers are sometimes subdivided by a mass of paperlike material which they manufacture by mixing bits of soil and vegetable detritus with a secretion which hardens and causes the mass to become solid. The North American species of <u>Liometopum</u> will tend aphids and coccids. This ant will feed upon any insects it can capture. According to Creighton, it has a preference for nests at higher levels (4000-7000 feet) than <u>Liometopum occidentale</u>, and therefore, keeps its range largely separate from the latter. This is rather difficult to understand, since both <u>luctuosum</u> and <u>occidentale</u> are foothill ants.

For <u>Liometopum occidentale luctuosum</u> Wheeler no records were established in the ecotone or meadow; however, in the forest in 1965 seven sites were located; one of these persevered for all three years. In fact, in 1966 only three nests were discovered, and in 1967 only the nest which extended through all three summers. Of the seven colonies, then, which were present in 1965, only one nest (14%) was stable for the duration of this research study; another 14% was still present in 1966, but absent in 1967, therefore, stability of nests in this species was poor.

The third and last of the Dolichoderinae species is <u>Tapinoma</u> <u>sessile</u> (Say). Its range is southern Canada and the entire United States except the southwestern deserts, and it can certainly be classified as abundant, based on 199 records established by 1963.

In view of its broad range, it can be considered ubiquitous, for this ant is obviously extremely adaptable and tolerant, but it is absent from tundra and in exposed areas in forests near timberline. Also, there are none of these ants in saltbush deserts of Colorado. Grogg observed plesiobiotic relations in this ant on two occasions, for in one case, Tapinoma was living under the same rock in close proximity to a colony of Lasius (A.) murphyi Forel, and in the other, with a colony of Lasius (A.) claviger (Roger). Creighton writes that Dr. M. R. Smith's observations of Tapinoma sessile are that it is not at all particular about its nest sites, which it changes frequently, and will nest in the soil, with or without a covering object, under bark, and in all sorts of preformed cavities, and it also becomes a pest in houses. Furthermore, it has a remarkable elevational tolerance, since it occurs from sea level to sub-alpine areas. The workers usually forage in files and are ownivorous, although they appear to prefer honey-dew and sweet foods when they can get them. Tapinoma is an emergetic ant; it is not timid but less combative and bad tempered than many ants. This is why on occasion it forms compound nests with other ants. The Tapinoma odor is due to butyric acid.

Insofar as <u>Tapinoma sessile</u> (Say) is concerned, in my study area it proved ubiquitous, since it was found in all three quadrats; however, it was by far most abundant in the wooded region. The ecotone had six nests in 1965, but these were vacated in 1966 and 1967; however, one colony was discovered for each of the latter two years. The forest in 1965 yielded 19 nests, none of which continued throughout the three summers. In fact, only two were found at the same

sites in 1966, but in that year three new nests were located. The wooded quadrat in 1967 rendered eight nest sites none of which were occupied by Tapinoma sessile in 1965 or 1966, for most of these had been occupied by other species in 1966; only two were new nest sites. Even before one considers the meadow, it is obvious that great nest instability exists in Tapinoma sessile. It is of interest that in 1965, of the six sites which housed sessile, not a single one of these six sites had any species of ant in 1966. This condition was a very common occurrence in most species in my study, in that where nests were occupied by a particular species in one year, very often the next year when that species was not present, no other ant was present at that site either. Tables I - IX demonstrate this time and again. In 1965 there were two nests in the meadow; in 1966 there were two located at different sites, and in 1967 there was only one colony which was discovered at a new site.

From the descriptions of the results of the collections of the ants in my quadrats, over the period of three years, from 1965 through 1967, it is quite obvious that the majority of ants in my study were quite unstable insofar as their nesting habits are concerned.

Creighton writes that <u>Tapinoma sessile</u> (Say) changes its nest sites frequently. Wheeler (1910) writes of the migration of <u>Formica sanguinia</u> which often has summer and winter residences, and uses the expression "analagous to the city and country homes of wealthy people." The summer nests are built in open, sunny locations where food is plentiful and conditions are most favorable for rearing the brood, whereas, the winter nests are built in secluded

spots in the woods and are used as hibernacula or, very rarely, for protection from the excessive heat of summer. Wheeler says that the migration of ants from one nest to another is determined upon, and initiated by, a few workers, which are either more sensitive to adverse conditions, or of a more alert and venturesome nature, than the majority of their fellows. After they select a site, these workers begin to deposit their brood, queen, males, fellow workers, and even their myrmecophiles.

Two types of ant nests are distinguished according to Wheeler (1910)—the temporary and permanent, without any corresponding differences in architecture. Wheeler compares this to Forel's distinction of monodomous and polydomous colonies. The nest of the former colony is a single circumscribed unit, whereas a polydomous colony spreads over several nests, the inhabitants of which remain in communication with one another and may visit back and forth.

Brian (1965) claims that most mortality of nests occurs before they are even started, since it depends upon the survival of the queen. Most causes of nest loss are still unknown. Brian writes that a favorite method of studying ant colony survival is to mark and measure their mounds. In species that construct mounds, those mounds which disappeared probably did so through shading, through human interference, or from unknown causes. Talbot (1961) noticed that many mounds of a species of <u>Formica ulkei</u> in Michigan were formed and then abandoned. These mounds may not have necessarily died--they may have returned to their parent colony, and so, perhaps, represent trial buds that proved unsuccessful. Scherba (1961, 1963) studied the population dynamics of <u>Formica opaciventris</u>,

which reproduces by budding following pleometrosis, and between 1957 and 1959 small mounds had as high as 35% mortality. The large mounds are destroyed by destructive forces, shading, exhaustion of local resources, predation, and other causes still unknown. More data for many other species are needed to determine the causes of nest mortality. Trampling and cultivation destroys many ant nests and severely restricts the areas that are inhabitable.

Although Pickles (1937 and 1938) described changes of ant nest distribution, and Brian (1952) worked on the replacement of nests under stones, Yasuno (1965) writes that there is no quantitative investigation on the change of the ant population. To analyze the population dynamics of the ant population, a consecutive study was accomplished at the Kayano grassland on Mt. Hakkôda, which is situated in the northern part of the main island of Japan, during the period 1957-1961. Yasuno found that the ant population is in a dynamic state. The number of nest-mounds of <u>Formica truncorum</u> <u>yessensis</u> change seasonally, but the number of nests of this species and <u>Camponotus herculeanus japonicus</u> seems to be constant. As for <u>Formica fusca japonica</u>, the number of nests is also nearly constant. A stable state is maintained apparently in the number of nests, but stability of individual nests is very low, especially in <u>F. fusca</u> <u>japonica</u>.

Lincoln Index

Another purpose of this study was to determine the value of the Lincoln Index in determining the number of individuals in an ant colony. Table XV presents the results of this work, and with exception of one colony of <u>Camponotus</u>, it can be seen that the density determined by actual count is greater than that figure derived from the use of the Lincoln Index. In other words, the use of the index in my study resulted in great underestimation of the size of the colony.

Andrewartha and Birch (1954) write that it is important that either the initial marking or the subsequent catching be done evenly over the area selected for study, because of the following assumptions which the index requires: a. the marked individuals redistribute themselves at random to the unmarked ones; b. the marked ones are neither more nor less readily caught than the unmarked ones; and c. between the times of release and recapture there have been no gains or losses by births, deaths, or migration.

Golley and Gentry (1964) used the mark-recapture procedure with the southern harvester ant, <u>Pogenomyrmex badius</u>, and the technique produced contradictory results. These workers found that the use of the index depended on the following assumptions: a. all normal workers forage, b. marked and unmarked ants mix in the hill before the second sample is withdrawn, and c. the radioactive phosphorus which was used for labeling the ants, adheres to the tagged ants until the sample is taken, yet is not transferred to other ants within the hill. The tagging and excavation experiment was unsuccessful for this species, since the radioactive phosphorus spread to other members of the colony. Also, tagged ants were only in the upper part of the hill, with the majority in the uppermost 10 centimeters, and this suggests that the ants do not mix within the hill (at least for short periods of time) and that all the

workers do not participate in foraging activity. It has long been known that ant workers, like honey bees, remain in the nest as nurses for a time after hatching from the egg and begin foraging as they age. Incidentally, Golley and Centry found that movements of ant hills is a common phenomenon in these ants. The authors use the term movements of ant hills to mean migration from old to new hills. The results of Golley and Gentry's work aupport Chew (1960), who worked with <u>Pogonomyrmex occidentalis</u> (Cresson) in Arizona. Chew, using similar methods, found that only about one half of the workers were active at the surface and that random mixing within the colony did not take place. I consulted Dr. Chew at the University of Southern California, Los Angeles, and it is Chew's opinion that since many workers do not forage, the Lincoln Index underestimates the size of the colony, and therefore the index may be useful in estimating the density of foraging workers.

Climatic Conditions

Brian (1965) found that in woodlands the colony densities of various ants are remarkably high, and figure VIII and table XII show that the wooded area had not only the largest number of nests per year, but also the greatest species diversity as well. This quadrat was in a ponderosa pine forest facing north by northwest and is a very favorable location for ants. The meadow also faced north by northwest, but upon examination of figure IX and table XIII it can readily be seen that there is a distinct paucity of ants here in comparison with the forest. The ecotone faced east by southeast but mostly east. This quadrat had a considerable number of nests and species each year but not to the extent of the wooded quadrat. According to Wheeler (1910), colonies of ants in hilly or mountainous country are usually more abundant on the east and south exposures. My study proves to be an exception to the above, but as Gregg (1963) writes, general observations do appear to support Wheeler, although the whole question needs further study. Gregg also found that ants which are confined to tree-covered environments or which use both arborescent and open environments form the bulk of the ant fauna of Colorado. This statement is certainly supported by my findings. He adds that grassland and desert occupy the more level and more arid regions, and their ants contribute a much smaller proportion of the total fauna.

As noted previously, temperatures were taken two to three inches below the surface of the ground, at the surface, six inches above it, and five feet from the ground. As stated by Gregg (1963), in spite of the importance of atmospheric heat to all forms of life, the temperature of the soil is of particular significance to myriads of organisms, because so many are in direct contact with it. The invertebrates in soil faunas, of which ants form a conspicuous part, are subject to soil temperatures virtually throughout their lives, except for short periods like the nuptial flights of ants. Gregg believes it is logical to consider surface temperature as properly a phase of soil temperature, since it is a measure of heat generated by contact of light with the soil. Further, surface temperatures may reach critical upper limits of tolerance for minute organisms before either the air or the internal soil temperatures, and the soil surface is the place where so many of these organisms carry on

a large percentage of their activities. Hence, its importance should not be ignored. The air temperature immediately above the surface also has an important influence on the activities of these small terrestrial animals, and consequently these measurements were included in my work. Air temperatures show greater fluctuation than soil temperatures.

Even a cursory examination of the tables showing raw data reveals a distinct variation from one quadrat to the other for each of the four different types of temperature recording areas. As noted previously, tables XIX, XX, and XXI show weekly morning, noon, and afternoon temperature averages in all quadrats for the various localities at which temperatures were measured. In the morning, the woodland subsurface temperatures consistently averaged lower than the meadow temperatures but higher than those in the ecotone. The noon and afternoon subsurface temperatures showed the forest intermediate between the ecotone and the meadow.

Insofar as the weekly morning surface temperature averages are concerned, in three of the five weeks (60%) the forest temperatures averaged below those of the meadow and the ecotone. According to Chauvin (1967), forest insects do not receive anything like the same radiation as those which live out in the open air, since the forest floor receives only a low percentage of the sun's radiation. The forest, however, was intermediate between the ecotone and meadow in averages of my weekly noon surface temperatures. The morning temperature means for the station six inches above the ground were lower in the forest than in the ecotone and meadow for

three weeks of the five, and at noon and in the afternoon the forest temperatures were intermediate between the ecotone and the meadow.

The meadow consistently had the highest temperatures of all three strata--subsoil, surface, and six inches above the surface. Radiant energy from the sun is a source of heat for insects, and both visible light and infrared rays will serve to heat the body above the surroundings. The radiant energy can cause the temperature to go high enough to cause the insect to retreat to a shady spot such as the nest and therefore curtail its activities. This is probably an important factor in limiting the number of colonies in the meadow, since insects may find themselves beyond their optimum temperature due to the sun's rays reaching the soil. Weekly averages for temperatures taken five feet from the ground revealed similarities to the other temperature-recording stations. It should be remembered that the stratum five feet from the ground should not have as great an influence on the ant as the other levels, but for ants which forage on plants at that height there would be an influence.

The ponderosa pine forest's subsurface weekly morning temperature averages were higher than those of the soil's surface in the forest in four of the five weeks. In the ecotone this was the case in only two of the five weeks, and in the meadow the situation was like that of the forest. The deeper soil apparently retains some heat from the previous day, whereas the surface has cooled considerably during the night. The morning soil warmth would probably stimulate activity in the adult ants and growth in the young. Weekly temperature averages six inches above the ground in the morning were generally higher than the surface temperatures in every week and in every quadrat, but one week saw the same temperatures at both recording stations in the forest, and one week in the meadow the surface temperature was higher than that of the stratum six inches above the ground.

At noon subsurface weekly temperature averages in all quadrats were lower than at the other three recording stations. In the meadow the surface temperatures were higher than those taken six inches above its surface in three out of five weeks (one week they were the same). This fact would limit foraging activities if temperatures were above the optimum. In a comparison of the noon temperatures five feet above the ground with the surface reading, the meadow had higher weekly average temperatures in every case at the surface. In the forest only one week saw the surface temperature warmer than the temperature taken five feet above the ground, and in the ecotone the surface was consistently cooler than the temperature five feet above the ground. This observation is another possible reason for a greater number of ant colonies in the forest and ecotone.

In the afternoon the forest had a higher weekly average surface temperature as compared to its subsurface reading in three of the five weeks, but the surface temperature in the meadow was not higher than the subsurface in any week. The ecotone had two out of five weeks of surface temperature averages higher than subsurface. This fact would appear to indicate that the forest has a more constant daily temperature which permits a longer foraging and activity

period. Chauvin (1967) says that during the day, the underwood having quickly reached its equilibrium, this condition will remain with remarkable consistency throughout the day. During the night the temperature stays uniform all through the forest with two zones of minimum temperature, one on the floor where heavier cold air sinks, and the other in the treetops. This is in agreement with my results, since the majority of the five weeks revealed lower surface temperatures in the wooded area, in the mornings, than in the other two quadrats.

Since weekly maximum and minimum temperatures may be of even greater significance than weekly temperature averages as noted proviously, tables XXII, XXIII, and XXIV were prepared. These show weekly morning, noon, and afternoon temperature extremes in all quadrats at each of the four temperature recording locations. Insofar as temperature extremes were concerned, there was a much greater range from maximum to minimum temperatures from the highest station (5' above ground) to the lowest station (subsurface) and in that order.

Means were calculated for the weekly maximum and minimum morning subsurface temperatures. The results for average maximum temperatures are: ecotone--63.6, forest--64.8, and meadow--66.5; average minimum temperatures: ecotone--59.0, forest--60.4, and meadow--61.8. The forest was intermediate between the ecotone and the meadow, and the forest appears to have morning subsurface temperatures which are less variable than those in the other quadrats. The means of the weekly maximum surface temperatures in the morning are; ecotone--67.5, forest--68.6, and meadow--68.1. It is apparent that

the forest maximum surface temperature is generally higher than in the ecotone or meadow, which makes possible a longer daily period of activity for the ants in this quadrat. However, the slight temperature differences may or may not be significant. The ants usually are exposed to a higher morning temperature, and may, therefore, begin foraging and other activities earlier in the day.

The lowest weekly morning temperature recorded for any quadrat at the three stations in or near the ground during the recording period was for the ecotone (46.5°) . The forest was intermediate between the ecotone and meadow in the means of the weekly temperature minimums for the two stations below and at the ground's surface. For stations at, below, or six inches above the ground, the highest weekly morning temperature recorded was in the forest (80.9°) . The ecotone had 80.5° as its maximum, and the meadow had $.78.3^{\circ}$. This temperature in the forest was at six inches above the surface, and this would probably indicate an earlier start in activity for the ants in the forest, that is, if the temperature differences are significant. Four-tenths of 1° between the scotone and forest is not a large difference, although the difference between the forest's 80.9° and the meadow's 78.3° is greater.

It can be seen in table XXIII, which shows weekly noon temperature maximums and minimums, that the forest was intermediate between the meadow, which generally had higher maximums and minimums than the forest, and the ecotone, which had the lowest maximums and minimums.

Means were calculated from the weekly figures for the noon temperature extremes. These are

Subsurface

Maximum

ecotone-64.6
forest-68.3
meadow73.0

Minimum

ecotone-60.1 forest--62.4 meadow--66.5

Surface

Maximum

ecotone-80.3 forest--84.5 meadow--87.0

Minimum

ecotone-63.6 forest--65.1 meadow--69.6

Six Inches Above the Surface Maximum

> ecotone-81.1 forest--85.3 meadow--87.2

Minimum

ecotone-64.4 forest--66.4 meadow--67.4

It is apparent from these figures that the forest has a middle posi-

From table XXIV we can see that for afternoon weekly temperature extremes at both the surface and six inches above the ground the wooded quadrat often had higher maximums, but minimums were between those for the ecotone and the meadow.

Means from the weekly figures were calculated for the afternoon temperature extremes. These are

Subsurface

Maximum

```
ecotona-66.6
forest--71.9
meadow--75.0
```

Minimum

ecotone-61.2 forest--64.7 meadow--67.3

Surface

1. A. I. A.

Maximum

ecotone-77.5 forest--81.2 meadow--81.1

Minimum

ecotona-61.2 forest--63.8 meadow--66.4

Six Inches Above the Surface Maximum ecotons-73.6

forest--80.5 meadow-81.4

Minimum

ecotone-60.9
forest61.1
meadow-65.2

It may be noted that in the afternoon on the surface the mean temperature maximum for the forest was 31.2° - higher than the ecotone, and slightly higher than the meadow; this observation may serve as evidence for a longer daily temperature range for foraging and other activities. Minimum mean temperatures for the woodland were consistently between those for the meadow and the ecotone. Since much of the activity of ants occurs on the soil and a few inches above, it would appear that temperature is a very important factor in the Mooded quadrat, since it is very favorable there much of the time. One can conclude from these tables that there is considerable variability between the three samples taken daily from each quadrat. Upon examination of table XXXII, which shows the average daily and weekly per cent of moisture in all quadrats in the first three of the four weeks, it may be seen that the average per cent moisture was highest in the meadow, but it was highest in the ecotone the last week, and in the forest the per cent moisture was higher than in the ecotone for one of the four weeks. Based upon the four weekly figures for all quadrats, the monthly mean percentage of moisture is as follows: ecotone 11.52, forest 10.51, and meadow 12.76. Obviously, the woodland had a smaller percentage of moisture during this period. Perhaps the soil moisture or humidity is more ideal in the forest, since it is true that the rate of development in some organisms is retarded at high humidities. Also, it is probable that excess moisture may cause mortality in terrestrial insects due to fungus and other related pathological agents.

According to Gregg, moisture is perhaps less limiting than temperature; however, the meadow may be an example of a case of excessive moisture. Certainly, the moisture and temperature combination provides suitable conditions for ant colony formation in the ecotone, but these two factors may interact in the wooded area more advantageously than in the ecotone, for there were so many more ants in the forest, and also more species. However, other factors may also be influential in the wooded quadrat, such as soil, vegetation, and food availability. For instance, according to Chauvin (1967) the rain which reaches the soil after leaching the leaves on the trees has been changed chemically and in particular has become richer in mineral salts. It may contain 4 to 20 times as much calcium and 10 to 50 times as much potassium as the rain that falls on open ground. This helps to modify the litter of dead leaves, and the insects living there cannot fail to be affected by it. This would have only an indirect effect upon ants, however.

Foraging Activities

Four species of ants were observed insofar as their foraging activities are concerned, and Camponotus (T.) vicinus Mayr ant workers were often observed climbing trees in their search for food. Formica fusca Linnaeus workers were seen foraging on and under the bark of the ponderosa pines. In my observations of Formica obscuripes Forel, I found that it probably forages for considerable distances. Only one nest of this species was discovered in 1965, and in the quadrat in which the nest was located, the meadow, foragers were seen as far away as approximately 75 meters from the nest. Columns of formicines of this species were seen moving along trails carrying food and thatching materials, and the columns were followed and traced to the mound nests. Since in 1965 only one colony of this species was located in the meadow, it is very probable that the foragers which were followed originated from the observed nest. Creighton (1950) writes that Iridomyrmex pruinosus analis (E. Andre) are very active in spite of their rather small size, and they forage in files. This investigator observed these foraging files on two occasions. The ants appear to follow a scent path produced by a pheromone, as in one case I drew a finger across their path. which caused great confusion.

Nest Evacuation

There may be many reasons for ant nest evacuation. In my own research area, it was observed that often wherever ant nests had been vacated, a white mold was seen under the stone where the nest had formerly been. Possibly secretions of the mold may be undesirable to ants. Then, too, quite frequently where nests had been vacated a large spider would be found under the rock, and since spiders are predaceous, they may have been a major cause for departure from a nest. I always tried to be careful in sampling the nests, but this practice may have been responsible for the ants vacating the nest too. Flooding may have eliminated some nests, since run-off in the spring is sometimes excessive. Microclimatic conditions, food sources, and ant territorialism must certainly be considered in evaluation of causes of nest abandonment.

Compound Nests

~ :-

The ants in these compound nests were found under the same stone, but their colonies were separate (plesiobiosis, which has been defined in the Results section). Of the subfamily Myrmicinae, <u>Myrmica schencki emeryana</u> was one of the 12 species of ants which had at least one compound nest. I found two compound nests for <u>Myrmica emeryana</u> in which were associated <u>emeryana</u> and <u>Lasius niger</u> <u>neoniger</u> Emery in one case, and <u>emeryana</u> and <u>Formica (N.) pallidefulva</u> in the other. <u>Aphaenogaster (A.) subterranea valida</u> wheeler was found near another species of ant under the same rock upon six different occasions. The ants with which <u>valida</u> was associated in close proximity were <u>Tapinoma sessile</u> (Say), <u>Lasius (G.) brevicornis</u> <u>microps</u> Wheeler (two occasions), <u>Formica fusca Linnaeus (two occa</u> sions), and <u>Lasius (C.) umbratus aphidicola</u> (Walsh). <u>Solenopsis (D.)</u> <u>molesta validiuscula</u> Emery was found associated with other species on four separate occasions. In three cases, <u>Formica (N.) pallide-</u> <u>fulva Latreille</u>, and <u>Solenopsis validiuscula</u> were under the same rock, and in another instance, there were three species in plesiobiosis; these were <u>Formica pallidefulva</u>, <u>Solenopsis validiuscula</u>, and <u>Tapinoma sessile</u>. The last of the myrmicine ants having compound nests was <u>Leptothorax rugatulus</u> Emery with one nest in which it was closely associated with <u>Lasius (C.) brevicornis microps</u> Wheeler.

Of the subfamily Formicinae, Camponotus (T.) vicinus Mayr on one occasion was found associated with another species under the same rock, the other species being Lasius (A.) latipes (Walsh). Although Gregg states that Lasius alienus americanus Emery has been found living in plesiobiosis with Formica (P.) neogagates Emery, I did not find any cases where Lasius Americanus lived in a compound nest with any other ant. Another formicine Lasius niger neoniger Emery was associated with another species under the same rock in two cases. These species are Myrmica schencki emeryana Forel and Lasius (C.) brevicornis microps Wheeler. I found Lasius (C.) brevicornis microps wheeler in six different compound nests of two species per site in association with Lasius niger neoniger Emery, Tapinoma sessile (Say), Formica fusca Linnaeus, Leptothorax rugatulus Emery, and Aphaenogaster (A.) subterranea valida Wheeler (two records). Another member of the subfamily Formicinae, Lasius (C.) umbratus aphidicola (Walsh) was found by Gregg (1963) on one occasion
associated with Leptothorax rugatulus Enery in at least a plesiobiotic relationship. In 1965 in the ecotone I found Lasius aphidicola under a rock associated with Formica (N.) pallidefulva Latreille. when the rock was removed, the two species did not appear to interfere with each other in any way. Gregg cites a nest of mixed Formica (R.) wheeleri Creighton and Formica (P.) neogagates Emery with a quantity of aphidicola which were all dead; presumably the aphidicola served as food for the other ants. Three members of the genus Formica of this subfamily (Formicinae) revealed at least one compound nest. Formica (P.) limata Wheeler was located in the forest in 1967 and in a compound nest with Tapinoma sessile (Say). Formica (N.) pallidefulva Latreille was found in apparent plesiobiosis with these species: Lasius (C.) umbratus aphidicola (Walsh), Myrmica schencki emeryana Forel, and Tapinoma sessile (Say). For the third member of the genus Formica, Formica fusca Linnaeus, Gregg recorded one instance of plesiobiosis and this was with Myrmica lobicornis lobifrons Pergande under a rock. I discovered two cases; one species with which Formica fusca was associated being Lasius (C.) brevicornis microps Wheeler, and the other species Aphaenogaster (A.) subterranea valida Wheeler.

Of the subfamily Dolichoderinae, one species, <u>Tapinoma</u> <u>sessile</u> living with at least one other species under the same rock on seven different occasions. The species associated with <u>Tapinoma</u> <u>sessile</u> were <u>Formica</u> (N.) <u>pallidefulva</u> Latreille, <u>Aphaenogaster</u> (A.) <u>subterranea valida</u> Wheeler (two records), <u>Lasius</u> (C.) <u>brevicornis</u> <u>microps</u> Wheeler, <u>Formica</u> (P.) <u>limata</u> Wheeler, and <u>Solenopsis</u> (D.) <u>molesta validiuscula</u> Emery. In one case, <u>pallidefulva</u>, <u>validiuscula</u>, and <u>sessile</u> were all under the same rock. Gregg also found a case where <u>Tapinoma</u> was a third member of three colonies under the same rock and <u>Solenopsis</u> (<u>D.</u>) <u>molesta validiuscula</u> Emery, which was also present, may have been lestobiotic (thieving) on the other two ants. In my records of three species in close proximity, <u>Solenopsis</u> may have borne the same relationship to the other two ants.

Polydomous Nests

Of the six species of ants which had probably formed as branches from the original colony, there were five belonging to the subfamily Formicinae, and the other species was a member of the Myrmicinae. The myrmicine ant is Aphaenogaster (A.) subterranea valida Wheeler, and I observed polydomous nests for valida on three occasions in the wooded quadrat. Of the Formicinae, in at least one instance, several nests of <u>Carponotus</u> (T.) vicinus found in the forest were very close to each other, and workers could be seen visiting other nests (polydomous situation). I observed Lasius niger meoniger Emery living in polydomous nests on two occasions in the wooded quadrat, and another species of Lasius, Lasius (C.) brevicornis microps Wheeler living in one polydomous nest in the same quadrat. Of the two Formica species which were found in polydomous nests, Formica fusca Linnaeus was observed on two occasions in the forest, and Formica (N.) pallidefulva Latreille was seen in a polydomous nest condition on three occasions in the ecotone.

101

Census of Colonies

I planned to count at least one colony from each species, and three, if possible, so that an average might be calculated. Unfortunately, not all species were counted although the majority were tabulated. There were several reasons for not counting all species, one being that many nests simply were vacated. Also, since in 1967 no stakes were inserted beside the newly discovered nests, these were difficult to locate again, and then, too, a few older stakes were missing. When this occurred, it was not difficult to identify the previous year's nest site, because of the preparation of detailed maps (see figures VII to IX) which had numbers for the various stakes. These numbers were not included in the figures, since they would have made them too cluttered, and, also they were not necessary for the purposes of the figures. In some cases, more than three nests were counted for a particular species. This was due to the fact that taxonomic identification was not accomplished until after the field " work, and I sampled nests on the basis of previous records. Since a great deal of nest change and migration occurred, the result was seen in duplication of species counts. It is important to note that the actual counts of the colonies of the various ants would have been somewhat higher if all the foraging workers could have been included. However, since the absence of some workers occurred for all species, results will still be comparable to those which would have been achieved had all workers been counted.

In an examination of the tables for species composition (XI to XIII), it can be observed that in the ecotone in 1965 Formica (N.) pallidefulva Latreille was the dominant species insofar as colony numbers are concerned; however, colony-size average for this species was low at 76 members. Two other species, <u>Camponotus</u> (T_{\cdot}) vicinus Mayr and Aphaenogaster (A.) subterranea valida Wheeler, were both common in the quadrat and had higher individual colony population counts. The former had 100 individuals for average size, and the latter had 471. Therefore, all three of these species possibly exerted a dominant influence on associated ant colonies in the ecotone. Solenopsis (D.) molesta validiuscula Emery, Lasius (C.) umbratus aphidicola (Walsh), and Tapinoma sessile (Say) all were quite common in the quadrat, and the remaining ten species were rare and uncommon. This observation appears to confirm the wellknown ecological principle that there are more rare or uncommon species than there are abundant species and that a few very common species may be far greater in numbers of individuals. In the ecotone in 1966, similar results were found as in 1965 but Aphaenogaster valida did not have as large a proportion of nests in comparison with Formica pallidefulva and Camponotus vicinus as before. In the ecotone in 1967 pallidefulva still led with the greatest number of colonies, but Solenopsis (D.) molesta validiuscula Emery now surpassed vicinus, and colony size in validiuscula averaged 221 which, of course, exceeded vicinus.

In the wooded quadrat in 1965 <u>Aphaenogaster (A.) subterranea</u> <u>valida</u> Wheeler was probably the dominant species with 25.4% of the colonies. This ant's colony size averaged 471 individuals, which compounded whatever influence it may have exerted in the quadrat. <u>Lasius niger meoninger Emery</u> was probably also important as a dominant, as it had 21.5% of the colonies, and its colony size averaged 399. <u>Lasius (C.) brevicornis microps</u> Wheeler and <u>Tapinoma sessile</u> (Say) also were probably important in their dominance in the quadrat. In 1966 in the forest the relationships were like those in 1965 except that <u>Camponotus (T.) vicinus</u> Mayr perhaps was somewhat more important as a dominant than <u>Tapinoma</u>. In 1967 in the forest, with exception of the fact that <u>Leptothorax rugatulus</u> Emery (average colony size 231) might have taken over <u>vicinus's</u> place in ecological importance in the quadrat, the relationships were the same as for 1965 and 1966.

In 1965 in the meadow <u>Lasius</u> (C.) <u>umbratus aphidicola</u>(Walsh) was dominant with 35.3% of the colony total, and these colonies were large with an average of 1852 individuals per nest so their importance was probably considerable. <u>Solenopsis</u> (D.) <u>molesta validiuscula</u> Emery (average colony size 221) and <u>Lasius alienus americanus</u> Emery (nest size 43) both possibly were important with 23.5% and 17.6% of colony total respectively. In the meadow in 1966 <u>Solenopsis validiuscula</u> now became the dominant (32% of colonies) over <u>Lasius aphidicola</u> (24%), and <u>Lasius americanus</u>'s percentage dropped to eight per cent, a fact which caused it to be considered uncommon or rare. In 1967 in the meadow <u>aphidicola</u> again probably took over the leadership in dominance with 40% of the colonies, whereas <u>Formica</u> (N.) <u>pallidefulva nitidiventris</u> Emery and <u>Solenopsis validiuscula</u> tied with 15% each.

104

The species Formica obscuripes Forel in the meadow, <u>Cremato-gaster lineolata</u> (Say) in the forest, and <u>Liometopum occidentale</u> <u>luctuosum</u> Wheeler in the forest had low percentages of quadrat colony totals which would classify them as rare in the quadrat; however, the fact that they had the following average colony sizes would probably indicate that they had considerable ecological significance: <u>obscuripes-3,330, lineolata-3,481</u>, and <u>luctuosum-1,215</u>. Since <u>obscuripes</u> is also a larger ant and forages far afield, it should certainly be influential in its area.

As has already been noted, from one year to the next there appear to be changes in the numbers and ecological importance of various ants in their respective areas. It is of interest, therefore, to examine the 25 species and subspecies discovered in this study and determine the changes for these in terms of the nest numbers in 1965 and in 1967. The question is whether they remained constant in number. It has already been noted that most of the 25 species showed considerable individual nest instability.

Table XI shows species composition of the ecotone in all three years of the study, and of the 16 species found, two had the same number of nests in 1967 as in 1965. These are <u>Formica obscuriventris clivia</u> Creighton and <u>Pheidole pilifera coloradensis</u> Emery. Four increased in number: <u>Lasius (A.) latipes</u> (Walsh) and <u>Crematogaster lineolata</u> (Say) had no colonies in 1965 or 1966 but one each in 1967, while <u>Camponotus (T.) vicinus</u> Mayr and <u>Solenopsis (D.)</u> <u>molesta validiuscula</u> Wheeler also increased their colony numbers. The rest of the species declined in nest number from 1965 to 1967. but, of course, in some cases, as in <u>F</u>. <u>lasioides</u> and <u>Iridomyrmex</u>, the decline was only one colony.

Table XII shows species composition for the wooded quadrat, and in this table, which depicts a total of 20 species, two had the same nest total in 1965 as in 1967. These species are Formica (N.) <u>pallidefulva</u> Latreille and Formica (P.) <u>lasioides</u> Emery. There are five species which showed an increase in nest total for the quadrat. <u>Iridomyrmex pruinosus analis</u> (E. Andre'), Formica (P.) <u>limata Emery</u>, and <u>Myrmica schencki emeryana</u> Forel each had one colony in 1967 but no nests in 1965 or 1966, and <u>Solenopsis</u> (D.) <u>molesta validiuscula</u> Emery and <u>Leptothorax rugatulus</u> Emery also showed increases. All others showed decreases in nest numbers. In the forest, as in the ecotone, there are a number of species which have almost the same nest total in 1967 as in 1965 with a decline of only one, nest.

Table XIII shows species composition of the meadow, and this table depicts a total of 10 species, three of which increased their nest totals. These are Formica (N.) pallidefulva <u>nitidiventris</u> Emery, Formica obscuripes Forel, and Formica fusca Linnaeus. The latter had no nests in 1965 or 1966 and only one in 1967. The other quadrat species decreased, but again, as in the other two quadrats, there are several species which decreased their 1965 nest total by only one nest in 1967.

Nuptial Flights

Sudd (1967) states that in temperate lands the mating flight is usually late in the year, and if he refers to late summer as being late in the year, a considerable number of my species support his thesis. Although no such flights were actually seen by this observer, the results are based on collection and observation of winged males and females in the various ant colonies.

In the ecotone in 1966, the following species were noted:

Lasius (C.) umbratus aphidicola (Walsh) Aphaenogaster (A.) subterranea valida Wheeler Camponotus (T.) vicinus Mayr (three nests) Lasius alienus americanus Emery

In the wooded quadrat in 1966 the following species had winged members:

Aphaenogaster (A.) subterranea valida Wheeler (mine nests) Lasius (C.) brevicornis microps Wheeler (four nests) Lasius niger meoniger Emery (three nests) Camponotus (T.) vicinus Mayr (two nests)

In the meadow in 1966, there were the following species with winged ants:

Lasius (C.) umbratus aphidicola (Walsh) (three nests) Solenopsis (D.) molesta validiuscula Emery (four nests)

In the ecotone in 1967 the following species were observed:

<u>Iasius (C.)</u> <u>umbratus aphidicola</u> (Walsh) <u>Formica (N.) pallidefulva</u> Latreille (three nests) <u>Iasius alienus americanus</u> Emery <u>Crematogaster lineolata</u> (Say)

SUMMARY

There were ten objectives or purposes of this study listed in the introduction. This summary will briefly relate the outcome of the research in terms of these objectives.

1. Twenty-five species and subspecies of ants were discovered in the research area, which consisted of three quadrats in a meadow, a wooded area, and an ecotone. Of these 25 species, nine are species found in two contiguous quadrats, which provides evidence that the plains-foothill boundary is an important biological boundary. Two additional species in the meadow did not penetrate the ecotone, which may help support the thesis that ants provide evidence of a boundary at the lower edge of the Foothill Zone.

2. Individual colonies of most of the species of ants found in this study are not very stable.

3. The Lincoln Index is not a very effective tool in estimating the size of ant colonies. It results in an underestimation of the size of the colony, since the Lincoln Index presupposes that all workers forage.

4. Moisture and temperature have very important effects upon ants. Temperature is especially significant for fossorial species, which includes the majority of ants. Temperature and moisture in the wooded area were especially favorable for ants, since the greatest number of colonies, as well as the largest variety of species, were located there. 5. Foraging activities, which were quite variable, were noted for four species of ants.

6. Causes for exodus from an ant nest may be excessive moisture, which results in flocding and subsequent growth of mold, disturbance to nests, food shortage, and possibly other causes.

7. A considerable number of ant species, of the 25 discovered, live in plesiobiosis, or compound nest associations.

8. Polydomous nests were found in six species of formicids.

11

10

9. Probable dominants among ants were noted for the three quadrats for each of the three years.

10. A considerable number of ant species have nuptial flights in late summer.

LITERATURE CITED

- Allee, W. C., A. E. Emerson, O. Park, T. Park and K. P. Schmidt. 1949. Principles of animal ecology. W. B. Saunders Co., Philadelphia, Pa. 837 pp.
- Andrewartha, H. G. and L. C. Birch. 1954. The distribution and abundance of animals. University of Chicago Press, Chicago, Illinois. 782 pp.
- Armin, L. C. 1963. A study of the family Carabidae (Coleoptera) in Boulder County, Colorado. Doctoral thesis. University of Colorado. 465 pp.
- Barrett, K. E. J. and J. C. Felton. 1965. The distribution of the wood ant, Formica rufa Linnasus (Hymenoptera:Formicidae), in southeast England. Entomologist 8:181-191.
- Borchert, H. F. 1956. The ant fauna of foothill canyons in the vicinity of Boulder, Colorado. Master's thesis. University of Colorado. 57 pp.
- Brian, M. V. 1952. The structure of a dense natural ant population. J. Anim. Ecol. 21:12-24.

1965. Social insect populations. Academic Press, New York. 135 pp.

Brian, M. V., J. Hibble and D. J. Stradling. 1965. Ant pattern and density in a southern English heath. J. Anim. Ecol. 34:545-555.

1

- Browne, J. T. 1958. A distributional study of the ant fauna of Gregory Canyon, Boulder, Colorado. Master's thesis. University of Colorado. 91 pp.
- Byars, L. F. 1936. An ecological study of the ants of the plains region of Boulder County, Colorado. Master's thesis. University of Colorado. 52 pp.
- Chauvin, Rémy. 1967. The world of an insect. McGraw-Hill Book Co., New York. 256 pp.

Chew, R. M. 1959. Estimation of ant colony size by the Lincoln Index Method. J. New York Entomol. Soc. 67:157-161.

1960. Note on colony size and activity in <u>Pogonomyrmex</u> occidentalis (Cresson). J. New York Entomol. Soc. 68:81-82.

- Cole, A. C. 1932. The thatching ant, Formica obscuriventris Forel Psyche 39:30-33.
- Creighton, W. S. 1950. The ants of North America. Bull. Mus. Comp. Zool. Harvard University. Vol. 104:585 pp.
- Delfin, E. D. 1954. An entomological survey of Gregory Canyon, Boulder, Colorado. Master's thesis. University of Colorado. 61 pp.
- Elton, C. S. 1932. Territory among wood ants (Formica rufa L.) at Picket Hill. J. Anim. Ecol. 1:69-76.
- Forel, A. H. 1928. The social world of the ants compared with that of man. Putnam, New York. 2 Vols.
- Francoeur, André. 1966. La faune myrmécologique de l'érablière à sucre (<u>Acertum saccarophori</u>, Dansereau) de la région de Québec. Naturaliste Can. 93:443-472.
- Golley, F. B. and J. B. Gentry. 1964. Bioenergetics of the southern harvester ant, <u>Pogonomyrmex badius</u>. Ecology 45(2):217-225.
- Gösswald, K. and W. Kloft. 1961. Beziehungen einiger im Walde lebender Mäuse zu Waldameisennestern. Waldhygiene 4:22-23.
- Gregg, R. E. 1947. Altitudinal indicators among the Formicidae. Univ. Colo. Stud., Ser. D, 2:385-403.

______ 1963. The ants of Colorado. University of Colorado Press, Boulder, Colorado. 792 pp.

Jackson, C. H. N. 1936. Some new methods in the study of <u>Glossina</u> <u>morsitans</u>. Proc. Zool. Soc. London. 811-896.

ī :

- 1939. The analysis of an animal population. J. Anim. Ecol. 8:238-246.
- Knight, C. B. 1965. Easic concepts of ecology. Macmillan Co., New York. 468 pp.
- Lincoln, F. C. 1930. Calculating waterfowl abundance on the basis of banding returns. U. S. Dept. Agric. Circ. No. 118.
- Merriam, C. H. 1898. Life zones and crop zones of the United States. Bull. U. S. Dept. Agric., No. 10, 79 pp.
- Messenger, P. S. 1959. Bioclimatic studies with insects. Annu. Rev. Entom. 4:183-206.
- Odum, E. P. 1959. Fundamentals of ecology. 2nd ed. W. B. Saunders Co., Philadelphia, Pa. 546 pp.

In the forest in 1967 there were these species with winged

ants:

~:

. (

Lasius (C.) brevicornis microps Wheeler <u>Aphaenogaster (A.) subterranea valida</u> Wheeler (two nests) <u>Lasius niger sitkaensis</u> Pergande <u>Myrmica schencki emeryana</u> Forel <u>Iridomyrmex pruinosus analis</u> (E. André) <u>Lasius niger neoniger Emery</u>

The meadow had no nests with winged ants in 1967. It is of interest to note that some species appeared on the list in 1967, but did not in 1966.

11

÷

10

- Odum, E. D. and A. J. Pontin. 1961. Population density of the underground ant, <u>Lasius flavus</u>, as determined by tagging with P³². Ecology 42:186-188.
- Osborn, W. S. 1953. Estimates of alpine grasshopper densities as determined by modifications of the Lincoln Index. Master's thesis. University of Colorado. 53 pp.
- Pickles, W. 1937. Populations, territories and biomasses of ants at Thornhill, Yorkshire, in 1936. J. Anim. Ecol. 6:54-61.
- 1938. Populations, territories and biomasses of ants at Thornhill, Yorkshire, in 1937. J. Anim. Ecol. 7:370-380.
- Robbins, W. W. 1910. An introduction to the study of the ants of northern Colorado. Univ. Colo. Stud. 7(4):215-222.
- Rodeck, H. G. 1964. Natural history of the Boulder area. University of Colorado Museum, Boulder, Colorado. 100 pp.
- Scherba, G. 1961. Nest structure and reproduction in the moundbuilding ant <u>Formica opeciventris</u> (Emery) in Wyoming. J. New York Entomol. Soc. 69:71-78.
 - 1963. Population characteristics among colonies of the ant Formica opaciventris (Emery) (Hymenoptera:Formicidae). J. New York Entomol. Soc. 71:219-232.
- Schneirla, T. C. 1949. Army-ant life and behaviour under dryseason conditions. 3. The course of reproduction and colony behaviour. Bull. Amer. Mus. Natur. Hist. 94:1-81.
 - 1957. A comparison of species and genera in the ant subfamily Dorylinae with respect to functional pattern. Insectes sociaux 4:259-298.
- Smith, M. E. 1962. The distribution of ants in the vicinity of Valmont Butte, Valmont, Colorado. Master's thesis. University of Colorado. 67 pp.
- Sudd, J. H. 1967. An introduction to the behavior of ants. St. Martin's Press, New York. 200 pp.
- Talbot, M. 1934. Distribution of ant species in the Chicago region with reference to ecological factors and physiological toleration. Ecology 15:416-439.

1961. Mounds of the ant Formica ulkei at the Edwin S. George Reserve, Liwingston County, Michigan. Ecology 42:202-205.

1966. Flights of the ant Aphaenogaster Treatae. J. Kan. Entomol. Soc. 39:67-77. Taussig, W. H. 1962. An ecological study of <u>Formica neorufibarbis</u> <u>gelida</u> Wheeler in the alpine tundra of Colorado. Master's thesis. University of Colorado. 72 pp.

- Weber, N. A. 1935. The biology of the thatching ant Formica rufa obscuripes Forel in North Dakota. Ecol. Mono. 5:166-206.
- Wheeler, G. C. and J. Wheeler. 1963. The ants of North Dakota. University of N. Dakota Press, Grand Forks. 326 pp.

(Hymenoptera; Formicidae). J. Kan. Entomol. Soc. 38:55-61.

Wheeler, W. M. 1910. Ants, their structure, development, and behavior. Columbia University Press, New York. 663 pp.

1917. The mountain ants of western North America. Proc. Amer. Acad. Arts Sci. 52:457-569.

- Wigglesworth, Vincent B. 1965. The principles of insect physiology. Methuen and Co. Ltd., London. 741 pp.
- Wilson, E. O. 1955. A monographic revision of the ant genus Lasius. Bull. Mus. Comp. Zool. 113:1-199.
- Wilson, E. O. and G. L. Hunt. 1966. Habitat selection by queens of two field-dwelling species of ants. Ecology 47(3):485-487.
- Yasuno, M. 1965. The study of the ant population in the grassland at Mt. Hakkoda. IV. The stability of the ant population. Sci. Rep. Tohoku Univ. Ser. IV (Biol.) 31:75-81.

1965. The study of the ant population in the grassland at Mt. Hakkoda. V. The interspecific and intraspecific relation in the formation of the ant population, with special reference to the effect of the removal of <u>Formica truncorum</u> <u>yessensis</u>. Sci. Rep. Tohoku Univ. Ser. IV (Biol.) 31:181-194.

÷

APPENDIX

• ;

.

10

•

-:

í

;

• (

TABLE I

LIST OF ANT SPECIES FOUND IN 1965 IN QUADRAT 1

	Colony number	Date	Species
-	e indre i de la principa de la	June	
	1	15	Lasius (C.) umbratus aphidicola (Walsh)
	2	15	Tapinoma sessile (Say)
	埠		Formica (N.) pallidefulva Latreille
			Solenopsis (D.) molesta validiuscula Emery
	3	15	Lasius (C.) umbratus aphidicola (Walsh)
	4	15	Formica (N.) pallidefulva Latreille
	5	15	Formica (N.) pallidefulva Latreille
	6	15	Lasius (C.) umbratus aphidicola (Walsh)
	7	15	Formica (N.) pallidefulva Latreille
			Tapinoma sessile (Say)
	8	15	Formica (P.) lasioides Emery
	9	15	Formica (N.) pallidefulva Latreille
	10	15	Formica (N.) pallidefulva Latreille
	11	15	<u>Gamponotus (I.) vicinus</u> Mayr
	12	10	Formica (N.) pallidefulva Latreille
	13	16	Formica (P.) lasioides Emery
	14	15	Camponotus (T.) vicinus Mayr
	15	16	Formica (N.) pallidefulva Latreille
	10	16	Lasius (C.) umbratus aphidicola (Walsh)
	17	10	Formica (N.) pallidefulva Latreille
	19	16	Formica (N.) pellidefulva Latreille
	73 ~	10	Camponotus (T.) vicinus Mayr
	20	10	Camponotus (I.) vicinus Mayr
	22	10	Formica (N.) pallidefulva Latreille
	22	10	Iridomyrmex pruinosus analis (E: Andre)
	23	10	Tapinoma sessile (Say)
.,	24	10	Solenopsis (D.) molesta validiuscula Emery
	26	10	Apnaenogaster (A.) subterranea valida kheeler
	20	36	Appaenogaster (A.) subterranea validà bheeler
	28	10	<u>Iapinoma sessile</u> (Say)
	20	10	Abnaenogaster (A.) subterranea valida kheeler
	30	10	Aphaenogaster (A.) subterranea valida kheeler
	37	16	Aphaenogaster (A.) subterranea valida cheeler
	32	16	Lasing allenus americanus imery
	33	116	Lapinora Sessile (Day)
	34	76	Lacing (C) palliderulva Latrellie
	34	16	Formica (N.) pallidefulva Latreille
-	محمد میں اور اور اور اور اور اور اور اور اور اور		LEDIUS MEREI MEONIZET MERY

"when the colony number is omitted, this indicates that more than one species was found beneath the same rock.

•	Colony number	Date	Species
-	Well-followeb-followeb-vie	June	
	36	16	Formica (P.) pallidefulva Latreille
	37	16	Pheidole pilifera coloradensis Emery
	38	16	Formica obscuriventris clivia Creighton
	39	16	Formica (N.) pallidefulva Latreille
	40	16	<u>Solenopsis</u> (D.) <u>rolesta validiuscula</u> Emery
	41	16	Formica (N.) pallidefulva Latreille
	42	16	Lasius niger neoniger Emery
	43	16	Lasius (C.) unbratus aphidicola (Walsh)
		July	
	44	6	Formica (N.) pallidefulva Latreille
	45	6	Formica (N.) pallidefulva Latreille
	46	6	Camponotus (T.) vicinus Mayr
	47	6	Camponotus (T.) vicinus Mayr
	48	6	Aphaenogaster (A.) subterranea valida Wheeler
	49	5	Formica (N.) pallidefulva Latreille
	50	0.	Lasius (A.) claviger coloradensis Wheeler
	51	6	Lasius alienus americanus Emery
	52	0	Aphaenogaster (A.) subterranea valida Wheeler
	22	0	Formica (N.) palliderulva Latreille
	54	0	Lasius alienus americanus Emery
	22	0	Apnaenogaster (A.) subterranea valida wheeler
	50	, O,	Formica (N.) palliderulva Latreille
	21	6	Formica (N.) pallidefulva Latreille
	. <u>50</u>	6	Formica (N.) palliderniva Latreille
	60	6	Formica (N.) palliderulva Latreille
	61	6	Formica (N.) palliderulva Latreille
(62	6	Formica (N.) palliderulva Latreille
	63	2	Toiter (N.) palliceluiva Latreille
	64	7	Lacing aligning analis (L. Andre)
	65	2	Tapinoma cassila (Sam)
	66	2	Equitor (N.) pollidenture Latraille
	67	2	Rownice (N.) pollidefulve Letroille
	68	2	Solenonsia (D.) molecta validiucala imame
	69	7	Formica (N.) pollidefulva Latrailla
	70	. 7	Solenonsis (D.) molesta validinamia Smooth
	77	7	Maria schendel emericana Foral
	72	2	Campanatus (T) status Kamp
	73	2	Lacine alienne americanus France
	74	7	Camonatus (P.) vicinus Meur
	75	2	Salanansie (II.) vicinius mayi
	76	2	Solenonsis (D.) molecta validiucula Emery
	77	7	Forsign (N.) rollidefulty Latrolla
	78	2	Formice (N.) mallidefulty Latreille
	10	(TATTA (10) Parting TAPLATTA

TABLE II

LIST OF ANT SPECIES FOUND IN 1965 IN QUADRAT 2

	Colony number	Date	Species
•		June	
	1	21	Leptothorax rugatulus Emery
	2	21	Crematogaster lineolata (Say)
	3	21	Crematogaster lineolata (Say)
	4	21	Formica (N.) pallidefulva nitidiventria Fmery
	5	21	Crematogaster lineolata (Say)
	6	21	Camponotus (T.) vicinus Mayr
	7	21	Aphaenogaster (A.) subterranea valida Wheeler
	8	21	Aphaenogaster (A.) subterranea valida wheeler
	9	21	Aphaenogaster (A.) subterranea valida wheeler
	10	21	Aphaenogaster (A.) subterranea valida Wheelar
	11	21	Aphaenogaster (A.) subterranea valida Wheeler
	12	21	Lasius niger neoniger Emery
	13	21	Aphaenogaster (A.) subterranea valida wheeler
	14	21	Aphaenogaster (A.) subterranea valida Wheeler
	15	21	Camponatus (T.) vicinus Mayr
	16	21	Solenopsis (D.) molesta validiuscula Emery
	17	21	Formica (N.) pallidefulva Latreille
	18	21	Lasius (C.) brevicornis microps Wheeler
	19	21	Formica (N.) pallidefulva Latreille
	20	21	Formica (N.) pallidefulva Latreille
	21	21	Tapinoma sessile (Say)
	22 -	21	Tapinoma sessile (Say)
	23	22	Aphaenogaster (A.) subterranea valida Wheeler
	- 24	22	Aphaenogaster (A.) subterranea valida Wheeler
	25	22	Solenopsis (D.) molesta validiuscula Emery
	20	22	Aphaenogaster (A.) subterranea valida Wheeler
	21	22	Formica fusca Linnaeus
	20	22	Tapinoma sessile (Say)
	29	22	Aphaenogaster (A.) subterranea valida Wheeler
	30	23	Tapinoma sessile (Say)
	-	00	Aphaenogaster (A.) subterranea valida wheeler
	20	23	Apnaenogaster (A.) subterranea valida Wheeler
	32	4)	Lasius (C.) unoratus apaidicula (Walsh)
	34	22	Toring (C) houring micros liberter
	24	:02	Liandown agaidmtals incrops wheeler
	36	22	Traine bies bourger Incruosom Mueeter
	37	23	Lastus niger neoniger Emery
	21	4)	restro under ucourdet tweeth

*When the colony number is omitted, this indicates that more than one species was found beneath the same rock.

Colony		
number	Date	Creater
		2 hectes
	June	
38	23	Lasius (C.) brevicornia microne literal an
39	23	Lasius niger neoniger Wrowy
40	23	Camponotus (T) ricinus Norm
41	23	Laging (A) latinon (Walch)
	~	Camponotus (T) misimus Kom
42	23	Lasius niger popriger Trans
43	23	Aphanacaton (A) mittery
Lile	23	Lastun niges near a subterranea valida Wheeler
45	23	Lasing niger neoniger Emery
46	23	Lasing (A) Johinge (Halah)
1.7	23	Componetus (T) relations Marsh
48	23	Lading nices needless Exert
110	23	inhomogrator (A) subtamine molide lite ?
50	22	Aphaenogaster (A.) subterranea valida wheeler
5	24	Asids (C.) Drevicornis microps wheeler
52	24.	Appaenogaster (A.) subterranea valida Wheeler
52	24	Tapinona sessile (Say)
55	24	Liometopum occidentale luctuosum Wheeler
54	24	Tapinoma sessile (Say)
22	24	Aphaenogaster (A.) subterranea valida Wheeler
50	24	Llometorum occidentale luctuosum Wheeler
51	24	Liometopum occidentale luctuosum Wheeler
58	24	Crematogaster lineolata (Say)
.59	24	Crematogaster lineolata emeryana Creighton
60	24	Camponotus (T.) vicinus Mayr
61	24	Aphaenogaster (A.) subterranea valida Wheeler
, 62	24	Aphaenogaster (A.) subterranea valida Wheeler
63	24	Aphaenogaster (A.) subterranea valida Wheeler
64	24	Camponotus (T.) vicinus Mayr
65	24	Aphaenogaster (A.) subterranea valida wheeler
66	24	Aphaenogaster (A.) subterranea valida Wheeler
67	24	Aphaenogaster (A.) subterranea valida Wheeler
68	24	Lasius niger neoniger Emery
69	24	Lasius niger sitkaensis Pergande
70	24	Aphaenogaster (A.) subterranea valida Wheeler
71	:24	Aphaenogaster (A.) subterranea valida Wheeler
72	24	Camponotus (T.) vicinus Mayr
73	24	Aphaenogaster (A.) subterranea valida Wheeler
74	24	Lasius niger sitkaensis Pergande
75	24	Lasius niger neoniger Emery
76	24	Aphaenogaster (A.) subterranea valida Wheeler

Colony	7	
number	- Date	Species
Collectory and a top of the second	June	
77	24	Lasius niger naoniger Americ
		Lasius (C.) previcornia mignora libral
78	24	Lasius (C.) brevicornis microps wheeler
79	24	Lasius (C.) bravicornis microps wheeler
80	24	Lasius (C.) brevicornis microps khoolor
81	24	Lasius niger neoniger Emery
82	25	Lasius niger sitkaensis Perganda
83	25	Lasius (C.) umbratus aphidicola (Walch)
84	25	Lasius niger neoniger Enery
85	25	Lasius niger neoniger Deery
86	25	Lasius niger neoniger Fmery
87	25	Lasius niger neoniger Frerv
88	25	Lasius niger neoniger Emery
89	25	Lasius niger neoniger Emery
90	25	Aphaenogaster (A.) subterranea valida Wheeler
91	25	Formica fusca Linnaeus
92	25	Tapinoma sessile (Say)
93	28	Aphaenogaster (A.) subterranea valida Wheeler
94	28	Lasius (C.) brevicornis microps sheelar
95	28	Tapinoma sessile (Say)
96	- 28 ·	Aphaenogaster (A.) subterranea valida wheeler
97	28	Lasius (A.) latipes (Walsh)
-98	28	Lasius (C.) brevicornis microps Wheeler
99	28	Liometopum occidentale luctuosum Wheeler
100	28	Tapinoma sessile (Say)
101	28	Lasius niger neoniger Emery
102	28	Formica fusca Linnaeus
103	128	Aphaenogaster (A.) subterranea valida Wheeler
104	28	Lasius niger neoniger Emery
105	28	Aphaenogaster (A.) subterranea valida Wheeler
106	28	Tapinoma sessile (Say)
		Lasius (C.) brevicornis microps Wheeler
107	28	Aphaenogaster (A.) subterranea valida Wheeler
108	28	Lasius niger neoniger Emery
109	28	Lasius niger neoniger Emery
110	28	Aphaenogaster (A.) subterranea valida Wheeler
111	29	Aphaenogaster (A.) subterranea valida Wheeler
112	29	Formica fusca Linnaeus
113	29	Formica fusca Linnaeus
114	29	Lasius (C.) brevicornis microps Wheeler

LIST OF ANT SPECIES FOUND IN 1965 IN QUADRAT 2

Colony	r Date	Species	
	June		
115	29	Tapinoma sessile (Sav)	
116	29	Lasius (C.) brevicarnia mianana libration	
117	29	Lasius (C.) brevicornis microns wheeler	
118	29	Lasius (C.) brevicornia microps Wheeler	
119	29	Lasius (C.) brevicornie micropy Wheeler	
120	29	Formica fusca Linnaeus	
		Lasius (C.) bravicornia microna libert	
121	29	Aphaenogaster (A.) subtermanea malde the	
122	29	Lasius (C.) brevicornis microna libral	
123	29	Lasius (C.) brevicornia microna libralan	
124	29	Lasius niger neoniger Themy	
125	29	Formica (P.) lasioides Emery	
126	30	Aphaenogaster (A.) subterranea walida Whasley	
127	30	Leptothorax rugatulus Emery	
128	30	Lasius (A.) latipes (Welch)	
129	30	Lasius niger neoniger Emery	
130	30	Lasius niger neoniger Fmery	
131	30	Lasius niger neoniger Fmery	
132	30	Aphaenogaster (A.) subterranea valida khealer	
133	30	Formica fusca Linnaeus	
134	30	Formica fusca Linnaeus	
135	- 30 ·	Lasius niger neoniger Emery	
136	30	Lasius niger neoniger Emery	
137	30	Aphaenogaster (A.) subterranea valida Wheeler	
138	30	Aphaenogaster (A.) subterranea valida Whaeler	
139	30	Lasius (C.) brevicornis microps Wheeler	
140	30	Aphaenogaster (A.) subterranea valida bhaeler	
141	30	Lasius niger neoniger Emery	
142	: 30	Lasius niger neoniger Emery	
143	30	Lasius niger neoniger Emery	
T44	30	Lasius (C.) brevicornis microps Wheeler	
71.4	July		
145	1	Lasius (C.) brevicornis microps Wheeler	
146	1	Lasius (C.) brevicornis microps wheeler	
147	1	Lasius niger neoniger Emery	
148	: 1	Leptothorax rugatulus Emery	
149	1	Leptothorax rugatulus Emery	
150	1	Leptothorax rugatulus Emery	
151	1	Lasius niger neoniger Emery	
152	1	Lasius niger neoniger Emery	

121

Colony		
number	Date	Species
	.Tul v	
153	J	Tootun ménur
154	1	Lasius niger neoniger Emery
155	7	Lasius niger neoniger Emery
156	1	Lasing niger neoniger Emery
157	7	Lasius niger neoniger Emery
158	î	Lasius niger neoniger Emery
159	- 7	Lasius niger neoniger Emery
160	7	Lasius niger neoniger Emery
161	1	Lasius niger neoniger Emery
162	î	Lasius niger neoniger Emery
163	7	Lasius niger neoniger Emery
164	7	Lasius (C.) brevicornis microps Wheeler
165	7	Lasius (C.) previcornis microps Wheeler
166	5	Camponotus (T.) vicinus Mayr
167	7.	Camponotus (T.) vicinus Mayr
168	7	Lamponotus (T.) vicinus Mayr
169	7	Leptotnorax rugatulus Fmery
170	2	Lionatous (T.) vicinus Mayr
171	2	Componetopum occidentale luctuosum Wheeler
172	2	Taninoma and Vicinus Mayr
173	2,	Appropriate (Say)
174	2	Tanipore and A.) subterranea valida wheeler
175	2	Taninoma sessile (Say)
176	2	Formica (N) malles and
177	2	Tapinoma sessila (Saw)
.(Aphaenogaster (A) subtemporter att
178	2	Aphaenogaster (4.) subterranea valida wheeler
179	2	Tapinoma sessile (Saw)
180	2	Crematogaster lineolata (Sour)
181	2	Tapinoma sessile (Say)
182	2	Liometopum occidentale Juctucour Libral
183	2	Lasius niger neoniger Fmery
184	2	Lasius (C.) unbratus aphidicola (Walsh)
185	2	Aphaenogaster (A.) subterranea valida Uhealar
186	2	Aphaenogaster (A.) subterranea valida Wheelar
187	2	Tapinoma sessile (Say)
188	2	Formica (N.) pallidefulva Latreille
189	2	Aphaenogaster (A.) subterranea valida Wheeler
		Anternational An

LIST OF ANT SPECIES FOUND IN 1965 IN QUADRAT 2

Colony	063.9	
number	Date	Species
	July	
190	2	Aphaenogaster (4.) subtermanes molde these
191	2	Aphaenogaster (A) subtornance valida wheeler
192	2	Appaenogaster (A) subterranea Valida Wheeler
193	2	Aphaenogaster (A.) subterranea valida wheeler
194	2	Tapinoma seccila (Sec)
195	2	Camponotus (T.) ricinus Marrie
196	2	Aphaenogration (1) white mayr
197	2	Formica (N.) mallideaular interview valida wheeler
198	2	Camponotus (T) misimus K
199	2	Formica (N.) Dallidefulva Latreilla
		THE THE PARTY PROTOTING

11

-

10

.17

~:

;

• (

123

TABLE III

number	Date	Species
n fil fan ie fan de ferste ferste ferste skriederen. F	July	
1	.6	Formica obscuripes Forel
2	6	Lasius alienus americanus Emery
3	6	Lasing alienus americanus Emery
4	6	Lasius (C.) umbratus aphidicola (Malah)
5	6	Lasius (C.) unbratus aphidicola (Walsh)
6	6	Solenopsis (D.) molesta validiuscula Emery
7	6	Solenopsis (D.) molesta validiuscula Emery
8	6	Lasius (C.) umbratus aphidicola (Walsh)
9	. 6	Solenopsis (D.) molesta validiuscula Dmerv
10	6	Solenopsis (D.) molesta validiuscula Emery
11	6	Lasius (C.) umbratus aphidicola (Walah)
12	6	Lasius (C.) umbratus aphidicola (Malah)
13	6	Formica fusca argentea wheeler
14	6	Lasius (C.) unicratus aphidicola (Walsh)
25	5	Lasius (C.) umbratus aphidicola (Walsh)
16	6.	Lasius (A.) latires (Walsh)
17	6	Lasius (C.) umbratus aphidicola (Malsh)
13	6	Lasius alienus americanus Amery
19	6	Solenopsis (D.) polesta validiuscula Emery
20	6	Lasius alienus americanus Emery
21	6	Solenopsis (D.) molesta validiuscula Emery
22	. 6 .	Solenopsis (D.) molesta validiuscula Emery
23	6	Lasius alienus americanus Emery
- 24	6	Lasius (A.) latipes (Malsh)
25	6	Lasius (C.) unbratus aphidicola (Walsh)
26	7	Lasius (C.) umbratus aphidicola (Walsh)
27	7	Lasius alienus americanus Emery
28	7	Tapinoma sessile (Say)
29	1 7	Tapinoma sessile (Say)
30	7	Lasius (C.) unbratus aphidicola (Walsh)
31	7	Lasius (A.) latipes (Walsh)
32	7	Lasius (C.) umbratus aphidicola (Walsh)
33	7	Formica (N.) pallidefulva nitidiventris Inery
34	7	Solenopsis (D.) molesta validiuscula Emery

TABLE IV

Colony number	Date	Species
1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 13 14 5 16 7 8 9 20 1 22 23 24 5 26 7 8 9 30 1 32 33 4 5 5 6 7 8 9 40	Aug. 1111111111111112222222222222222222222	Lasius (C.) umbratus aphidicola (Walsh) Formica (N.) pallidefulva Latreille Gamponotus (T.) vicinus Mayr Formica (P.) lasioides Emery Camponotus (T.) vicinus Mayr Formica (N.) pallidefulva Latreille Formica (N.) pallidefulva Latreille Formica (N.) pallidefulva Latreille Gamponotus (T.) vicinus Mayr Formica (N.) pallidefulva Latreille Iridomyrmex pruinosus analis (E. André)

	Colony	Data	
		Dave	Species
		Aug.	
	41	2	
	42	2	Formica (N.) pallidefulva Latreilla
	43	2	
	44	2	Formica (N.) pallidefulva Latreille
	45	2	Formica (N.) pallidefulva Latreille
	46	2	Camponotus (T.) vicinus Mayr
	47	2	Camponotus (T.) vicinus Mayr
	48	2	Aphaenogaster (A.) subterranes milde thesi
	49	2	the subject and a valua wheeler
	50	2	
	51	. 3	Formica (N.) pallidafulva Latvoilla
	52	3	Formica (N.) pallidefulva Latrailla
	53	3	A PARTY PARTY INCIDENTIA
	54	3	Lasius alienus americanus Enour
	55	3	Aphaenogaster (A.) subtempone molting
	56	3	Subcerranea valida wheeler
	57	3	
	58	3	
	59	3	Formica (N.) pollidobular Lateration
	60	3	Formica (N.) Dallidatulwa Latrellie
	61 -	3.	dallideldiva Latrellie
	62	3	Formica (N.) pallidatulus Laborator
	-63	3	Iridomyrmey province analis (D
	64	3	Lasius alienus americanus David (E. Andre)
	65	3	aner reality
(66	3	
	67	3	Solenonsis (D.) molecte molecular
	68 i	3	more the star validiuscula Emery
	69	3	Formica (N.) pollidoonim total
	70	3	Latreille
	71	3	Myrmica schencki mammun n
	72	4	Camponotus (T) viginus V
	73	4	VICINUS Mayr
	74	4	Camponotus (T.) vicinus Nam
	75	: 4	the second (re) arething ways.
	76	4	
	77	4	
	78	4	

LIST OF ANT SPECIES FOUND IN 1966 IN QUADRAT 1

Colony number	Date	Species
	Aug.	New Colonies for 1966
79	4	Formica (N.) pallidefulva Latraillo
80	4	Aphaenogaster (A.) subterrance malide it.
81	4	Aphaenogaster (1.) subterranea valida wheeler
82	4	Formica (N.) pallidefulva Latavilla
83	4	Myrmica schencki emeruana Forel
		Lasius niger neoniger Emerry
84	4	Formica (N.) pallidatulma Latroilla
85	4	Formica (N.) pallidefulue Latreille
86	4	Myrica schencki emericana Famal
		Formica (N.) pallidefulra Introdata
87	4	Formica (N.) pallidefulva Latrellie
88	4	Lasius alienus americanus The
89	5	Camponotus (T) misimus Manary
90	5	Pheidole pilifers coloradanata D
91	5	Tapinoma sessila (Say)

et 1

۰.

-

10

-:

;

. (

Ł

127

TABLE V

number Date Species	
number Date Species Aug.	eeler eeler

_	Colony number	r Date	Species
	40	Aug. 8	Aphaenogaster (A.) subterranea valida Wheeler
	41	8	
	42	8	
	45	8	•••••
	45	8	
	46	8	
	47	8	방 <mark>문방</mark> 같이 있는 것은 것은 것은 것은 것을 가지 않는 것을 많이 없다. 것을 많이 있는 것을 것을 했다.
	48	8	
	49	8	Aphaenogaster (A.) subtermanes maid to the a
	50	8	the state with subversalies valida wheeler
	51	8	
	52	8	
	53	9.	Liometopum occidentale luctuosum Wheeler
	54	9	Camponotus (I.) vicinus Mayr
	55	9	Aphaenogaster (A.) subterranea valida Wheeler
	57	9	Liometopum occidentale luctuosum Wheeler
	58	9	Componeture (A) and the second
	59	9	Vicinus Mayr
	60	9	Camponotus (T.) vicimie Marra
	61	9	Camponotus (T.) vicinus Mayr
	62	9	Aphaenogaster (A.) subterranea valida Wheeler
	63	9	ATTIC MICELOL
• •	64	9	Formica (N.) pallidefulva Latreille
	65	9	Aphaenogaster (A.) subterranea valida Wheeler
	67	. 9	Aphaenogaster (A.) subterranea valida Wheeler
	68	9	non Tradina ut
	69	2	Lasius niger neoniger Emery
	70	9	
	71	9	Aphaenogaster (A) subtaneous attant
	72	10	Camponotus (T.) vicinus Norm
	73	.10	The second start
	74	10	
	75	10	Lasius niger neoniger Emery
	76	10	
	77	10	
	10	10	

Colony number	Date	Species
79 80 81 82 83 84 85 86 87 88 89	Aug. 10 10 10 10 10 10 10 10 10	Lasius (C.) brevicornis microps Wheeler Lasius (C.) brevicornis microps Wheeler Lasius niger sitkaensis Pergande Leptothorax rugatulus Emery Lasius niger neoniger Emery
90 91 92 93 94 95 96 97 98 99	10 11 11 11 11 11 11 11	Formica fusca Linnaeus Aphaenogaster (A.) subterranea valida Wheeler Lasius (C.) brevicornis microps Wheeler Aphaenogaster (A.) subterranea valida Wheeler Lasius (A.) latipes (Walsh)
100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117	$\begin{array}{c} 11\\ 11\\ 11\\ 11\\ 11\\ 11\\ 11\\ 11\\ 11\\ 12\\ 12$	Lasius (C.) brevicornis microps Wheeler Lasius niger neoniger Emery Lasius niger neoniger Emery Lasius (C.) brevicornis microps Wheeler Aphaenogaster (A.) subterranea valida Wheeler Lasius niger neoniger Emery Aphaenogaster (A.) subterranea valida Wheeler Aphaenogaster (A.) subterranea valida Wheeler Lasius (C.) brevicornis microps Wheeler Lasius (C.) brevicornis microps Wheeler Lasius (C.) brevicornis microps Wheeler Lasius (C.) brevicornis microps Wheeler

	0.7		
	Colony		
	number	Date	Species
			- <u>F</u> - + - + - + - + - + - + - + - + - + -
		Aug.	
	118	12	Toping (C) browing with a star
	110	10	Lasids (C.) Drevicornis microps wheeler
	172	12	Leptothorax rugatulus Amery
			Lasius (C.) brevicornis microps Wheeler
	120	12	Lasius (C.) brevicornis microps Wheeler
	121	12	
	122	12	Lasius (C.) brevicornis microps wheeler
	123	12	Lasius (C.) brevicornia microns Wheeler
	124	12	
	125	12	Formice (P.) Jaciaidan Energy
	126	12	Appaparaton (A) white a state
	127	73	Aphaenogaster (A.) subterranea valida wheeler
	109	12	hepcoundrax rugatorius Fmery
	120		
	129	13	
	130	13	
	131	13	
	132	13	
	133	13	· · · · · · · · · · · · · · · · · · ·
	134	13	
	135	13	
	136	13	
	137	13	den alter and a second and a
	138	73	
	130	12	
		10	
	140	13	Apnaenogaster (A.) subterranea valida Wheeler
.(141	13	Lasius niger neoniger Emery
	142	13	
	143	13	
	144	13	
	145	13	Lasius (C.) brevicornis microns Wheeler
	146	14	Lasius (C.) bravicornis microna Whaalow
	147	14	Lasius niger neoniger Rmerry
	148	14	The state of the s
	749	74	Tactus (C) busicionestis missiones In 2
	7.50	7/1	LEGIUS (U.) Drevicornis microps Wheeler
	10	2.1	
	171	14	
	152	14	부모님 이렇게 잘 잘 들었다. 그는 것이 아니는 것이 가지 않는 것이 가지?
	153	14	Lasius niger neoniger Emery
	154	14	
	155	14	Lasius niger neoniger Emery
			anterestation successibility and a second se

Colony	Date	Species
	Aug.	
156	14	Lasius niger neoniger Freme
157	14	Lasius niger neoniger Frame
158	14	
159	14	
160	14	Lasius niger neoniger Emery
161	14	Lasius niger neoniger Emery
162	14	Lasius niger neoniger Emery
163	14	Aphaenogaster (A.) subtermance molting
164	14	Aphaenogaster (A.) subterranea valida wheeler
165	14	Camponotus (T.) vicious Kom
166	14	violing rayr
167	14	2mlage
168	15	Leptothorax mestulue Emere
169	15	Camponotus (P.) vicinus Marry
170	15	A A A A A A A A A A A A A A A A A A A
171	15	Formica fusca Timague
172	15	manifer a volume Laiting (1)
173	15	
174	15	
175	15	
176 -	15	
177	15	Aphaenogaster (1.) subtermoned and the
178	15	Any Subterranea Valida Wheeler
179	15	Tapinoma sessile (Sar)
180	15	Cay)
181	15	
182	15	
183	15	tertes.
184	15	Lasius (C.) umbratus anhidiana (u. a.)
185	15	Aphaenogaster (1) autorationa (Walsh)
186	16	Succerranea valida Wheeler
187	16	
188	16	<mark>같은</mark> 이 같은 것은 것이 있는 것은 것이 같은 것이 있는 것이 없는 것이 없는 것이 없다. 것이 있는 것이 없는 것이 없다. 가지 않는 것이 없는 것이 없다. 것이 없는 것이 없 것이 없는 것이 없 않이 없는 것이 없 않이 않는 것이 없는 것이 없는 것이 않는 것이 없는 것이 않이
189	16	
190	16	Aphaenogaster (1) with any
191	16	Aphaerogastar (A) subterranea valida Wheeler
192	16	Aphaenogaster (A.) Subterranea valida Wheeler
193	16	subterranea valida Wheeler
194	16	
	~~	

Colony		
number	Date	Species
-	Aug.	
195	16	2 <u>1.4</u> 1 양성 전 전 전 전 전 전 전 전 전 전 전 전 전 전 전 전 전 전
196	16	
197	16	Aphaenogaster (A.) subterranea valida Whealer
198	16	Camponotus (T.) vicinus Mayr
199	16	Formica (N.) pallidefulva Latreille
		New Colcnies for 1965
200	16	Aphaenogaster (A.) subterranea valida Wheeler
201	16	Aphaenogaster (A.) subterranes valida Wheeler
202	16	Camponotus (T.) vicinus Mayr
203	16	Formica fusca Linnaeus
204	16	Aphaenogaster (A.) subterranea valida wheeler
		Formica fusca Linnaeus
205	17	Aphaenogaster (A.) subterranea valida Wheeler
206	17	Tapinoma sessile (Say)
207	17	Aphaenogaster (A.) subterranea valida Wheeler
208	17	Tapinoma sessile (Say)
209	17	Aphaenogaster (A.) subterransa valida Wheeler
210	17	Lasius niger neoniger Emery
211	17	Lasius niger neoniger Emery
212	17	Aphaenogaster (A.) subterranea valida Wheeler
213	17	Aphaenogaster (A.) subterranea valida' Wheeler
214	17	Tapinoma sessile (Say)
215	17	Formica (N.) pallidefulva Latreille
210	17	Aphaenogaster (A.) subterranea valida Wheeler
279	17	<u>Camponotus (T.) vicinus Mayr</u>
210	17	Aphaenogaster (A.) subterranea valida Wheeler
220	17	Abnaenogaster (A.) subterranea valida Wheeler
223	10	Lasius niger neoniger Emery
222	10	Apnaenogaster (A.) subterranea valida Wheeler
223	10	Lasius (C.) previcornis microus Wheeler
221	10	Laslus niger sittaensis Pergande
225	10	Ashaan niger neoniger Emery
226	18	Aphaenogaster (A.) subterranea valida Wheeler
227	18	Lestus niger neoniger Amery
228	18	Lastus niger neoniger Emery
229	18	Lasing (C) brunie subterranea valida Wheeler
and a grant of the second seco	10	Lagrus (b.) Drevicornis microps Wheeler

TABLE VI

	Colony		
	number	Date	
		20000	Species
-		Ano	
	7	20	Demise 1
	2	20	Formica obscuripes Forel
	~	20	Solenopsis (D.) molesta validiuscula Emery
	2	20	
	4	20	
	5	20	
	6	20	
	7	20	Solenopsis (D.) molesta validinamila man
	8	20	Lasius (C.) umbratus aphidicals (U.T.)
	9	20	Solenopsis (D.) moleste malidicula (Malsh)
	10	20	Lista validiuscula Emery
	11	20	Lasing (C) unhandres and the a
	12	20	aphidicola (Walsh)
	13	20	
	74	20	Transformer () h h h h h
	75	20	Lasius (A.) Latipes (Walsh)
	16	20	Lasius (A.) latipes (Walsh)
	10	20	Formica obscuripes Forel
	17	20	Lasius (C.) umbratus aphidicola (Walsh)
	18	20	Lasius alienus americanus Emery
	19	20	temate
	20	20	
	21	20 .	
	22	20	Solenopsis (D.) molesta validiuscula Emery
	-23	20	
	24	20	
	25	20	2 ¹
(26	21	Tapinoma sessile (Saw)
	27	21	conditioned to a second s
	28	21	
	29	21	
	30	21	Lacine (C) unhactive and it a first
	31	27	Solemonsia (D.) malartus aphicicola (Walsh)
	32	27	Dolenobsis (D.) molesta Validiuscula Emery
	33	27	Ensuine (N)
	3/4	27	Formica (M.) parlidefulva nitidiventris Emery
	14	hinds	<u>Colenopsis</u> (D.) <u>molesta validiuscula</u> Emery
		3	New Colonies for 1966
	35	21	Lasius alienus americanus Emery
	36	21	Lasius (C.) umbratus aphidicola (Walsh)
	37	21	Formica (N.) pallidefulva nitidiventria Emery
	38	21	Solenopsis (D.) molesta validiuscula Emerry
	39	21	Tapinoma sessile (Say)
	40	21	Solenopsis (D.) Folesta valtdinemila Frame
	41	21	Lasius (C.) umbratus aphidicals (Unlah)
			Aphaenogaster (A.) subtamarca militaria
			And Supression (A.) Supression Valida Wheeler

TABLE VII

Colony number	Date	Species
	July	
1	31	Lasius (C.) umbratus aphidicola (Walsh)
2	31	
3	31	
4	31	Formica (N.) pallidefulva Latreille
5	31	
6	31	
7	31	
8	31	
9	31	Formica (N.) pallidefulva Latreille
10	31	Formica (N.) pallidefulva Latreille
11	31	Camponotus (T.) vicinus Mayr
12	31	
13	31	Formica (P.) lasioides Emery
14	31	Camponotus (T.) vicinus Mayr
15	31	Formica (N.) pallidefulva Latreille
16	Aug.	
16	1	Solenopsis (D.) molesta validiuscula Emery
17	1	······
18	l	
19	1	Camponotus (T.) vicinus Mayr
20	1	Camponotus (T.) vicinus Mayr
21	1	Formica (N.) pallidefulva Latreille
22	1	Iridomyrmex pruinosus analis (E. Andre)
23	1	definition in the second se
24	1	ell al O .
25	1	
26	1	
27	1	transpo
28	' 1	
29	1	Aphaenogaster (A.) subterranea valida Wheeler
30	2	Aphaenogaster (A.) subterranea valida Wheeler
31	2	Lasius alienus americanus Emery
32	2	
33	2	denter la
34	2	titleres
35	2	exects
36	2	
37	2	
38	2	Formica obscuriventris clivia Creighton
39	2	
40	2	Solenopsis (D.) molesta validiuscula Emery
Colony number	Date	Species
---	------	---
1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997	Aug.	
41	2	
42	2	
43	2	
2424	2	<u>Solenopsis (D.) molesta validiuscula</u> Emery Formica (N.) pallidefulva Latreille
45	2	
46	2	<u>Camponotus</u> (<u>T.</u>) <u>vicinus</u> Mayr
47	3	Camponotus (T.) vicinus Mayr
48	3	Aphaenogaster (A.) subterranea valida Wheeler
49	3	Aphaenogaster (A.) subterranea valida Wheeler
50	3	Lasius (C.) umbratus aphidicola (Walsh)
51	3	Formica (N.) pallidefulva Latreille
52	3	Formica (N.) callidefulva Latreille
53	3	Formica (N.) pallidefulva Latreille
54	3	alterna and a second
55	3	
56	3	
57	3	Vite
58	3	
59	3	
61	3	Formica (N.) pallidefulva Latreille
62	3	1
63	3	Formica (N.) pallidefulva Latreille
64	3	Lasius (A.) latipes (Walsh)
65	3	
66	3	diverse
67	3	
68 69	3	Solenopsis (D.) molesta validiuscula Emery
70 71	3	Solenopsis (D.) molesta validiuscula Emery
72 73	3	Camponotus (T.) vicinus Mayr
74	33	Camponotus (T.) vicinus Mayr
76	3	
77	ñ	
78	3	termine .
79	2	
	,	

Colony	Date	Species
deterministic constant for the second start	Aug.	
80	4	Aphaenogaster (A.) subterranea valida Wheeler
81	4	
82	4	
83	4	
84	4	
85	4	
86	4	Formica (N.) pallidefulva Latreille
87	4	Formica (N.) pallidefulva Latreille
88	4	Terrer and a second sec
89	4	
90	4	Pheidole pilifera coloradensia Emery
91	4	the second s
		New Colonies for 1967
92	4	Formica (N.) pallidefulva Latreille
93	4	Formica (N.) pallidefulva Latraille
94	4	Tapinoma sessile (Say)
95	4	Solenopsis (D.) molesta validiuscula Emerry
96	4	Solenopsis (D.) molesta validiuscula Emery
97 .	× 4 ×	Solenopsis (D.) molesta validiuscula Emery
98	4	Formica (N.) pallidefulva Latreilla
- 99	4	Formica (N.) pallidefulva Latreille
100	4	Formica (N.) pallidefulva Latreille
Sec.		Solenopsis (D.) molesta validiuscula Emery
101	4	Formica (N.) pallidefulva Latreille
102	4	Solenopsis (D.) molesta validiuscula Emery
103	i 4	Formica (N.) pallidefulva Latreille
104	4	Camponotus (T.) vicinus Mayr
105	4	Lasius alienus americanus Emery
106	5	Formica (N.) pallidefulva Latreille
107	5	Crematogaster lineolata (Say)
108	5	Aphaenogaster (A.) subterranea valida Wheeler
	;	

TABLE VIII

Colony	Date	Species
	Aug.	
1	9	Crematogaster lineolata (Sav)
2	9	
3	9	Crematogaster lineolata (Say)
4	9	
5	9	
6	9	feeter.
7	9	Aphaenogaster (A.) subterranea valida Wheeler
8	9	
9	9	
10	9	Aphaenogaster (A.) subterranea valida Wheeler
11	9	
12	9	
13	9	
14	9	Aphaenogaster (A.) subterranea valida Wheeler
15	9.	Section Se
16	9	
17	9	Tapinoma sessile (Say)
		Formica (P.) limata Wheeler
18	9	Lasius (C.) brevicornis microps Wheeler
		Aphaenogaster subterranea valida Wheeler
19	. 9 .	Solenopsis (D.) molesta validiuscula Emery
		Formica (N.) pallidefulva Latreille
20	9	Solenopsis (D.) molesta validiuscula Emery
21	9	Solenopsis (D.) Molesta validiuscula Emery
22	9	Solenopsis (D.) molesta validiuscula Emery
. 23	9	Camponotus (T.) vicinus Mayr
24	9	
25	1 9	
20	2	Camponotus (T.) vicinus Mayr
28	2	
20	9	Leptothorax rugatulus Emery
27	9	4-mile
37	9	*****
32	0	
33	0	
34	0	
35	0	
36	10	Anhaennogester (A.) subtermones wolids Wheeler
-	~~~	Warner and an announdies Auting Mucatel

	Colony number	Date	Species
		Aug.	
	37	10	
	38	10	
	39	10	
	40	10	
	41	10	
	42	10	
	43	10	
	44	10	
	45	10	
	46	10	Lasius niver neonigen Theme
	47	10	HIGHY HIGH HIGHY
	48	10	Lentothoray muchtulus Descent
	49	10	Deperturian ragaculus Emery
	50	10	exists.
	57	10	Manimore analize (a.)
	52	10	ADDINOMA SESSILE (Say)
	53	10	
	Sh.	10	manual data data data data data data data da
	55	10	Camponotus (T.) vicinus Mayr
	55	10	Apnaenogaster (A.) subterranea valida Wheeler
	50	10	Liometopum occidentale luctuosum Wheeler
	21	10	
	50	10	Crematogaster lineolata (Say)
	27	10	
	60	10	
,	61	10	
'	62	10	
	63	10	
	64	10	
	65	10	Aphaenogaster (A.) subterranea valida Wheeler
	66	10	Tapinoma sessile (Sav)
	67	10	Aphaenogaster (A.) subterranea walida libertan
	68	10	(III) SAPOLITATICA VALLUA MIGOLOF
	69	10	
	70	10	
	71	10	
	72	11	Camponotus (T.) vicinus Marro
	73	11	The reality rays
	74	11	
	75	11	
	76	11	
	77	11	

	Colon	77	
	numbe	r Date	
			Species
		Aug.	
	78	11	Aphaenogaster (A.) subterranes wolde the
			Lasius (C.) brevicornia microne Wheeler
	79	11	Lasius (C.) bravicornis microns Wheeler
	80	11	TOTODS MUGGTEL
	81	11	
	82	11	Lasius niger sitkaensis Pergande
	83	11	Lasius (C.) umbratus aphidicola (Walsh)
	84	11	maisin)
	85	11	Leptothorax rugatulus Emery
	86	11	Leptothoraz rugatulus Emery
	87	11	Lasius niger neoniger Emery
	88	11	annon
	89	11	Minday,
	90	11	
	91	11	Formica fusca Linnaeus
	92	11	Antippe.
	93	11	Aphaenogaster (A.) subterranea valida Wheeler
	94	11	
	95	11	Tapinoma sessile (Say)
	90	11	Aphaenogaster (A.) subterranea valida Wheeler
	51	11	Lasius (A.) latipes (Walsh)
	50	11	divisions
	77	11	
	100	11	Lasius (C.) brevicornis microps Wheeler
• (102	11	Lasius niger neoniger Emery
	102	77	Servery .
	104	177	Tondan
	105	17	Lasius niger neoniger Emery
	106	17	
	107	12	
	108	12	Aphaenogaster (A.) subterranea valida Wheeler
	109	12	Toping without the second seco
	110	12	Approximation figer meeniger Emery
	111	12	Aphaenogaster (A.) subterranea valida Wheeler
	112	12	
	113	12	
	114	12	Anhannanaton ())
	115	12	And a subterranea valida Wheeler
-			

Colony number	Date	Species	
116	Aug.		
117	12	Lasius (G.) brevicornis microps wheeler	
119	12	Lasius (C.) brevicornis microps Wheeler	
110	12	Lasius (C.) <u>Drevicornis microps</u> Wheeler	
120	12	Lasius (C.) brevicornis microps Wheeler	
121	12	Lasids (C.) Drevicornis microps wheeler	
122	12	diamps	
123	12	Lasius (C.) brevicornis microne Wheeler	
124	12	The state of the s	
125	12	Lasius (C.) umbratus aphidicola (Walsh)	
126	12		
127	12		
128	12		
129	12	per elles	
130	13	Lasius niger neoniger Emery	
122	13	Annotan	
133	12	Travers and Provide The	
134	13	Formica Iusca Linnaeus	
135	13		
136	- 13		
137	13		
138	13		
139	13	Formica (P.) lasioides Emery	
140	13		
141	13		
142	13		
143	13		
144	13	Testing (0.) here is	
145	12	Lasius (C.) previcornis microps Wheeler	
140	13	Lasius (C.) previcornis microps Wheeler	
148	13	lasius niger neoniger Emery	
149	13	55 Marine	
150	13	(Polyment	
151	13	55-au	
152	13	Leptothorax rugatulus Emery	,
153	14	Lasius niger neoniger Emery	
154	14	Lasius niger neoniger Emery	
155	14	Lasius niger neoniger Emery	

Colon	У	
numbe	r Date	
		Species
	Aug.	
156	14	Lasius niger neonigen Unema
157	14	Monthe Monthed Miery
158	14	
159	14	
160	14	
161	14	Lasius niger neoniger Drome
162	14	Lasius niger neoniger mery
163	14	THORY THOUT AND THEY
164	14	
165	14	-
166	14	
167	14	
168	14	etiteur
169	14	Camponotus (T) ristory V
170	14	(1) VICINUS MAY
171	14	
172	14	
173	14	Arbana (2)
174	14	Leptothoray mustulus Engen
175	14	There's there's there's
176	- 14 .	Revue
177	14	
178	14	
179	14	Leptothorax migatulus Frame
180	15	Aphaenogaster (A.) subtemprove molities
181	15	Formica (N.) pallidefulva Latreille
182	15	Solenopsis (D.) molesta validiusmia Droma
183	15	the second secon
184	15	Lasius (C.) umbratus aphidicola (Walch)
185	15	(Marsh)
186	15	
187	15	
188	15	
189	15	
190	15	Aphaenogaster (A.) subterranea valida Wheelan
191	15	And a state of the second
192	15	Aphaenogaster (A.) subterranea valida Whealaw
193	15	and Anticipation Activity Anticipation
194	15	Crematogaster lineolata (Sav)
195	15	

Colon	y	
number	r Date	Species
	Aug.	
196	16	Aphaenogaster (A.) subterranea walida Wheeler
197	16	A A A A A A A A A A A A A A A A A A A
198	16	
199	16	
200	16	
201	16	Aphaenogaster (A.) subterranea valida khaalan
202	16	ATTICA MILEDICI.
203	16	Formica fusca Linnaeus
204	16	Tapinoma sessile (Sav)
205	16	Aphaenogaster (A.) subterranes wallda Wheeler
206	16	Lasius niger neoniger mary
207	16	and the second s
208	16	
209	16	Tapinoma sessile (Sav)
210	16	Lasius niger neoniger Fmery
211	16	Lasius niger neoniger Emery
212	16	the second
213	16	
214	16	
215	16	
216	-16 .	Aphaenogaster (A.) subterranea valida Wheeler
217	16	
218	16	Aphaenogaster (A.) subterranea validaiWheeler
219	16	Aphaenogaster (A.) subterranea valida Wheeler
220	16	Lasius niger neoniger Emery
(221	16	Aphaenogaster (A.) subterranea valida Wheeler
222	16	Lasius (C.) brevicornis microps Wheeler
223	16	
224	16	Lasius niger neoniger Emery
225	17	Aphaenogaster (A.) subterranea valida Wheeler
225	17	
227	17	Lasius niger neoniger Emery
228	17	Aphaenogaster (A.) subterranea valida Wheeler
229	17	Lasius (C.) brevicornis microps wheeler
230	17	Camponotus (T.) vicinus Mayr
231	17	
232	17	Camponotus (T.) vicinus Mayr
233	17	Lasius (C.) umbratus aphidicola (Walsh)
234	17	
235	17	Lasius niger neoniger Emery

LIST OF ANT SPECIES FOUND IN 1967 IN QUADRAT 2

Colony	T Det	
nuaser	Date	Species
	Aug	
236	17	Formica fusca Linnaeus
237	17	
230	17	Lasius (C.) brevicornis microps Wheeler
240	17	Approved to a la l
241	17	Aphaenogaster (A.) subterranea valida Wheeler
242	17	Lasius Higer neoniger shery
243	17	Formica fusca Linnaeus
244	17	Aphaenogaster (4.) subternance molide litera
245	17	Lasius niger neoniger Fnerv
246	17	the second
247	17	Aphaenogaster (A.) subterranea valida Wheeler
248	17	
249	17	Lasius niger neoniger Emery
250	18	lasius (C.) umbratus aphidicola (Walsh)
252	18	Lasius (A.) claviger coloradensis Wheeler
253	18	Aphaenogaster (A.) subterranea valida Wheeler
254	18	
255	18	
256	- 18 '	
257	18	Aphaenogaster (A.) subterranea valida Libeeler
258	18	TATAGE MICOLOI
		t do
		<u>New Colonies for 1967</u>
259	18	Aphaenogaster (A.) subterranes valida theolog
260	18	Formica (N.) pallidefulva Latreilla
261	18	Formica (N.) pallidefulva nitidiventria Emery
262	18	Formica (N.) pallidefulva Latreille
203	18	Crematogaster lineolata (Say)
265	10	Leptothorax rugatulus Emery
266	10	Legionary rugatulus Emery
267	18	Lasius (C.) previcornis microps Wheeler
268	18	Anhaenogaster (A) aubtomorps Wheeler
269	18	Lasius niger sitkaensis Perganda
270	18	Lasius (C.) brevicornis microne theolog
271	18	Aphaenogaster (A.) subterranes welder thealer
		NUCOLOGIA VALUE MACOLO

LIST OF ANT SPECIES FOUND IN 1967 IN QUADRAT 2

Aug. 272 18 Formics (M.) pallidefulva Latreille 273 18 Lasius (C.) brevicornis microps Wheeler 274 19 Aphaenogaster (A.) subterranea valida Wheeler 275 19 Myrmics schencki emerysna Forel 276 19 Aphaenogaster (A.) subterranea valida Wheeler 277 19 Iridesyrmex Drinous snalis (E. Andre) 278 19 Lasius niger neoniger Emery 280 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Jasius niger neoniger Emery 284 19 Faulnora sessile (Say) 285 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Faminora sessile (Say) 285 19 Lasius (C.) brevicornis microps Wheeler 289 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Lasius (C.) brevicornis microps Wheeler 289 19 Kornica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Lasius (C.) brevicornis microps Wheeler 297 20 Tabinoma sessile (Say) 298 20 Formica (M.) subterranea valida Wheeler 299 20 Abhaenogaster (A.) Subterranea valida Wheeler 296 20 Lasius (C.) brevicornis microps Wheeler 297 20 Tabinoma sessile (Say) 298 20 Formica (M.) subterranea valida Wheeler 299 20 Abhaenogaster (A.) Subterranea valida Wheeler 290 20 Lortothorax rugatulus Emery 291 20 Lasius (C.) brevicornis microps Wheeler 292 20 Abhaenogaster (A.) Subterranea valida Wheeler 293 20 Formica (M.) Subterranea valida Wheeler 294 20 Abhaenogaster (A.) Subterranea valida Wheeler 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) Subterranea valida Wheeler 307 20 Lortothorax rugatulus Emery 308 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr	Colony	Date	Species
Aug. 272 18 Formics (N.) pallidefulva Latreille 273 18 Lasius (C.) brevicormis microps Wheeler 274 19 Aphaenogaster (A.) subterranea valida Wheeler 275 19 Myrmics schencki emeryana Forel 276 19 Aphaenogaster (A.) subterranea valida Wheeler 277 19 Lindomyrmex pruinosus analis (E. Andre) 278 19 Lasius niger neoniger Emery 279 19 Aphaenogaster (A.) subterranea valida Wheeler 280 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Taminoma cessile (Say) 285 19 Lasius (C.) unbratus anhidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Lasius (C.) brevicornis microps Wheeler 280 19 Lasius (C.) brevicornis microps Wheeler 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Iasius (C.) brevicornis microps Wheeler 285 19 Lasius (C.) brevicornis microps Wheeler 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius niger neoniger Emery 288 19 Aphaenogaster (A.) subterranea valida Wheeler 299 19 Lasius niger neoniger Emery 291 19 Lasius niger neoniger Emery 292 19 Lasius (C.) brevicornis microps Wheeler 294 20 Leptothorax rugatulus Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (N.) pallidefulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Leptothorax rugatulus Emery 301 20 Leptothorax rugatulus Emery 302 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) Subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walan)	-		
 272 18 Formica (N.) pallidefulva Latreille 273 18 Lasius (C.) brevicornis microps Wheeler 274 19 Aphaenogaster (A.) subterranea valida Wheeler 275 19 Myrmica schencki emeryana Forel 276 19 Aphaenogaster (A.) subterranea valida Wheeler 277 19 Iridomyrmex Envirosus schelis (E. Andre) 278 19 Lasius niger neoniger Emery 279 19 Aphaenogaster (A.) subterranea valida Wheeler 280 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Taminora sessile (Say) 285 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Formica fusca Linnaeus 291 19 Lasius (C.) brevicornis microps Wheeler 289 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (N.) pallidefulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Leptothorax rugatulus Emery 201 20 Leptothorax rugatulus Emery 202 20 Leptothorax rugatulus Emery 203 20 Formica fusca Linnaeus 204 20 Camponotus (T.) vicinus Mayr 205 20 Camponotus (T.) vicinus Mayr 205 20 Camponotus (T.) vicinus Mayr 205 20 Lasius (A.) Latipes (Walsh) 	-	Aug.	
 273 18 Lasius (G.) brevicornis microps Wheeler 274 19 Aphaenogaster (A.) subterranea valida Wheeler 276 19 Aphaenogaster (A.) subterranea valida Wheeler 277 19 Iridomyrmen pruinosus analis (E. Andre) 278 19 Lasius niger neoniger Emery 279 19 Aphaenogaster (A.) subterranea valida Wheeler 278 19 Lasius niger neoniger Emery 280 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Lasius niger neoniger Emery 284 19 Tarinoma sessile (Say) 285 19 Lasius (C.) Unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) Unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) Unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) Unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) Unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) Drevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 290 19 Kormica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Lasius (C.) brevicornis microps Wheeler 297 20 Taphoma sessile (Say) 298 20 Formica (M.) mallidafulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida	272	18	Formica (N.) pallidefulva Latreille
 274 19 Abhaenogaster (A.) subterranea valida Wheeler 275 19 Myrmica schencki emeryana Forel 276 19 Abhaenogaster (A.) subterranea valida Wheeler 278 19 Lasius niger neoniger Emery 279 19 Abhaenogaster (A.) subterranea valida Wheeler 280 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Lasius niger neoniger Emery 285 19 Lasius (C.) unbratus achidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) unbratus achidicola (Walsh) 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Lasius (C.) bravicornis microns Wheeler 289 19 Lasius niger neoniger Emery 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius niger neoniger Emery 296 20 Lasius niger neoniger Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallicarNiva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 291 20 Lasius niger neoniger Emery 292 30 Lasius niger neoniger Emery 293 40 Lasius niger neoniger Emery 294 40 Lasius niger neoniger Emery 295 20 Lasius (C.) bravicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallicarNiva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 291 20 Leptothorax rugatulus Emery 202 20 Leptothorax rugatulus Emery 303 20 Formica (M.) subterranea valida Wheeler 304 30 Gamponotus (T.) victuus Mayr 304 30 Gamponotus (T.) victuus	273	18	Lasius (C.) brevicornis microps Wheeler
 275 19 Myrmica schencki emeryana Forel 276 19 Aphaenogaster (A.) subterranea valida Wheeler 278 19 Lastus niger neoniger Emery 279 19 Aphaenogaster (A.) subterranea valida Wheeler 280 19 Lastus niger neoniger Emery 281 19 Lastus niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Taphona gesalle (Say) 285 19 Lastus (C.) unbratus achidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lastus (C.) unbratus achidicola (Walsh) 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Eornica fusca Linnaeus 290 19 Fornica fusca Linnaeus 291 19 Lastus niger neoniger Emery 292 19 Lastus niger neoniger Emery 293 19 Lastus niger neoniger Emery 294 19 Lastus niger neoniger Emery 295 20 Lastus niger neoniger Emery 296 20 Lertothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Fornica (M.) pallicafulya Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 296 20 Lertothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Fornica (M.) pallicafulya Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 201 20 Leptothorax rugatulus Emery 202 20 Leptothorax rugatulus Emery 203 20 Fornica fusca Linnaeus 204 20 Leptothorax rugatulus Emery 205 20 Lastus (G.) bravicornis Mayr 206 20 Aphaenogaster (A.) subterranea valida Wheeler 206 20 Aphaenogaster (A.) subterranea valida Wheeler 206 20 Aphaenogaster (A.) subterranea valida Wheeler 207 20 Aphaenogaster (A.) subterranea valida Wheeler 208 20 Fornica (M.) pallicafulya Latreille 209 20 Aphaenogaster (A.) subterranea valida Wheeler 202	274	19	Aphaenogaster (A.) subterranea valida Wheeler
 276 19 Aphaenogaster (A.) subterranea valida wheeler 277 19 Iridomyrtex pruinous snalis (E. Andre) 278 19 Lasius niger neoniger Emery 279 19 Aphaenogaster (A.) subterranea valida wheeler 280 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida wheeler 283 19 Aphaenogaster (A.) subterranea valida wheeler 284 19 Taninoma sessile (Say) 285 19 Lasius (C.) unbratus achidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida wheeler 287 19 Lasius (C.) unbratus achidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida wheeler 287 19 Lasius (C.) bravicornis microps wheeler 288 19 Aphaenogaster (A.) subterranea valida wheeler 289 19 Kornica fusca Linnaeus 290 19 Kornica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius (C.) bravicornis microps wheeler 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) bravicornis microps wheeler 296 20 Lestothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Aphaenogaster (A.) subterranea valida wheeler 299 20 Aphaenogaster (A.) subterranea valida wheeler 296 20 Lestothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Aphaenogaster (A.) subterranea valida wheeler 299 20 Aphaenogaster (A.) subterranea valida wheeler 299 20 Lasius (T.) yotinus Mayr 200 20 Aphaenogaster (A.) subterranea valida wheeler 201 20 Lestothorax rugatulus Emery 202 20 Lestothorax rugatulus Emery 203 20 Formica fusca Linnaeus 204 20 Gamonotus (T.) vicinus Mayr 205 20 Gamonotus (T.) vicinus Mayr 206 20 Aphaenogaster (A.) subterranea valida wheeler 207 20 Formica fusca Linnaeus 208 20 Aph	275	19	Myrmica schencki emeryana Forel
 277 19 <u>Iridemyrtex pruinosus analis</u> (E. Andre) 278 19 <u>Iasius niger neoniger Emery</u> 279 19 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 280 19 <u>Iasius niger neoniger Emery</u> 281 19 <u>Iasius niger neoniger Emery</u> 282 19 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 283 19 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 284 19 <u>Taphnema Sessile (Say)</u> 285 19 <u>Iasius (C.) unbratus aphidicola (Walsh)</u> 286 19 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 287 19 <u>Iasius (C.) unbratus aphidicola (Walsh)</u> 286 19 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 287 19 <u>Iasius (C.) brevicornis microps</u> Wheeler 288 19 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 289 19 <u>Fornica fusca Linnaeus</u> 290 19 <u>Fornica fusca Linnaeus</u> 291 19 <u>Iasius niger neoniger Emery</u> 292 19 <u>Iasius niger neoniger Emery</u> 293 19 <u>Iasius (C.) brevicornis microps</u> Wheeler 294 19 <u>Iasius niger neoniger Emery</u> 295 20 <u>Iasius (C.) brevicornis microps</u> Wheeler 296 20 <u>Lestothorax rugatulus Emery</u> 297 20 <u>Tapinoma sessile (Say)</u> 298 20 <u>Fornica (M.) pallidefulva Latreille</u> 299 20 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 201 20 <u>Iastothorax rugatulus Emery</u> 302 20 <u>Leptothorax rugatulus Emery</u> 303 20 <u>Fornica fusca Linnaeus</u> 304 20 <u>Camponotus (T.) vicinus Mayr</u> 305 20 <u>Camponotus (T.) vicinus Mayr</u> 306 20 <u>Aphaenogaster (A.) subterranes valida Wheeler</u> 307 20 <u>Lasius (A.) Jatipes (Walah)</u> 	270	19	Aphaenogaster (A.) subterranea valida Wheeler
 275 19 Lasius niger neoniger Emery 279 19 Aphaenogaster (A.) subterranea valida Wheeler 280 19 Lasius niger neoniger Emery 281 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Tarinora sessile (Say) 285 19 Lasius (C.) unbratus achidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) unbratus achidicola (Walsh) 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Formica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (B.) mallidafulva Latreille (Say) 298 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 201 20 Leptothorax rugatulus Emery 202 20 Leptothorax rugatulus Emery 203 20 Formica fusca Linnaeus 204 20 Leptothorax rugatulus Emery 205 20 Lasius (C.) brevicornis Materianea valida Wheeler 205 20 Lasius (C.) brevicornis microps Vheeler 206 20 Aphaenogaster (A.) subterranea valida Wheeler 209 20 Aphaenogaster (A.) subterranea valida Wheeler 209 20 Aphaenogaster (A.) subterranea valida Wheeler 209 20 Aphaenogaster (A.) subterranea valida Wheeler 201 20 Leptothorax rugatulus Emery 302 30 Formica fusca Linnaeus 304 30 Gamonotus (T.) vicinus Mayr 305 30 40 30 Gamonotus (T.) vicinus Mayr 305 40 40 Gamonotus (T.) vicinus	211	19	Iridomyrmex pruinosus analis (E. Andre)
 279 19 Anhaenogaster (A.) subterranea valida Wheeler 280 19 Lasius niger neoniger Enery 281 19 Lasius niger neoniger Enery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Tapinoma sessile (Say) 285 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Lasius (C.) brevicornis microps Wheeler 289 19 Formica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Enery 292 19 Lasius niger neoniger Enery 293 19 Lasius (C.) brevicornis microps Wheeler 294 19 Lasius niger neoniger Enery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Enery 297 20 Tapinoma sessile (Say) 298 20 Formica (N.) pallidafulva Latretile 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Eornica (N.) pallidafulva Latretile 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 201 20 Leptothorax rugatulus Enery 302 20 Formica (N.) subterranea valida Wheeler 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	278	19	Lasius niger neoniger Enery
 19 Lasius niger neoniger Emery 19 Lasius niger neoniger Emery 19 Aphaenogaster (A.) subterranea valida Wheeler 19 Aphaenogaster (A.) subterranea valida Wheeler 19 Aphaenogaster (A.) subterranea valida Wheeler 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Formica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Lasius (G.) brevicornis microps Wheeler 297 20 Tapinoma sessile (Say) 298 20 Formica (W.) pallidafulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 201 20 Leptothorax rugatulus Emery 302 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) Subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	219	19	Aphaenogaster (A.) subterranea valida Wheeler
 19 Lasius niger neoniger Emery 282 19 Aphaenogaster (A.) subterranea valida Wheeler 283 19 Aphaenogaster (A.) subterranea valida Wheeler 284 19 Taninoma sessile (Say) 285 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Formica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius (C.) brevicornis microps Wheeler 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Lasius (C.) brevicornis microps Wheeler 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallidafulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallidafulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 201 20 Leptothorax rugatulus Emery 202 20 Leptothorax rugatulus Emery 203 20 Formica (M.) subterranea valida Wheeler 204 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 304 20 Camponotus (T.) vicinus Mayr 305 20 Lasius (A.) Latipes (Walsh) 	280	19	Lasius niger neoniger Enery
 Aphaenogaster (A.) subterranea valida Wheeler Aphaenogaster (A.) subterranea valida Wheeler Tapinoma sessile (Say) I asius (C.) umbratus aphidicola (Walsh) Aphaenogaster (A.) subterranea valida Wheeler P Aphaenogaster (A.) subterranea valida Wheeler P Aphaenogaster (A.) subterranea valida Wheeler P Aphaenogaster (A.) subterranea valida Wheeler P Formica fusca Linnaeus Formica fusca Linnaeus Formica fusca Linnaeus I asius (G.) brevicornis microps Wheeler I asius (G.) brevicornis microps Wheeler Lasius niger neoniger Emery Lasius (G.) brevicornis microps Wheeler Lasius (A.) pallidafulva Latreille Aphaenogaster (A.) subterranea valida Wheeler Aphaenogaster (A.) subterranea valida Wheeler Aphaenogaster (A.) subterranea valida Wheeler Leptothorax rugatulus Emery Leptothorax fusca Linnaeus Gamponotus (T.) vicinus Mayr Gamponotus (T.) vicinus Mayr Aphaenogaster (A.) subterranea valida Wheeler Aphaenogaster (A.) subterranea valida Wheeler Aphaenogaster (A.) subterranea valida Wheeler 	281	19	Lasius niger neoniger Emery
 283 19 Aphaencgaster (A.) subterranea valida Wheeler 284 19 Tapinoma sessile (Say) 285 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Formica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius niger neoniger Emery 296 20 Lasius (C.) brevicornis microps Wheeler 297 20 Lasius (S.) brevicornis microps Wheeler 298 20 Formica (N.) pallidafulya Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 301 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Gamponotus (T.) vicinus Mayr 305 20 Gamponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	282	19	Aphaenogaster (A.) subterranea valida Wheeler
 19 Taplnoma sessile (Say) 285 19 Lasius (C.) unbratus aphidicola (Walsh) 286 19 Aphaenogaster (A.) subterranea valida Wheeler 287 19 Lasius (C.) brevicornis microps Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 288 19 Aphaenogaster (A.) subterranea valida Wheeler 289 19 Formica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallidafulva Latreille : 299 20 Aphaenogaster (A.) subterranea valida Wheeler 301 20 Leptothorax rugatulus Emery 302 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	203	19	Aphaenogaster (A.) subterranea valida Wheeler
 19 Lasius (C.) unbratus aphidicola (Walsh) Aphaenogaster (A.) subterranea valida Wheeler 19 Lasius (C.) brevicornis microps Wheeler 28 19 Aphaenogaster (A.) subterranea valida Wheeler 28 19 Aphaenogaster (A.) subterranea valida Wheeler 28 19 Fornica fusca Linnaeus 290 19 Fornica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius niger neoniger Emery 296 20 Leptothorax rugatulus Emery 298 20 Formica (N.) pallidafulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 299 20 Aphaenogaster (A.) subterranea valida Wheeler 300 20 Aphaenogaster (A.) subterranea valida Wheeler 301 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 304 20 Lastus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lastus (A.) Latipes (Walsh) 	204	19	Tapinoma sessile (Say)
 Aphaenogaster (A.) subterranea valida Wheeler 19 Lasius (C.) brevicornis microps Wheeler Aphaenogaster (A.) subterranea valida Wheeler Aphaenogaster (A.) subterranea valida Wheeler Pormica fusca Linnaeus Lasius niger neoniger Emery Pormica fusca Linnaeus Pormica fusca Linnaeus<	205	19.	Lasius (C.) umbratus aphidicola (Walsh)
 267 19 Lasius (C.) brevicornis microps Wheeler 288 19 Arhaenogaster (A.) subterranea valida Wheeler 289 19 Formica fusca Linnaeus 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius niger neoniger Emery 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinowa sessile (Say) 298 20 Formica (M.) pallidefulva Latreille 299 20 Aphaenogaster (A.) subterranea valida Wheeler 300 20 Aphaenogaster (A.) subterranea valida Wheeler 301 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	200	19	Aphaenogaster (A.) subterranea valida Wheeler
 Aphaenogaster (A.) subterranea valida wheeler Formica fusca Linnaeus Formica fusca Linnaeus Formica fusca Linnaeus Formica fusca Linnaeus Isius niger neoniger Emery Lasius (G.) brevicornis microps Wheeler Pormica (M.) pallidefulva Latreille Formica (M.) pallidefulva Latreille Aphaenogaster (A.) subterranea valida Wheeler Lastothorax rugatulus Emery Lastothorax rugatulus Emery Lastothorax rugatulus Emery Camponotus (T.) vicinus Mayr Camponotus (T.) vicinus Mayr Camponotus (T.) vicinus Mayr Aphaenogaster (A.) subterranea valida Wheeler Lasius (A.) Latipes (Walsh) 	287	19	Lasius (C.) brevicornis microps Wheeler
 19 Formica fusca Linnaeus 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius (C.) brevicornis microps Wheeler 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallidefulva Latreille 299 20 Aphaenogaster (A.) subterranes valida Wheeler 300 20 Leptothorax rugatulus Emery 301 20 Leptothorax rugatulus Emery 302 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Gamponotus (T.) vicinus Mayr 305 20 Gamponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	208	19	Aphaenogaster (A.) subterranea valida Wheeler
 290 19 Formica fusca Linnaeus 291 19 Lasius niger neoniger Emery 292 19 Lasius niger neoniger Emery 293 19 Lasius (C.) brevicornis microps Wheeler 294 19 Lasius niger neoniger Emery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallidefulva Latreille 299 20 Aphaenogaster (A.) subterranes valida Wheeler 300 20 Aphaenogaster (A.) subterranes valida Wheeler 301 20 Leptothorax rugatulus Emery 302 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	209	19	Fornica fusca Linnaeus
 19 Lasius niger neoniger Emery 19 Lasius niger neoniger Emery 19 Lasius (C.) brevicornis microps Wheeler 19 Lasius (C.) brevicornis microps Wheeler 19 Lasius (C.) brevicornis microps Wheeler 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (N.) pallidefulva Latreille 299 20 Aphaenogaster (A.) subterranes valida Wheeler 300 20 Aphaenogaster (A.) subterranes valida Wheeler 301 20 Leptothorax rugatulus Emery 302 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranes valida Wheeler 307 20 Lasius (A.) Latipes (Walsh) 	290	19	Formica fusca Linnaeus
29219Lasius niger neoniger Emery29319Lasius (C.) brevicornis microps Wheeler29419Lasius niger neoniger Emery29520Lasius (C.) brevicornis microps Wheeler29620Leptothorax rugatulus Emery29720Tapinoma sessila (Say)29820Formica (M.) pallidefulva Latreille29920Aphaenogaster (A.) subterranes valida Wheeler30020Aphaenogaster (A.) subterranea valida Wheeler30120Leptothorax rugatulus Emery30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler	291	. 19	Lasius niger neoniger Emery
29519Lasius (C.) brevicornis microps Wheeler29419Lasius niger neoniger Enery29520Lasius (C.) brevicornis microps Wheeler29620Leptothorax rugatulus Emery29720Tapinoma sessile (Say)29820Formica (M.) pallidefulva Latreille29920Aphaenogaster (A.) subterranes valida Wheeler30020Aphaenogaster (A.) subterranea valida Wheeler30120Leptothorax rugatulus Emery30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler	292	19	Lasius niger neoniger Emery
 19 Lasius niger neoniger Enery 295 20 Lasius (C.) brevicornis microps Wheeler 296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (M.) pallidefulva Latreille 299 20 Aphaenogaster (A.) subterranes valida Wheeler 300 20 Aphaenogaster (A.) subterranea valida Wheeler 301 20 Leptothorax rugatulus Emery 302 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Gamonotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 	293	19	Lasius (C.) brevicornis microps Wheeler
29520Lasius (C.) brevicornis microps Wheeler29620Leptothorax rugatulus Emery29720Tapinoma sessila (Say)29820Formica (N.) pallidafulva Latreille29920Aphaenogaster (A.) subterranes valida Wheeler30020Aphaenogaster (A.) subterranea valida Wheeler30120Leptothorax rugatulus Emery30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler	294	19	Lasius niger neoniger Enery
296 20 Leptothorax rugatulus Emery 297 20 Tapinoma sessile (Say) 298 20 Formica (N.) pallidefulva Latreille : 299 20 Aphaenogaster (A.) subterranes valida Wheeler 300 20 Aphaenogaster (A.) subterranea valida Wheeler 301 20 Leptothorax rugatulus Emery 302 20 Leptothorax rugatulus Emery 303 20 Formica fusca Linnaeus 304 20 Camponotus (T.) vicinus Mayr 305 20 Camponotus (T.) vicinus Mayr 306 20 Aphaenogaster (A.) subterranea valida Wheeler 307 20 Lasius (A.) Latipes (Walsh)	295	20	Lasius (C.) brevicornis microps Wheeler
 297 20 <u>Tapinoma sessila</u> (Say) 298 20 Formica (N.) pallidafulva Latreille ; 299 20 <u>Aphaenogaster (A.) subterranea valida</u> Wheeler 300 20 <u>Aphaenogaster (A.) subterranea valida</u> Wheeler 301 20 <u>Leptothorax rugatulus</u> Emery 302 20 <u>Leptothorax rugatulus</u> Emery 303 20 <u>Formica fusca Linnaeus</u> 304 20 <u>Camponotus (T.) vicinus</u> Mayr 305 20 <u>Camponotus (T.) vicinus</u> Mayr 306 20 <u>Aphaenogaster (A.) subterranea valida</u> Wheeler 307 20 <u>Lasius (A.) latipes</u> (Walsh) 	. 290	20	Leptothorax rugatulus Emery
29820Formica (M.) pallidefulva Latreille29920Aphaenogaster (A.) subterranes valida Wheeler30020Aphaenogaster (A.) subterranea valida Wheeler30120Leptothorax rugatulus Emery30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Gamponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	297	20	Tapinoma sessila (Say)
29920Aphaenogaster (A.) subterranes valida Wheeler30020Aphaenogaster (A.) subterranes valida Wheeler30120Leptothorax rugatulus Emery30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	298	,20	Formica (N.) pallidefulva Latreille
30020Aphaenogaster (A.) subterranea valida Wheeler30120Leptothorax rugatulus Emery30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	299	20	Aphaenogaster (A.) subterranes valida Wheeler
30120Leptothorax rugatulus Emery30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	300	20	Aphaenogaster (A.) subterranea valida Wheeler
30220Leptothorax rugatulus Emery30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	301	20	Leptothorax rugatulus Emery
30320Formica fusca Linnaeus30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	302	20	Leptothorax rugatulus Emery
30420Camponotus (T.) vicinus Mayr30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	303	20	Formica fusca Linnaeus
30520Camponotus (T.) vicinus Mayr30620Aphaenogaster (A.) subterranea valida Wheeler30720Lasius (A.) latipes (Walsh)	304	20	Camponotus (T.) vicinus Mayr
306 20 <u>Aphaenogaster (A.) subterranea valida</u> Wheeler 307 20 <u>Lasius (A.) latipes</u> (Walsh)	305	20	Camponotus (T.) vicinus Mayr
307 20 Lasius (A.) latipes (Walsh)	306	20	Aphaenogaster (A.) subterranea valida Wheeler
	307	20	Lasius (A.) latipes (Walsh)

TABLE IX

Colony number	Date	Species
1 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 10 11 2 3 4 5 6 7 8 9 21 22 3 4 5 6 7 8 9 30 11 2 3 4 5 6 7 8 9 30 31 32 33 4 5 6 7 8 39 30 31 32 33 4 5 6 7 8 39	Aug. 666666666666666666666666666666666666	Formica obscuripes Forel Solenopsis (D.) molesta validiuscula Emery Lasius alienus americanus Emery Solenopsis (D.) molesta validiuscula Emery Lasius (C.) umbratus aphidicola (Walsh) Lasius (C.) umbratus aphidicola (Walsh) Hasius (A.) latipes (Walsh) Formica obscuripes Forel Lasius (C.) umbratus aphidicola (Walsh) Hasius (C.) umbratus aphidicola (Walsh) Hasius (C.) umbratus aphidicola (Walsh) Formica (M.) pallidefulva nitidiventris Emery Formica (M.) pallidefulva nitidiventris Emery Lasius (C.) umbratus aphidicola (Walsh)

LIST OF ANT SPECIES FOUND IN 1967 IN QUADRAT 3

Colony number	Date	Species
40 41	Aug. 7 7	Lasius (C.) umbratus aphidicola (Walsh) Lasius (C.) umbratus aphidicola (Walsh)
		New Colonies for 1967
42 43 44 45 46	7 7 7 7	Lasius (C.) umbratus aphidicola (Walsh) Solenopsis (D.) molesta validiuscula Emery Formica fusca Linnaeus Tapinoma sessile (Say) Formica (N.) pallidefulva nitidiventris Emery

-:

;

.(

....

..

• ;

-

	al Zonation	00 Upper Sonoran, Transition Canadian	05 Upper Sonoran, Transition, Canadian, Hudsonian	00 Transition	00 Upper Sonoran, Transition 00 Transition, Upper Sonoran,	0 Upper Sonoran, Transition, Canadian Fudernia	0 Upper Sonoran, Transition	0 Upper Sonoran, Transition, Canadian, Hudsonian	0 Upper Sonoran, Transition, Canadian	0 Upper Sonoran, Transition, Canadian, Hudsonian	0 Upper Sonoran, Transition, Canadian, Mudsonfan Arctic	Upper Sonoran, Transition, O Canddian, Hudsonian, Arctic
SPECIES	Altitudin: Range in feet	5,000-9,7	4,800-10,50	5,850-6,9(3,500-8,00	4.500-11.00	3,500-9,50	5,354-10,00	3,500-9,60	3,500-10,40	·3.500-12,400	4,600-12,200
ES AND ZONATION OF	Range	Rocky Mts. & N. & Midwest	No. U.S. & S. into Mts.	E. & S.E. U.S., Spotty in Midwest & Colo.	N. & E. U.S. Holarctic, N. & S. into Mts.	E., Midwest, & W. U.S.	N. U.S. and S. into Rockies	Mostly N. U.S. & S. into Rockies	rutuwest u.S. to W. Coast S. into Mex.	E. & Cent. U.S.	C.U TRAVEST U.S.	N. U.S. Transcontinental
RANG	Species	Subfamily FORMICINAE Formica (P.) limata Emery Formica (P.) lasiatan in.	Jereann Sentorcer () aphrony	allianer stratenerted to (N) A	Formica fusca Linnaeus	Formice fusce argentes Wheeler	F. Obscuriventris rlivis fund that the	Camponotus (T.) vicinus Mavr	Lasius allenus americamis Dueve	Lasius niger neoniger hmary	Lasius niver sitksensis Dominals	Terbanne

TABLE X

TABLE X (Continued)

. (

RANGES AND ZONATION OF SPECIES

	Zonation	Transition, Canadian	Upper Sonoran, Transition, Canadian	Upper Sonoran, Transition Upper Sonoran, Transition		Upper Sonoran Procest+1	Upper Sonoran, Transition	Upper Sonoran, Transition,	Veueuran, Indsonian	U. Sonoran, Transition,	Venagian Upper Sonoran, Transition	Upper Sonoran. Transition	Upper Sonoran, Transition	Upper Sonorse
	Altitudinal Range in feet	5.700-9.700	5,154-9,500	5,100-8,000 4,800-8,500		3,500-6,300	4,800-7,550	3.500-10,505	5.400-6 EDD	5,800-9,713	5.354-7.500	3,500-8,378	5,354-8,700	5,500-5,900
14	Range	N. U.S. Exten- sions S. & W. M. B. S.	Mq.E. & E. U.S.	Rocky Mountain N. U.S. into E. Rockies & W. Mts.		Western U.S.	E. edge of Rocky Mts., Calif. Mts.	Transcontinental U. S.	N.E. & E. U.S.	E. and Midwest U.S.	tocky Mts., N.W. ind W. U.S.	workocky Mts.	<pre>4. U.S., Rockies 5. to Black Hills</pre>	locky Mts. to Daks.
	Species	Lasius (C.) brevicornis microps Wheeler Lasius (C.) umbratus anhidicola (Walch)	Lasius (A.) claviver colonofordent un	Lasius (A.) latipes (Walsh)	Subfamily DOLICHODERINAE	Iridomyrmex pruinosus analis (E.Andre) Liometopum occidentala 1	Tapinoma sessila (carr)	(Apr) Attacks	Subfamily MYRMICINAE Crematogaster lineolata (Say)"	Aphaenogaster (A) anti-	Solenopsis (D.) molesta walida walida	Leptothorax rugatulus Emerv	Pheidole pilifera coloradancia un	

TABLE XI

• (

SPECIES COMPOSITION OF Q1

;

	19(55	ſ	966		1967	
Species	Colonies Number	Species Number	Colonie Number	s Species % Number	Coloni Number	Res.	Species Number
		ม้ เป็นการที่สามารถสามารถที่สามารถที่สามารถที่สามารถที่สามารถที่สามารถที่สามารถที่สามารถที่สามารถที่					
Lasius (C.) umbratus aphidicola (Walsh)	6 7	3	H	1.9	~	3.5	
L. alienus americanus Emery	2	-	m	5.8	~	3.5	
L. niger neoniger Emery	2	4	н	1.9			
I. claviger coloradensis Wheeler	1	2					
L. (A.) latipes (Walsh)					н	1.7	
Formica (N.) pallidefulva Latreille	32 39.	0	22 4	2.3	2	36.8	
F. (P.) lasioides Emery	2	4	-	1.9	Ч	1.7	
F. obscuriventris clivia Creighton	1	2	Ч	1.9	Ч	1.7	
Camponetus (T.) vicinus Mayr	8	8	6	7.3	6	15.8	
FORMICINAE	57 69.	8 11	38 7	3.0 7	37	64.7	2
Crematogaster lineolata (Say)					Ч	1.7	
Aphaenogaster (A.) subterranea valida Wheeler	8	8	9	1.5	9	10.5	
Solenopsis (D.) molesta validiuscula Emery	7 8	2	Ч	1.9	10	27.5	
Myruica schencki emeryana Forel	1	2	m	5.8			
Pheidole pilifera coloradensis Emery	1 1	2	ч	1.9	Ч	1.7	
MYRMICINAE	17 20.	4 2	11 2	1.1 4	18	31.4	4
Iridomyrnex pruinosus analis (E. Andre)	2. 2	4	2	3.8	Ч	1.7	
Tapinoma sessile (Say)	6. 2	e	-1	1.9	Ч	7.7	
DOLICHODERINAE	8	7 2	e	5.7 2	2	3.4	2
TOTALS	82 99.	8 14	52 9	9.8 13	57	99.5	13

TABLE XII

• (

SPECIES COMPOSITION OF Q2

1

:

		1965			1966	and a second second		1967	
Species	Coloni	es øg	Species	Colon	ies r %	Species	Color	16s	Species
								2	TECHNINA
Lasius (C.) umbratus aphidicola (walsh)	e	1.5		3	~		9	3.6	
Le niker neoniger Emery	77	21.5		30	20.1		00	200	
L. (A.) claviger coloradensis wheeler		•					1		
L. (A.) latipes (Walsh)	77	0.0		~	0		4 0	•	
L. (C.) brevicornis microne Wheelen	1 0	20		+ ;			N	7.5	
T nime sitiscants Damas 2	Ŋ.	7.77		2	14.1		22	13.3	
and a structure of the stand of	4	2.0		t	2.2		2	1.2	
allianter In. / particelulva Latreille	9	2.9		4	2.7		9	3.6	
r. (N.) Pallidelliva nitidiventris Emery	~	1.0					-	9	
r. (r.) 11mata Wheeler							-	N.	
Fe fusca Linnaeus	6	4-4		6	4 77		1 a	-	
F. (P.) lasioides Emery		v			- 0		D ;	2	
Camponotus (T.,) vicinus Mayr	2	0		14			4 (•	
FORMICINAR				ot	1.nt		ת	5.5	
Cremetorseter line leave leave	CTT	55.3	TO	87	58.4	6	88	53.2	12
Anhaeventer (1) 111 (387)	9	5.0		Ч	2.		5	3.0	
Munice of A. Subterranea valida Wheeler	52	25.4		48	32.2		E	26.7	
TATHLOR SCHERCKI EMERYANA POREL							-	9	
The received a set of the recta validius cula Emery	~	1.0					v	3.0	
reprovinces rugatulus Emery	9	2.9		v	4-6		10	20	
WI NUTCTNER	. 99	32.2	7	17	24.2	c	22	100	L
Iridomyrmex pruinosus analis (E. Andre)	1		-	5		2	5	0.01	n
Liometopum occidentale luctuogum Wheeler	6	7-5		~	0 0		-1 r	•	
Tapinoma sessile (Sav)	- 0			1			4	•	
TOL TCHODERTMAR	AT A	2.		5	3.4		ω	4.8	
	202	12.7	2	ω	5.4	2	TO	6.0	e
TOTALS	205 10	00.2	16	149 I	1.00	14	165	99.8	50

TABLE XIII

SPECIES COMPOSITION OF Q3

:

Species	T T T T	1965			1966			1967	
	19qunN	88	Species	Coloni	88	Species	Color	nies er 4	Species Number
Lasius (C.) umbratus aphidicola (Walsh)	12	35.3		4	1		c		
Le (A.) Jatipes (Walsh) L. alienus americanus Drame	3	8		2 (1)	t 00		r a	v 0 †	
Formica obscuripes Forel	0-	2.0		~	<u>م</u> د		-	n v	
r. fusca Linnaeus	•			N	Ø		67	10	
F. fusca arrentea Wheeler	Ч	2.9					-1	S	
Fe (Ne) Palliderulva nitidiventris Enery FORMICINAR	Ч	2.9		~	ω		(*	ч г	
Aphaenogaster (A.) subterranea valida Wheeler	54 1	10°4	9	14 5	9.	ŝ	16,	18	`0
<u>Solenopsis</u> (D.) molesta validiuscula Emery MYRMICINAR	80	3.5		-1 60 -1 60	t n		~	и Г	
Tapinoma sessile (Say)	α η	1. v v o	Ч	50	100	~) m	12	н
DOLLCHODERINAE	2 02	5	Ч	NO	an an	~	-	S	
TOTATS		•		2	>	-1	4	n	-1
	34 9	9.8	Ø	25 10	0	8	20	100	8

1

÷

TABLE XIV

ANT SUBFAMILY COMPOSITION

;

.

3

			FOR	ICINAE		L 13			MYRM Col	ICINAE				A	OLICHO	DERINA	E	
		.965	4	996	H.	67	H	965	П	966	F	67	-	965	19	99	15	67
PL AR	- CNI	X	NO.	%	No.	%	ė N	×	No.	X	No.	z	No.	R	No.	R	No.	82
3	53	69.4	38	23	37	2.49	17	20.7	1	21.12	18	31.4	ω	2.6	6	5.7	8	3.4
53	113	55.3	87	58.4	88	53.2	66	32.2	去	36.3	67	40.6	26	12.7	ω	5.4	OT	\$
3	57	70.4	74	56	16	80	8	23.5	6	36	ŝ	15	2	5.9	01	8	-	Ś
Colony Totals	13		139		THI	et î	16		42		88		36		13		13	

i

10

• (

TABLE XV ESTIMATION OF ANT COLONY SIZE BY THE LINCOLN INDEX

;

Colony	Markød First Dav	Recaptured Marked	second day Ilmurked	Colony density	Colony density
			N141441 1000	YADUT AA	by Agruat Count
Camponotus (T.) vicinus	50	4	56	750	60
Camponotus (T.) Vicinus	25	rt -	н	50	62
Camponotus (T.) vicinus	30	6	19	93	142
Camponotus (T.) vicinus	50	S	ω	130	273
Formica (N.) pallidefulva Tatreille	25	Ø	16	75	211
Formica (N.) pallidefulva Latreille	011	9 1 9	7	113	149
	٦.	: 2			

MORNING	TEMPERATURE	READINGS
---------	-------------	----------

	Time	Area	Subsurface	Surface	6" above	51 above	Clear
Date	A.M.		2-3" below		ground	J' accive	SKy
					E O CU PI	Eroun	condition
7/29	8:00	01	65.5	67 9	00 0		
		02	62.0	61	10.0	77.8	Clear
		03	(7.0)	04-5	73.1	75.0	
		(J)	07.0	67.9	77.9	78.3	
-							
7/30	8:00	QI	63.2	68.5	80.5	81.4	Clear
		Q2	65.5	76.4	80.9	87.5	04.002
		Q3	67.9	70.6	28.3	20.0	
				10.00	10.)	(7.01	
7/31	7:00	01	63.5	10 2	1.6 -	1.6	
		02	63 6	47.03	40.5	46.5	Overcast
		00	01.9	55.7	48.6	48.2	
		63	05+5	52.8	49.5	48.5	
0/2	1					-	
8/1	6:30	QI	59.8	57.5	54.5	55.5	Ptly Cldy
		02	59.5	55.0	52.9	52 /1	roth orth
		Q3	62.4	54.5	57 6	57 5	
				1.07	Juco	51.5	
8/2	6:30	01	61.6	62 6	11 -		
		02	6h o	03.0	00.3	67.8	Ptly Cldy
		00	04.4	62.9	69.3	68.5	
		6	66.5	63.6	62.3	63.9 .	
0/0							
013	6:30	Q1	61.8	60.9	61.0	61.9	Ptly Cldy
		Q2	63.4	60.7	60.1	60.0	. ord ord
	-11	. 03	66.1	63.5	62.4	62 1	
				-2-2	~~~~	UCOL	
8/4	6:30	C1	61.3	58 1	67 2	10 11	
		02	67.0	57 6	(2.6)	102.0	Fair
		03	65.0	21.0	01.04.0	63.5	
		0	03.4	01.0	61,1	64.6	
ale	6.70	07	10 .				
015	0:13	QL	68.5	66.4	64.3	70.0	Fair
	t	Q2	64.9	63.8	65.1	- 69.0	
		Q3	66.9	61.3	60.5	66 0	
						0.0	
8/6	6:30	01	63.8	63.0	66 7	60 0	
		02	66.0	6h E	66 0	00.0	Fair
		03	67 3	67	00.4	08.5	
		~	07.1	01.5	62.0	65.9	
8/7	6.45	07	11. 1				
977	0142	QL	04.0	64.2	69.3	71.0	Fair
		Q2	66.3	70.0	71.8	72.2	
		Q3	68.0	69.8	70.0	72.5	
						1~0)	
8/8	6:30	Ql	63.9	64.1	65 4	66 3	D13
		02	66.3	64.0	60.0	1.00	rely cldy
		03	66.0	64 0	03.2	05.9	
			00.9	04.0	00.1	67.2	

MORNING TEMPERATURE READINGS

Date	Time A.M.	Area	Subsurface	Surface	6ª above	5' above	Sky
					REGUNA	ground	condition
8/9	7:00	Q1	63.9	63.0	63.8	64 2	(7) and m
. *		02	66.0	64.9	63.8	Gla la	croudy
• •		03	67.9	67.2	66.5	65.4	- 9
						0):+	
8/10	7:30	01	60.1	57.4	55.1	54.2	Rain
1		02	61.1	59.0	55+9	55.0	
		93	62.9	60.8	56.7	55.1	
8/11	8:00	01	59.1	55.7	53.5	53.0	Cl ouder
		G2	59.3	56.1	54.3	53.3	CTOROY
		93	61.4	59.1	55.9	54:2	
0/20	_					2.000	
8/12	7:30	QI	60.1	61.5	62.3	63.4	Fair
1		02	60.8	57.7	61.5	63.1	
		3	62.2	62.2	65.6	66.9	
8/13	7:30	QI	61.1	62.5	64.8	66.5	Fair
		Q2	61.6	58.1	62.0	64.1	A CANA
		C3	62.0	61.9	66.2	. 67.1	
8/14	7:30	Q1	60.4	59.1	50.3	50 7	()]
		02	62.4	60.0	59.3	59.1	croudy
1		Q3	63.9	62.3	59.0	56.4	
8/15	7:30	01	58.7	57.2	62.9	di. L	-
		·Q2	59.0	53.8	60.2	60 4	Fair
		93	59.1	59.5	66.4	67 0	
				5705	0044	0/+0	
8/16	7:30	01	59.5	67.1	67.7	67.8	Fois
۰ ^۱		Q2	61.7	60.2	62.4	64.6	A Chidad
· · ·		93	62.0	59.8	68.9	69.0	
8/17	7:30	01	59.5	57.3	57.8	27 0	D41
• • • •		92	61.2	58.1	56.8	567	LETA CITA
		93	62.7	60.6	59.5	60.0	
8/18	7:30	01	58.2	63.7	61. 0	(1	
		02	60.3	50 0	60.0	04.9	Fair
		03	62.2	62 7	64.0	61.6	
		~	and l	UGAL	04+9	00.2	

. .

MORNING TEMPERATURE READINGS

	Time	Area	Subsurface	Surface	6" above	51 abov	a Skor
Dat	e A.M.	and the second se	2-3" below		ground	ground	condition
0/7						and a state of a state of a state of a	
0/1	9 7:30) Q1	59.1	58.8	59.8	59.7	Fair
		Q2	61.0	57.5	58.0	58.7	
		Q3	62.6	60.9	59.9	60.1	
0/0							
0/2	0 7:15	QI	60.1	64.3	66.2	67.5	Ptly Cldy
		Q2	63.1	64.4	67.0	69.8	
		23	63.3	65.0	67.5	71.1	
8/2	1 7:00	01	67.8	67 h	60.0	-	
		02	65.0	66 3	60.7	70.0	Fair
		03	65.9	68 0	69.7	69.9	
		~	0)0)	00.0	07+7	70.6	
8/22	?:15	Q1	60.9	60.9	61.7	61 D	Dial of the
		Q2	64.1	61.8	62.6	64. 7	rely clay
		Q3	64.8	63.0	65.2	66 5	
01-0					-) - 1		
8/23	7:15	QI	59.8	58.4	57.4	57.3	Rain
		Q2	62.1	55.9	55.6	55.0	LOAL
		Q3	62.7	60.3	58.9	58.8	
8/24	a.h.e	~					
0/24	7:45	Q1	57.7	57.1	65.1	63.0	Fair
		02	61.2	59.5	58.7	59.4	
	-1 >	03	59.8	59.9	60.3	59.2	
8/25	7.15	07	rP O	(2.2			
0/20	1123	02	50.9	61.3	64.4	64.9	Fair
		03	62.2	59.8	60.7	61.6	
			02.02	62.7	64.4	66.2	
8/26	7:30	Ql	60.2	65.7	66 7	110	
		Q2	63.3	62.7	66 /	60.9	rly Clay
•		Q3	63.6	66.8	69.2	60.3	.*
					09.2	09.7	
8/27	7:30	Q1	61.2	70.0	70-6	77.0 0	Touder
		Q2	64.9	70.1	71.5	72.1	Tourth
		Q3	66.6	69.7	71.0	71.2	
8/28	7.20	07	10 1				
0/20	1120	Q1	60.6	63.2	64.1	63.2 F	air
		02	03.1	61.8	60.7	61.1	
		S.	03.8	63.7	63.2	63.5	

- (

TABLE XVII

NOON TEMPERATURE READINGS

	and the second se	Time	Area	Subsurface	Surface	6" above	5' abov	e Sky
,	Date	P. M.		2-3" below		ground	ground	condition
	-1							
	7/29	12:00	Q1	66.1	85.5	86.2	85.6	Ptly Cldy
			Q2	67.9	86.2	91.5	90.5	real ord
			Q3	74.7	84.5	86.3	86 4	
	i de la composición d					000)	00.4	
	7/30	12:00	01	66.1	80.5	87 7	07 m	(7)
			02	67.2	82.4	SE IL	OL O	croudy.
			03	74.5	00.0		04.2	
					1409	00.4	03.2	
	7/31	12:00	01	59.9	50 0	60 1		
			02	62.3	57.7	00.4 ro.0	57.9	Ptly Cldy
			03	66.0	59.0	59.8	60.9	
				00.9	00.9	65.8	62.5	
	8/1	12.00	07	Ch r	(0.0	-1 1		
	-1-	****	02	67.0	09.9	74.4	74.8	Fair
			22	03.2	77.2	78.8	77.9	
			63	70.5	80.5	80.4	75.6	
	0/0	70.00	~					
	0/2	12:00	QI	64.3	70.1	75.8	75.1	Ptly Cldy
			Q2	65.9	78.4	77.2	77.9	
			Q3	67.9	79.9	82.4	80.1	
	010							
	8/3	12:00	Q1	64.0	72.3	72.7	72.6	Cloudy
			Q2	67.9	72.9	72.1	72.2	varung
	`		Q3	70.3	81.3	78.2	75.5	
	- 1-					•	()-)	
	8/4	12:30	01	65.2	76.2	81.3	182.5	Fain
			Q2	68.9	81.7	84.0	84.2	L'att.
			Q3	75.4	86.4	88.3	84. 2	
						000)	UT & L	
	8/5	12:00	Q1	64.7	75.9	77.3	87 ¢	CT and
	۰.	t	Q2	68.6	80.2	87 1	S 04 0	cronal
			03	72.1	83.8	85.0	00.0	
					0,00	07.7	07.0	
	8/6	12:00	01	65.1	76.1	70 6	03 3	
	•		02	68.9	85.2	(7.0	81.1	Ptly Cldy
			03	73.3		07.1	85.9	
	:		4.7	69.63	90.0	02.0	84.9	
	8/7	12:00	01	65 8	776 6	80.0		
	-11		02	60 /	70.0	80.9	83.1	Ptly Cldy
			03	07.4	04.2	86.3	87.8	
				()+1	72.3	54.9	89.6	
	8/8	12.00	07	GE E	00 h	01 -		
	-10 .	~~ 100	02	60.0	02.4	84.7	83.5	Fair
			02	07.7	87.0	85.6	85.2	
				78.0	97.2	96.9	88.8	
_								

TABLE XVII (Cont'd.)

MOUN IESTERAIURE READINE	NOON	TEMP	ERATURE	READINGS
--------------------------	------	------	---------	----------

	an a	an de la ser de la ser de la ser	Subsurface	ad - Brennington pur pri dit-the dag	6" abova	51 abova	Sky
Date	Time	Area	2-3" below	Surface	ground	eround	condition
						and the second second second	
8/9	12:30	QL	63.2	64.5	63.2	62.5	Detala
		02	65.9	62.9	62.0	67.6	UI 20020
		03	68.0	68.2	64.0	62.3	
						Viii Bab	
8/10	12:30	01	60.8	57.0	54.9	54.7	Pain
•		02	61.5	59.9	56.5	ch. 3	110.411
		03	63.6	60.2	56.7	55 5	
			-,		1002	2202	
8/11	11:30	01	59.1	57.4	56.7	55 E	Orrows
		C2	60.0	58.8	53 6	27.2	Asleast
		03	63.5	62.0	63 7	21+2	
				OL + V	ULeL	00.9	
8/12	12:00	01	61.8	72 7	75 3	06 0	D12 02 1
		02	65.4	76 0	1202	10.1	hera craa
		03	67.0	70.0	()=4	13.4	
			0169	11.3	OT*O	01.0	
8/13	12:00	C1	62.3	72 1	24 0	PH. O	
-//		02	64.2	77 3	1990 D	74.0	Farr
		03	69 h	77.0	10.2	78.5	
		*)	00.4	13.0	13.1	14+3	
8/14	12:00	07	63.7	62 3	62.0	100	A
		02	63.8	Ch 7	64.0	02.3	Overcast
		03	66 6	65 3	64.2	03.2	
-		4)	00.0	03.1	00.1	05.9	
-8/15	17.30	-01	50 8	66 12	60 1	50° 0	
-1-2		.02	62.6	69.2	Dy.I	1 70.0	rair
		03	64 0	00.3	70.1 ·	09.9	
			03.7	()+4	14.3	72.8	
8/16	11:30	03	60.8	60.2	677 7		-
-/	anal Jo	02	64.3	49.3	71.1	70.2	Fair
	•	03	65 1	00.4	70.4 3	72.2	
		~)	00.4	(1+2	73.1	72.3	
8/17	11.30	ó	60 7	66 E	60.0	1	
		02	63 6	20.0	60.7	07.1	Estly Fair
		03	69.7	70.0	Oyed.	07.9	
		a.J	0701	(U+T	1406	0.80	
8/18	12:00	01	62.2	211 3	75 3		
-1		02	63 5	(7+)	()=3	70.7	Astly Fair
		03	60 4	13.1	13.4	73.7	
		~)	07+0	10.3	70.3	78.1	

NOON TEMPERATURE READINGS.

			Subaugaca		68 -	F1 -	
Data	174-100	1	Subsuriace	C	o" above	5' above	Sky
Daus	1 Line	Area	C-1. DOTOM	Surrace	ground	ground	condition
8/19	12:30	67	61.3	72 7	27 2	00.1	Maka Data
-/-/		02	69.0	72 5	72.7	13.L	mstly fair
		03	27 1	(4.)	14.1	72.4	
		~)	(1.44	19.9	01.0	80.9	
8/20	11:30	01	62.1	77.3	28.3	77 0	Delles al and
		02	64.9	80.3	81 1	01 0	LUTA CTORY
		03	66 6	77 9	01.1	01.2	
		~)	05.0	11.0	02.0	82.3	
8/21	12:00	01	65.6	75.0	76.9	78.0	Matler Date
		02	66.8	76.3	78.8	80.2	Escry Larr
		03	72.5	86.2	81.0	00.5	
		\sim	(20)	00.2	04.0	00.0	
8/22	12:30	01	63.4	75.5	77.4	77.6	Phly Clay
		22	68.3	76.4	77-1	78 0	rory ordy
		23	20-8	85.7	Rit 7	22.0	
		~	1000		04.7	03.0	
8/23	12:30	01	61.0	66.6	67.3	68.1	Fate
		02	67.4	6611	68.0	66.3	7. CI-4-7
		93	69.8	75.1	20.0	20.5	
		-	.,	1 202	[0+]	(1)	
8/24	12:15	01	59.8	69.2	69.5	67.0	Fair
		Q2	60.7	68.1	68.0	68.8	& GLANE
	·: ` `	23	69.3	76.3	75.7	68 0	
		~		,,	13+1	00.9	
8/25	1:00	Q1	61.3	70.0	77.2	29.3	Ptly Cldy
		02	68.2	87.2	86.2	80.3	- and arad
		23	72.6	86.8	87.2	84.3	
			1	0010	01 420	04.)	
8/26	12:00	QL	62.1	80.6	80.4	79.9	Ptly Clay
	ı	22	66.3	82-8	84.3	82.2	rows origh
		23	69.7	89.9	85.6	97 3	
			4741	0707	03:0	01.)	
8/27	11:45	Q1	62.3	82.2	81.0	79.1	Ptly Clay
		22	65.3	84.2	82.2	81.8	rowy oray
		23	69.4	78.8	78.3	81.4	
- 1	_:						
8/28	12:00	Ql	62.4	72.6	76.5	76.7	Ptly Cldy
		Q2	68.5	77.4	78.8	79.3	
		23	70.0	81.4	81.8	77.7	
					~ m c V	(1•1	

. (

TABLE XVIII

AFTERNOON TEMPERATURE READINGS

			Subsurface		6ª above	5' above	Sky
Dat	e Time	Area	2-3" below	Surface	ground	ground	condition
7/20	0 1.20	07	(0.5				
114	7 4190	Q1	03.7	74.3	75.1	74.4	Ptly cldy
		Q2	00.5	77.2	75.5	74.5	
		Q3	75.2	78.3	77.5	75.9	
7/30	4:30	01	70 4	69. E	10 4	-	
112		02	67.8	40 r	09.5	70.1	Hd rained
		03	72 1	60.5	00.2	67.5	
			1204	01.7	07.5	70.8	
7/31	4:30	Q1	60.0	59.1	57.0	57.5	Data
		Q2	64.5	60.1	55.5	53 5	Rolli
		Q3	68.5	67.0	64.5	63.4	
				-,	04.5	03.4	
8/1	5:30	¢1	66.4	69.3	71.5	72.1	Ptly cldy
		Q2	68.8	70.3	71.2	72.3	
		Q3	73.5	73.2	73.1	74.4	
8/2	5:20	01	66.5	69.9	76.0		B . 1.
		02	67.7	77.3	70.2	11.4	Fair
		03	73.0	81.0	17+L	79.8	
			()+)	01.9	02.1	82.1	
8/3	5:30	Q1	66.0	65.5	62.8	65.3	Ptly cldy
		Q2	65.3	63.9	67.2	68.2	they they
		Q3	70.5	68.0	69.2	69.9	
8/4	5:00	01	67.5	83.1	82.9	100.0	D4.7
•		02	73.4	84.7	02.0	02.9	Ptly cldy
		03	74.7	27 7	03.0	03.0	
		47	1.7+2	CHEEL	0)=4	85.0	
8/5	4:45	Q1	66.8	76.1	78.3	78.8	Cloudy .
		Q2	68.3	77.9	78.0	77.8	erond .
		Q3	72.9	79.0	79.7	81.6	
8/6	5.00	01	11.0				
0/0	5:00	00	00.0	80.5	81.8	83.1	Ptly cldy
		62	12.5	83.0	81.9	81.9	
:		QJ	18.9	52.4	83.0	82.7	
8/7	4:45	Ql	70.2	79.0	77.9	77 6	Dela alda
		Q2	70.0	78.6	78.3	77 0	eery croy
		Q3	75.3	82.2	82.5	82.6	
8/8	5.00	07	60 0	-	-		
010	5100	02	60 1	70.3	76.1	77.9 1	ld rained
		03	Cy .4	71.1	77.0	78.3	
		\$	14.0	19.7	82.1	83.2	

.(

AFTERNOON TEMPERATURE READINGS

and fighter state		10 101-111-0-10 -10-	481-2424-444-7446-444-444-444-4	Subsurface		6" above	51 above	Sky
D	ate	Time	Area	2-3" below	Surface	ground	ground	condition
						and the state of t	1999-1999 1999 1999 1999 1997 1998 1996 1996 1996 1996 1996 1996 1996	
8	19	5:00	Q1	63.4	64.1	63.0	61.2	Rain
			02	65.7	63.2	63.1	60.3	
		1	23	68.4	68.9	65.1	61.6	
8	/10	5:00	Q1	60.0	57.1	54.4	57.2	Rain
			Q2	60.8	59.3	55.6	52.4	
			Q3	61.3	59.6	56.0	53.1	
8	/11	5:00	01	61.1	66.7	67.2	68.9	Ptly oldy
			Q2	65.4	66.5	67.8	69.1	roth croh
			03	66.9	66.1	63.4	60 0	
						0.78.7	07.7	
8	/12	4:45	01	63.7	66.0	68 E	77 2	Cl andre
			02	68.0	20.3	70.8	()	croudy
			03	72.5	21 2	72.0	(207	
			4)	1000	1204	1607	14.0	
8	/13	4:30	C1	64.2	66 1	70 E	70 0	CTA made
- 1			02	67 3	20.1	(0.)	12.0	Sit rain
			03	20.0	70.3	(1.)	().0	
			4)	1007	14.3	16.L	.14.1	
8	174	4.45	01	62 2	60 3	=6 0	100 0	Della
~,		1.19	02	66 2	60.2	50.9	51.3	Rain
	5		03	69 0	67.0	50.0	50.1	
			(),) (),)	00.0	2.7	02.3	04.3	
- 8	175	6.30	01	62 11	62 0	60 0	10: 0	*** 1
-,	-	0.00	102	67 2	66 6	6 2	1 00.0	Fair
			03	60 3	67.0	60.J	09.1	
				09.0	07.0	00.7	69.9	
8	176	5.30	67	63 0	70 0	70 P	00 0	T . 1
	~~~	1.00	02	69 5	10.7	14.1	13.9	Fair
			03	72 2	1.1.5	12.1	71.8	
				1606	(2)	75+1	76.3	
8/	177	5.00	01	62 7	72 2	22 0		
~1		2.00	02	60 0	14.6	1204	74.0	beth crah
			03	72 9	10.7	70.4	73.9	
÷			40	(2.0	0.1.0	79.9	78.8	
RI	18	5.30	07	61 0	73 4	00.0	500 C	
~	alu al	200	02	68 0	1.1.04	12.U	12.3	Fair
			12	00.9	1202	12.4	73.7	
			w.)	1.2.07	13.2	10.5	70.9	

. 4

AFTERNOON TEMPERATURE READINGS

_			Subsurface		6" above	5' above	Sky
Date	Time	Area	2-3" below	Surface	ground	ground	condition
8/19	5:00	01	62.1	72.2	73.8	75.2	Ptly oldy
-1		02	68.9	73.3	74-1	74.2	rang chuy
		93	73-9	76.1	78.0	77.7	
0.1-0							
8/20	5:00	QI	64.2	76.2	77+3	78.3	Fair
		Q2	70.9	81.4	80.2	79.8	
		Q3	74.1	79.1	81.0	84.9	
8/21	5:30	Cl	64.2	72.4	73.8	74-3	Ptly oldy
		02	70.1	72.3	72.9	73.0	rong crub
		¢3	74.7	74.9	73.3	73.9	
0/00	F.00	50	(1) 7	-			
0/24	3:00	02	04.1	73.2	73.9	73.5	Ptly cldy
		62	1302	77.3	75.2	75.9	
		63	74.4	75•7	75.8	76.0	
8/23	5:45	Q1	61.6	65.0	65.8	66.1	Fair
		02	68.3	67.0	66.8	66.8	
		Q3	70.2	70.4	70.0	69.2	
8/24	5:15	01	60.9	63.7	63.2	60.7	Foin
-,~		02	68.5	69.6	20.0	60.0	L.C.T.
-		Q3	72.3	74.8	76.1	75.1	
8/25	5.20	07	(n P	(m ).		-1.0	
0/25	2:20	02	62.1	07.4	74.3	174:8	Ptly cldy
		62	09.4	72.1	74.4	75.0	
		43	12+9	73.9	73+5	74.7	
8/26	5:00	01	63.7	76.4	77.9	78.3	Ptly cldy
	t.	Q2	69.8	80.4	80.0	80.1	
		03	72.8	81.2	79.8	80.4	
8/27	4:30	C1	64-1	72.5	72 8	א רס	Clouder
		02	70.7	72.5	20.9	70 1	CTORDA
		03	72.0	73.3	73.1	72.3	
-				12-2	1.2.4.44	1.063	
8/28	5:00	Ql	63.1	66.5	70.3	72.2	Ptly cldy
		Q2	63.6	72.2	71.4	72.6	
		63	77 7	MC 6	n6 h		

Date	Area	Subsurface	Surface	6" above ground	5' above ground	
7/29-8/4	Ql	62.4	60.8	64.1	64.8	
	Q2	62.8	61.8	63.8	64.4	
	Q3	65.8	62.1	63.3	64.1	
8/5-8/11	Q1	63.4	62.0	62.5	63.8	
	Q2	64.3	63.2	63.2	64.0	
	03	65.9	63.5	62.5	63.8	
8/12-8/18	Q1	59.6	61.1	62.9	63.4	
	Q2	61.0	58.3	60.4	61.7	
1	Q3	62.1	61.2	64.4	64.8	
8/19-8/25	C1	59.8	61.2	63.5	63.8	
	02	62.7	60.7	61.8	. 62.8	
	Q3	63.0	62.8	63.7	64.6	
8/26-8/28*	01	60.7	66.3	67.1	67.0	
	02	63.8	64.9	66.2	67.2	
	03	64.7	66.7	67.8	. 68.1	
	ι		.::	10	1	

-

#### TABLE XIX

WEEKLY MORNING TEMPERATURE AVERAGES IN ALL QUADRATS

*For only a three day period.

. 1

;

Date	Area	Subsurface	Surface	6" above ground	5 ¹ above ground
7/29-8/4	61	64.3	73 E	76 0	01 0
11-1-01+	02	66.2	76.8	70. Ju	12.1
	03	77.5	87 8	97 /1	(0.)
		(1.)	OT.O	01+4	10.2
8/5-8/11	Q1	63.5	70.0	71.0	71.6
	02	66.3	74.0	73.6	74.0
	Q3	70.5	79.5	77.9	75.7
8/12-8/18	01	61.4	69.0	70 0	20 2
0/20-0/20	02	63.9	71.0	10.9 m c	10.1
	03	67 7	72 5	(10)	(1.)
		01.1	()0)	(4+)	13.2
8/19-8/25	01	62.1	72.2	74.0	74-4
	Q2	65.0	75.3	76.0	75.3
	Q3	70.4	81.0	81.0	78.8
0/0/ 0/00*					
8/25-8/28	Q1	62.3	78.5	79.3	78.6
	Q2	67.0	81.5	81.8	81.1
	03	69.7	83.4	81.9	80.1
				11	

;

### WEEKLY NOON TEMPERATURE AVERAGES IN ALL QUADRATS

TABLE XXI

*For only a three day period.

Area	Subsurface	Surface	6 [#] above ground	5 [•] above ground	
<b>G1</b>	65-8	20.0	70 7	77 h	
02	67.7	71.6	21 2	72.4	
03	72.9	73.9	74.0	74.5	
QL	65.1	70.5	71.2	71.2	
Q2	67.4	71.4	71.7	71.0	
Q3	71.1	73.8	73.8	73.5	
Q1	62.9	67.4	68.7	69.9	
Q2	67.2	69.6	69.8	69.9	
93	71.1	71.0	72.5	73.5	
Q1	62.8	69.9	72.4	73.1	
Q2	69.9	73.3	73.5	73-5	
Q3	73.2	74.9	75.4	75.9	
01	63.6	71.8	23.7	71.0	
' 02	69.7	75-0	74.7	31 0	
03	72.0	76.7	76.4	. 76.7	
	Area Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q3 Q1 Q2 Q3 Q1 Q2 Q3 Q3 Q1 Q3 Q3 Q1 Q3 Q3 Q3 Q1 Q3 Q3 Q3 Q3 Q3 Q3 Q3 Q3 Q3 Q3	Area         Subsurface           Q1         65.8           Q2         67.7           Q3         72.9           Q1         65.1           Q2         67.4           Q3         71.1           Q1         62.9           Q2         67.2           Q3         71.1           Q1         62.8           Q2         69.9           Q3         73.2           Q1         63.6           Q2         69.7           Q3         72.0	AreaSubsurfaceSurfaceQ1 $65.8$ $70.0$ Q2 $67.7$ $71.6$ Q3 $72.9$ $73.9$ Q1 $65.1$ $70.5$ Q2 $67.4$ $71.4$ Q3 $71.1$ $73.8$ Q1 $62.9$ $67.4$ Q2 $67.2$ $69.6$ Q3 $71.1$ $71.0$ Q1 $62.8$ $69.9$ Q2 $69.9$ $73.3$ Q3 $73.2$ $74.9$ Q1 $63.6$ $71.8$ Q2 $69.7$ $75.0$ Q3 $72.0$ $76.7$	AreaSubsurfaceSurfacegroundQ1 $65.8$ $70.0$ $70.7$ Q2 $67.7$ $71.6$ $71.2$ Q3 $72.9$ $73.9$ $74.0$ Q1 $65.1$ $70.5$ $71.2$ Q2 $67.4$ $71.4$ $71.7$ Q3 $71.1$ $73.8$ $73.8$ Q1 $62.9$ $67.4$ $68.7$ Q2 $67.2$ $69.6$ $69.8$ Q3 $71.1$ $71.0$ $72.5$ Q1 $62.8$ $69.9$ $72.4$ Q2 $69.9$ $73.3$ $73.5$ Q3 $73.2$ $74.9$ $75.4$ Q1 $63.6$ $71.8$ $73.7$ Q2 $69.7$ $75.0$ $74.1$ Q3 $72.0$ $76.7$ $76.4$	AreaSubsurfaceSurfaceSurface $ground$ $ground$ Q165.870.070.771.4Q267.771.671.271.4Q372.973.974.074.5Q165.170.571.271.2Q267.471.471.771.0Q371.173.873.873.5Q162.967.468.769.9Q267.269.669.869.9Q371.171.072.573.5Q162.869.972.473.1Q269.973.373.573.5Q162.869.972.475.9Q163.671.873.774.0Q269.775.074.131.9Q372.076.776.476.7

# WEEKLY AFTERNOON TEMPERATURE AVERAGES IN ALL QUADRATS

TABLE XXII

*For only a three day period.

.(

•

#### TABLE XXIII

## WEEKLY MORNING TEMPERATURE EXTREMES IN ALL QUADRATS

		Subsu	rface	Sur	face	6" abor	re surf	5' abo	ve surf
Date	Area	Max	Min	Max	Min	Max	Min	Max	Min
-las ali									
7/29-8/4	QI	65.5	59.8	68.5	49.3	80.5	46.5	81.4	46.5
	02	65.5	59.5	76.4	55.0	80.9	48.6	81.5	48.2
	03	67.9	65.2	70.6	52.8	78.3	49.5	79.7	48.5
8/5-8/11	01	68.5	59.1	66.4	55.7	69.3	53.5	21.0	53.0
	02	66.3	59.3	70.0	56.1	71.8	54.3	72.2	53.3
	Ø	68.0	61.4	69.8	59.1	70.0	55.9	72.5	54.2
8/12-8/18	01	61.1	58.2	67.1	57.2	67.7	57.8	62.8	57.0
	02	62.4	59.0	60.2	53.8	62.4	56.8	64 6	56 3
	Q3	63.9	59.1	62.3	59.5	68.9	59.0	69.0	56.4
8/19-8/25	01	61.8	57.7	67.4	57.1	69.9	57 14	20.0	67 3
	02	65.0	61.0	66.3	55.9	60.7	66.6.	40.0	56 0
	23	65.9	59.8	68.0	59.9	69.9	58.9	70.6	58.8
8/26-8/28*	01	67.2	60 2	20.0	62 0	<b>20</b> (	(1. )	-	10
0/20-0/20	100	41.0	62 7	70.0	(7.0	70.0	04.1	71.0	03.2
	00	16 2	63.6	/V.L	01.0	71.5	60.7	72.1	61.1
-	~>	0.00	03.0	09.7	03.7	71.0	63.2	71.2	63.5
	L				• ( ⁷	alo	1		

-

*For only a three day period.

• (

i

		Subsu	rface	Surf	ace E	aboy	e surf	51 abo	ve surf
Date	Area	Max	Min	Max	Min	Max	Min	Max	Min
7/29-8/4	Q1	66.1	59.9	85.5	59.9	86.2	60.4	85.6	57.9
	Q2	68.9	62.3	86.2	59.0	91.5	59.8	90.5	60.9
	03	75.4	66.9	90.9	68.9	88.4	65.8	86.4	62.5
8/5-8/11	01	65.8	59.1	82.4	57.0	84.7	54.9	83.5	54.1
	Q2	69.9	60.0	87.0	58.8	86.3	56.5	87.8	54.3
	Q3	78.0	63.5	97.2	60.2	96.9	56.1	89.6	55.5
8/12-8/18	01	63.3	57.8	74.3	62.1	75.3	63.0	76.7	62.3
	02	65.4	62.6	77.1	64.1	78.2	69.1	78.5	63.2
	Q3	69.6	65.9	78.5	65.1	81.8	66.1	81.0	65.9
8/19-8/25	Q1	65.5	59.8	77.3	66.6	78.3	67.3	79.3	67.0
	Q2	69.0	60.7	87.2	66.1	86.2	68.0	81.2	66.3
	Q3	72.6	66.6	86.8	75.1	87.2	70.9	84.3	68.9
*									
8/25-8/28	01	62.4	62.1	82.2	72.6	81.0	76.5	79.9	76.7
~~	Q2	68.5	66.3	84.2	77.4	84.3	78.8	82.2	79.3
-	Q3	70.0	69.4	81.4	78.8	81.8	78.3	81.4	77.7
undAuto Britishiginkal na finanza a na panya	L				κF	do	. 1		

#### TABLE XXIII

### WEEKLY NOON TEMPERATURE EXTREMES IN ALL QUADRATS

*For only a three day period.

.(

;

#### TABLE XXIV

WEEKLY AFTERNOON TEMPERATURE EXTREMES IN ALL QUADRATS

		Subsur	face	Suri	ace 6	ó* abor	surf	51 abo	ve surf	-
Date	Area	Max	Min	Max	Min	Max	Min	Max	Min	
						\ \		and a second	1421-4994-45-49-49-49-49-49-49-49-49-49-49-49-49-49-	
7/29-8/4	01	70.4	60.0	83.1	59.1	82.8	57.0	82.9	57.5	
	Q2	73.4	64.5	84.1	60.1	83.8	55.5	83.8	53.5	
	03	76.2	68.5	81.9	67.0	83.2	64.5	85.0	63.4	
8/5-8/11	01	79.2	60.0	80.5	57.1	81.8	54.4	83.1	51.2	
	02	72.5	60.8	83.0	59.3	81.9	55.6	81.9	52.4	
	03	78.9	61.3	82.4	59.6	83.0	56 0	82.2	62 1	
	~	1007	~~~ V J	UL20T	3700	0).4	20.0	Olek	22+7	
8/12-8/18	Ql	64.2	61.9	72.2	60.1	73.2	56.9	74.0	57.3	
	Q2	69.9	61.3	76.9	60.3	75.4	56.6	73.9	56.1	
	Q3	72.8	68.0	81.0	61.9	79.9	62.3	78.8	64.3	
8/19-8/25	01	64.2	60.9	76.2	63.1	77.3	65.8	78.3	66 7	
	02	73.2	68.3	81.4	67.0	80.2	66 8	- Dd 9	66 9	
	03	74.4	70.2	20 1	70 4	01 0	00.0	(7.0	00.0	
	1	1-10-4	10.2	1701	10.4	01.0	70.0	84.9	09.2	
8/26-8/28*	07	64.7	62.3	06 h	11 1		-	-		
0/200/20	, 00	Totol C	(0 (	10.4	00,5	77+9	70.3	78.3	71.4	
54 E	42	10.7	00.0	80.4	72.2	80.0	70.9	80.1	70.4	
	63	72.8	71.1	81.2	73.3	79.8	73.1	80.4	72.3	
•							11			
	1				.:1		2		-	

*For only a three day period.

.

t

;

#### TABLE XXV

#### U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS STATION-BOULDER, COLORADO TIME OF OBSERVATION 5:30 P.M.

June		gato	oservation	Precipitation S. Boulder	
And some third, it is seen to be a second ball the second s	Max.	Min.	at Obsn.	Ins & Hundredths	
	a É				
1	75	57	62	.15	
2	77	48	68		
3	77	50	70		
4	75	51	51	.30	
5	62	44	57	1.20	
6	80	43	23		
7	80	60	74	-03	
8	74	49	61	- 06	
õ	74	5/L	69	T	
10	60	56	56		
11	67	57	67	24	
7.0	10	21	67	• 20	
12	71	21	OT 0	1	
13	02	20	12	-01	
14	79	58	58		
15	77	56	76		
16	77	56	61	.10	
17	76	55	70	•25	
18	78	55	75	T	
19:	85	57	81		
20	83	55	76		
21	- 81	53	80	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
22	. 89	65	72	The second se	
23	82	59	82	상업 전 전쟁이 많은 것이 많이 많이 많이 했다.	
24	82	55	79	.06	
25	82	59	75		
26	85	55	80		
27	80	59	79		
29	Oh.	50	0		
20	04	22	11		
29	83	59	05		
30	81	62	75	.10	
July				Sum 2.66	
· 1	89	58	85		
2 :	85	57	78		
3	88	54	87		
4	90	60	83		
5	89	61	85		
6	90	58	83		
		~			

.1

### TABLE XXV (Continued)

#### U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS STATION-BOULDER, COLORADO TIME OF OBSERVATION 5:30 P.M.

Date	Temp.	Boulde	r 24 Hrs.	Precipitation	
1965	Endin	g at Ob	servation	S. Boulder	
July	Max.	Min.	at Obsn.	Ins & Hundredths	
7	00	67	00		
8	80	65	81	(Th	
a	86	58 -	81	25	
10	Ol	50	80	• <i>2</i> )	
11	80	62	00	1	
12	07	60	00	10.	
12	07	05	04	.02	
1)	04	22	OT .	T	
14	89	54	87		
15	92	61	87		
10	94	65	89		
17	90	62	84		
18	90	67	87	·. ·	
19	88	65	81	•32	
20	86	60	81	T	
21	87	54	84	.75	
22	92	62	86	.49	
23	83	63	73	T	
24	74	60	69	1.07	
25 '	77	57	72	•98	
26	83	57	80	.10	
27	76	61	64	T	
28	L 80	55	71		
29	87	60	85	안정 이 전쟁에서 이 것이 없는 것이 없다.	
30	86	61	81	-28	
31	67	56	57	84	
1	-1	10	21	Sum 5.71:	
				Jean Jean	

;
### TABLE XXVI

### U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS STATION-BOULDER, COLORADO TIME OF OBSERVATION 5:30 P.M.

Date	Temp.	Boulds	r 24 Hrs.	Precipitation	an a the second seco
1966	Ending	; at Ob	servation	S. Boulder	
August	Max.	Min.	at Obsn.	Ins. & Hundredths	
			•		
1	81	66	67	.16	
2	84	67	67	.25	
3	84	60	75	.02	
4	81	59	77	.04	
5	87	54	82	. •	
6	92	61	82		
7	84	57	79		
8	79	56	65	-	
9	77	58	74	T	
10	82	54	78	T	
11	95	56	86		
12	89	60	69	T	
13	85	50	85	•	
14	90	61	83		
15	87	55	82	1	
16	96	63	94		
17	94	73	74		
18	87	56	82	•	
19	85	66	67	• 04	
20	. 77	57	72	· · · ·	
21	14	52	70	an do the	
22	1 70	40	74		
2)	70	26	71	Ŧ	
25	87	40	80	*	
26	07	27	00	3	
27	02	56	0/		
28	86	63	72		
29	88	58	69	02	
30	89	55	68	•05	
31	80	58	60	• • • • • • •	
		50	00		
;				Sum •13	

• ٩

## TABLE XXVII

### U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS STATION-BOULDER, COLORADO TIME OF OBSERVATION 5:30 P.M.

Date	Temp.	Boulde	r 24 Hrs.	Precipitation	Mate Artistana (sa Sa
Jul w	Mar	g at uo	servation	S. Boulder	
	rake	Pilli.	at Ubsn.	Ins. & Hundredths	
31	88	64	82		
August	80	12	1-		
2	09	01	67	k .	
2	04	00	84	$\mathbf{r} = \mathbf{f}$	
1	04	02	02	.12	
5	00	22	05		
5	94	50	80	T	
2	02	59	72	T	
6	60	59	70	T	
0	LO	66	69		
10	01	61	65	.02	
10	80	53	64	.19	
10	82	52	64		
75	88	55	66	•03	
1)	00	62	83	,· · · · · · · · · · · · · · · · · ·	
14	85	60	70		
15	85	59	75		
TO ,	86	56	64	T	
17	80	59	76	T	
10	79	57	70	T	
19	83	45	75	•76	
20	82	56	70	4.0	
21	85	54	76		
22	18	61	65	T	
23	90	63	87		
24	94	65	75	· · · · · · · · · · · · · · · · · · ·	
25	93	61	78	T	
20	.77	53	54		
27	90	51	77	.10	
28	84	64	72	T	
29	82	52	63	T	
30	69	52	53	1.83	
31	59	48	50	.80	
				Sum 3.85	

. (

### TABLE XXVIII

#### U.S. WEATHER BUREAU RECORD OF CLIMATOLOGICAL OBSERVATIONS STATION-BOULDER, COLORADO TIME OF OBSERVATION 5:30 P.M.

Date	Temp.	Boulde	r 24 Hrs.	Pre	cipita	ation	
1968	Ending	g at Ob	servation	5	. Bou	lder	
July	Max.	Min.	at Obsn	. Ins.	& Hund	dredths	Shartshart securit management of sources and sources and sources and sources and sources and sources and source
29	92	59	81				
30	88	64	66	•	.12		
31	68	50	59		.16		
August				Sum	.28		
1	83	51	80				
2	85	61	83				
3	84	58	75		.02		
4	92	55	84				
5	89	66	81	-			
6	91	61	84				
7	90	64	80				
8	88	64	82		T		
9	83	60	61		.55		
10	62	53	53	all day	1.53	rain &	drizzlə
11	76	51	71		.23		
12	82	51	75		.08		
13	84	54	71		.02		
14	72	56	66		.33		
15	83	53	79		.01		
16	83	54	79				
17	. 81	53	78				
18	83	60	78		1	•	
19	86	54	82				
20	88	54	85				2
21	86	64	75				
22	85	58	78		-		
23	84	16	81		• · · · · ·		
2)	86	50	74				
25	86	50	85				
26	87	51	82				÷.
20	82	67	72				
29	86	53	76		m		
20	25	50	75 59		1 25		
29	()	50	20		•4)		
20	14	60	14		Ŧ		
L	. 12	54	13	Cum	2 02	D-144(205	
				SUM	3.02		

- 1

TABLE XXIX SOLL MDISTURE IN QUADRAT 1 IN 1968

;

11 2	4												
Average		7.61		12.17		12.93		20.15	•		6•63		10.66
& Moisture	6.85	8.05	20.21	10.34	30.02 01 51	12.22	14.07	7-96	12.47	7.33	6.18 6.18	12-11	8.87 11.88
Sample Moisture	12.7	14.41	26.9	16.6	5-12	19.4		15.2	7.62	13.9	12.0	18.7	16.1
Sample	-1	n m	i el	01 00		~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	) -	4 02 0	n	-10	i m	Ч	a m
Dry Soil	185.3	176.6	175.3	160.5	162.6	158.8 160.3	C 771	190.9		190.8	194.3	166.3	181.6 152.4
Sample	-	2 5	~	2 5	-	2			, i	10	m	Ч	2 5
Date	7/31		8/1		8/2		8/3		0.10	+/0		8/5	
Wet Soil	198.0	190.6	202.2	177.1	183.9	178.2	161.4	206.1	6 606	203.0	206.3	185.0	170.5
Sample	40	n m		2	Ч	2	Ч	01 m	ſ	1 01	m	Ч	n m
Date	7/29		7/31		8/1		8/2		8/3			8/4	

TABLE XXIX (Continued) SOIL MOISTURE IN QUADRAT 1 IN 1968

;

• (

late	Sample	Wet Soll	Date	Sample	Dry Soll	Sample	Sample Moisture	% Moisture	Average &
17	Ч	190.9	8/12	н	158.3	Ч	32.6	20.50	
	2	195.9	•	2	160.1	0	34.8	20 00	
	m	194.1		3	157.3	1	36.8	23.39	11-27
								Proper	
/12	r-t	176.0	8/13	rt	156.0	rt	20-0	12.82	
	2	194.4		0	170.0	N	24.4	711 25	211 110
	3	197.2		e	169.6	3	27.6	16.27	0****
/13	Н	198.3	41/8	F	170.0	r	000		
	N	198.9		0	0.17	4 0		C0-0T	
	m	4.102		5	173.4	2	28.0	16.40	E2.71
411	~	6 600	1.10	•					
+	4 (	1.20%	CT/O	-4	1.87L	Ч	29.2	16.40	
	4 0	5-26T		2	163.9	2	28.6	27.45	16.85
	2	C.201		<u>رم</u>	156.2	3	26.1	16.71	
15	Ч	190.6	9T/8	, H	159.8	-	8 06	20.05	
	~	196.8	•	2	173.1	10	000	12024	1
	e	189.3		3	165.6	1 67	23.7	14.31	al.cr
,16	ы	208.5	8/17	н	181.0	-	27.5	01 21	
	~	182.3		~	160.7	0	21.6	13.444	13.04
	n	187.8		m	165.9	e	21.9	13.20	トル・ヘー

TABLE XXIX (Continued) SOLL MOISTURE IN QUADRAT 1 IN 1968

-1

Average & 9.54 5.60 8.58 6.56 6.61 21.93 & Moisture 9.29 8.62 10.70 5.39 2.55 6.58 6.60 27.87 20.71 17.22 Molsture Sample 16.2 15.2 20.1 10.0 14.1 10.8 22.23 41.8 Sample n n m Nm Date Sample Dry Soil 174.4 176.4 187.9 196.6 186.8 192.8 165.1 170.9 177.6 184.3 183.0 192.4 181.3 150.0 m ÿ N 8/6 8/7 8/10 8/8 8/9 11/8 Date Sample Wet Soil 190.6 191.6 208.0 207.2 196.8 204.5 179.2 185.3 193.2 195.8 193.8 206.9 193.6 189.1 188.4 191.8 184.2 193.3 ri N 3 2 3 N 8/5 8/6 8/7 8/10 8/8 8/9

TABLE XXIX (Continued)

. (

SOIL MOISTURE IN QUADRAT 1 IN 1968

••••

Average 2		13.27	14.13	11.84	10.22	11.53	04.6
& Moisture	31.48	16.19	12.47	11.56 10.77 13.18	9.79 11.80 9.08	13.52 9.95 11.11	10.09 9.24 8.86
Sample Moisture	18.8	22.0	22.6 26.6 25.2	20.4 20.4 23.4	18.0 21.8 17.0	22.8 17.7 20.9	18.3 16.7 16.5
Sample	н	N M	101	199	HNM	NNM	HNM
Dry Soll	163.8	181.2	181.3 169.2 177.5	176.5 188.4 177.6	183.8 184.8 187.3	168.6 177.9 188.2	181.3 177.4 186.3
Sample	-10	3	1 2 9	HNM	- 1 a'm	-Harm	ANM
Date	81/8		8/19	8/20	8/21	8/22	8/23
Wet Soil	182.6	203.2	203.9 195.8 202.7	196.9 208.7 201.0	201.8 206.6 204.3	191.4 195.6 209.1	199•6 194•1 202•8
Sample	40	3	100	100	395	Ham	NNM
Date	8/17	•	8/18	8/19	8/20	8/21	8/22

TABLE XXIX (Continued) SOIL MOISTURE IN QUADRAT 1 IN 1968

:

• (

*	2					
Avenace	0 10 10 10 10 10 10 10 10 10 10 10 10 10	10.95	13.65	10.90	6.76	6.15
& Moisture	12.28	10.19	14.41 14.26 12.29	9.60 9.60 41.11	6.04 7.47 6.78	5.50 6.83 6.11
Sample Moisture	21.3	19•3 18•9	24.3	21.5	11.1 14.5 12.8	10.0 12.8 11.6
Sample	Ч	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	Ham	4 19 M	Man	HNM
Dry Soil	173.4	182.0	179.1 170.4 184.7	179 <b>•9</b> 186 <b>•5</b> 183 <b>•</b> 2	183.5 194.0 188.8	181.7 187.2 189.9
Sample	-+ 0	3 60	MNM	1 N M	H NI M	-1 01 m
Date	8/24		8/25	8/26	8/27	8/28
Wet Soil	194.7	200.9	204.9 194.7 207.4	201.4 204.4 203.6	194.6 208.5 201.6	191.7 200.0 201.5
Sample	10	3	1 an	HNM	n n m	400
Date	8/23	•	8/24	8/25	8/26	8/27

TABLE XXX SOLL MDISTURE IN QUADRAT 2 IN 1968

-0

;

Averare &		51.6	(+•)			TO·ZT			81.0	07+2			7.20	1200			6 63 9	0.0			まっ	
% Moisture	0.06	6.32	5.83	6	TZ-CT	11.12		7.97	10.10	6.47		5.93	8.03	1.91		6.85	5.35	04.4		9.60	まる	10.27
Sample Moisture	4-11	10.1	11.2	0 10	0.00	20.0		12.6	13.2	13.7		6.0T	12.1	12.2		11.8	9.5	11.7		18.2	12.4	TOT
Sample	н	~	e	-		3		Ч	2	m	ſ	-1 -	N	m	f	-1	2	m	ŕ	-1 (	2 6	2
Dry Soil	123.2	159.9	192.2	14.6	178.5	179.8		1.58.1	129.7	144.6	6 E81		0.001	2.57	0 000	C.2/T	T/1.5:	151.9	- 2 081	0.00	156.7	
Sample	-1	~	3	ч	2	e	,	-1 (	N	<b>F)</b>	ſ	0	1 ¢	<b>0</b> 0	F	1 C	2 (	2	-	10	3 67	
Date	7/31			8/J			010	210			8/3				8/4	- 1-			8/5			
Wet Soil	134.6	0.074	4.000	175.8	198.5	8.64L	6 06L	0 071	C 27 C	C.0C+	194.6	162.7	166.4		184.1	187.0	7 696	0.01	207.8	170.3	172.8	
Sample	Ч¢	4 (*	2	-	2 1	2	н	10	1 07	`	Ч	2	5		ч	~	~	1	~	~	ñ	
Date	7/29			7/31			8/1	-			8/2				8/3				9/4			

TABLE XXX (Continued) SOLL MOISTURE IN QUADRAT 2 IN 1968

;

Average % 2.09 61.4 6.04 4.24 7.52 19.46 % Moisture 5.99 6.99 8.30 4.62 7.93 000 233 19.71 Sample Moisture 9.06 13.1 9.20 6.7 8.4 8.7 10.8 32.4 Sample Date Sample Dry Soil 192.0 124.1 164.7 159.0 167.4 170.1 178.7 135.2 i i 3 01/8. 8/11 8/6 8/7 8/8 8/9 Wet Soil 203.5 200.7 199.6 187.7 166.9 171.5 207.8 204.5 198.1 196.8 187.7 165.7 207.5 180.9 Date Sample NNM MNM ANM NO Nm - NM 8/10 8/5 8/6 8/7 8/8 8/8

TABLE XXX (Continued) SOIL MOISTURE IN QUADRAT 2 IN 1968

:

• (

1 8																		
Averace		10.21	T70(T		22.57			75.77			14.78			17.66			15.00	
% Moisture	14-41	14.50	16.62	24.39	20.36	22.95	15.88	15.53	14.09	15.23	15.40	13.71	18.74	18.58	15.65	14.62	15.10	15.28
Sample Moisture	24-9	24.3	26.8	39.0	32.9	33.6	25.4	27.9	22.6	27.3	23.9	21.7	30.4	30.4	26.9	22.2	26.5	27.4
Sample	н	2	3	н	~	e	н	~	e	н	N	б	7	~	б	Ч	~	3
Dry Soil	172.8	166.6	161.3	159.9	161.6	146.4	160.0	179.6	160.4	179.3	155.2	158.3	162.2	163.6 :	171.9	151.8	175.5	179.3
Sempl.e	н	2	б	н	~	n	ч	2	3	ч	2	èn:	ن ب	2	m	Ч	~	m
Date	8/12			8/13			8/1 ⁴			8/15		•	91/8			8/17		
Wet Soil	7.72L	190.9	188.1	198.9	194.5	180.0	185.4	207.5	183.0	206.6	T-6/T	0.031	192.6	194.0	198.8	0°7/1	202.0	206.7
Sample	н	2	m	Ч		n	-	~	n	Ч	20	n	Ч	N	2	-	21	2
Date	11/8	•		8/12			8/13			4T/8			8/15			8/16		

TABLE XXX (Continued) SOIL MOISTURE IN QUADRAT 2 IN 1968

•(

Í

;

~	R																						
Averace	D X X 40 A 12	51 55	THOCT			12.41			00 01	00.04			10.05				9.45				8.33		
% Moisture	10 67	10.21	14.30	5 5 5	13.11	62.11	10.01	20.83	04.0	10.37		9.23	10.51	10.42		9.19	8.30	10.86		7.98	8.71	8.30	
Sample Moisture	2.12	24.4	25.6		1.04	4.VL	(	20.5	17.6	18.3		C.CT	19.3	19.4		7.07	13.7	19.9		1.++	16.4	15.7	
Sample	Ч	2	m	F	1 0	4 67	•	Ч	~	ŝ	r	4	N	m	r	-1 (	N	3	r	4	2	e	
Dry Soll	172.6	182.6	179.0	178.5	2 1791	180.7		189.3	181.5	176.4	168.0		0.001	186.2	0.961		A.Cot	163.2	184.2		TOOT	189.2	
Sample	н	2	m	r,	~	3 m		н	2	ŝ	m	0	N C	20	13 m	0	2 6	2	Ч	10	2 0	n	
Date	87/8			8/19				8/20			8/21				8/22				8/23				
Wet Soil	194.3	207.0	204.0	201.9	184.0	203.0		209.8	1.99.1	1.467	183.5	202.9	200	0.000	192.4	178.7	203.7	+=//~	198.9	204.7	2040 0	× • • • > > >	
Sample	rt	Ne	2	ч	2	e		-	NC	n	Ч	~	(*	1	-1	2	~	`	ч	2	6	1	
Date	8/17			8/18				8/19			8/20				8/21				8/22				

TABLE XXX (Continued) SOLL MOISTURE IN QUADRAT 2 IN 1968

i

:

Average % 8.26 6.85 6.88 6.88 47.9 Sample Moisture & Moisture 8.19 8.30 7.05 2.52 5.82 8.89 892 892 5.92 Sample 14.9 10.5 13.8 12.7 12.7 14.2 1112 -1 NM MNM -1 NM MNM Nm Date Sample Dry Soil 182.0 183.1 177.2 195.5 181.2 184.9 195.5 192.8 188.4 180.4 193.3 11 MNM rt n m Nº M n nm HNM 8/24 8/25 8/26 8/27 8/28 Sample Wet Soil 196.9 198.3 191.9 209.3 193.0 197.8 208.2 205.5 202.6 190.9 205.0 HNM p-4 Nm co m Nm -Nm Date 8/23 8/24 8/25 8/26 8/27

TABLE XXXI SOIL NOISTURE IN QUADRAT 3 IN 1968

;

1	R																		
Chemory	A 49 49 49	00 00	07.OT	•	19.91	10.01			15.52			11.61			10 02	ol.nt		13.64	
& Moisture	0	101 01	64.6	0 1	17.89	16.31		15.40	15.79		13.30	12.41	9.02		24.4	10.93	13 68	13.43	13.50
Sample Moisture	0.71	23.2	15.5	0 10	28.6	28.7	1	C.C2	24.7		24.1	20.0	13.5	8 36	19-61	12.4	25.7	24.2	22.2
Samp1e	۲	101	3	-	1 01	e	•	4 0	N M		Ч	~	m	~	1 (1)	m	Ч	2	3
Dry Soil	181.9	185.7	104.4	1.04L	159.9	176.0	7.62.6	1 44 1	178.9	•	180.0	161.1	2.64L	169.3	171.5	159.2	179.6	180.2	104.4
Sample	, H	2	m	rt	~	3	F	10	ŝ	,	-	2 0	n	'i H	2	ŝ	Ч	~	m
Date	7/31			8/1			8/2	•		010	6/2			8/4			8/5		
Wet Soil	198.1	208.9	5.67L	162.0	188.5	1.000	191.1	181.1	206.4	- 100	T-tor	T. TOT	3.004	186.1	1.191.1	9.0/T	204.7	204°4	0.007
Sample	н	~	n		N 6		Ч	2	т	•	10	4 67	•	ч	~ ~	n	r-i (	N (*	2
Date	7/29			7/31			1/8			8/2	2			8/3			8/4		

TABLE XXXI (Continued) SOIL MOISTURE IN QUADRAT 3 IN 1963

.1

--

i

.

	1												-					
Average &		72.26	00.44		01 01	9C*07		8.28			5.35			8.57			20.60	
% Moisture	24-11	30-16	04-11	A RIV		101-11	8.17	8-03	8.63	5.46	5.53	5.06	3 <b>7.</b> 66	7.33	16.7	20.67	20.97	20.15
Sample	20-6	1.12	20.7	15.7		19.1	15.4	14.1	16.0	10.3	10.6	10.1	21.3	13.9	14.9	32.0	36.3	34.5
Samp].e	н	2	3	-	10	n e	Ч	N	m	н	03	m	ri	N	67	н	2	3
Dry Soil.	180.4	187.4	181.6	177.6	7744	172.0	188.6	175.6	185.5	188.5	7.191	199.5	182.7	189.6.	188.4	154.8	173.1	2.171
Samol.e	, <b>H</b>	~	e	Ч	~	3 m	Ч	~	m	ч	01	6	Ч	0	e.	ч	2	m
Date	8/6			8/7			8/8			8/8			8/10			<b>LL/8</b>		
Wet Soll	201.0	208.5	202.3	193.3	195.0	1.161	204.0	189.7	201.5	198.8	202.3	209-6	204.0	203.5	203.3	186.8	209.4	205.7
Sample	н	N	m	H	2	3	Ч	2	m	rl -	~	m	н	~	n	ri -	2	2
Date	8/5		•	8/6			8/7			8/8			8/9			8/10		

TABLE XXXI (Continued) SOIL MOISTURE JN QUADRAT 3 IN 1963

....

• 1

					_	
Averace 4	20.59	21.67	24°61	17.09	19.23	17.23
% Moisture	22,55 20,52 18,70	22.33 21.42 21.25	18.40 18.10 21-73	18.06 16.83 16.39	18.91 17.88 20.89	17°11 16°78 17°79
Sample Moisture	37.3 33.0 30.7	37•9 35•3	28.3 29.7 36.4	31.4	28.9	29.4 29.1 30.8
Samole	Ham	Ham	1 2 2 2	Han	HNM	нас
Dry Soil	165.4 160.8 164.1	169.3 164.8 165.2	153.8 164.1 167.6	173.9 164.6 180.0	159.7 133.7 136.9	171.8 173.4 173.1
Samplu	Ham	1 am	n n m	rd arm	ANM	Han
Dute	8/12	8/13	ħT/8	8/15	8/16	8/17
Wet Soll	202.7 193.8 194.8	207.2 200.1 200.3	182 <b>.)</b> 193 <b>.</b> 8 204 <b>.</b> 0	205.3 192.3 209 <b>.5</b>	189.9 157.6 165.5	201.2 202.5 203.9
Sample	HNM	HNM	n n m	HNM	нар	Ham
Date	11/8	8/12	8/13	8/14	8/15	8/16

TABLE XXXI (Continued) SOIL MOISTURE IN QUADRAT 3 IN 1962

ί

:

8				-													
Average	15.30			12.89			12.25			10.01			9.58			8.38	
% Moisture	15.50	15.70	13.31	13.09	12.27	12.83	12.49	24.12	70.11	10.70	37.06	10°01	8.65	9.65	8.18	8.51	8.44
Sample Moisture	23.7	25.2	24.0	22.3	19.7	23.6	22.8	19.7	19.7	19.9	20.0	19.5	24.2	18.2	15.3	16.1	15.8
Sample	40	т	Ч	N	С	Ы	2	m	Ч	0	ო	Ч	~	m	Ч	~	Э
Dry Soil	175.9	160.5	180.3	120°4	160.6	0°478T	182.5	172.5	6-24t	186.0	180.8	186.7	164.1 :	188.6	187.1	192.8	187.3
Sample	10	m	Ч	2	m	Ч	01	<u>n</u>	Ч	~	3	5	2	Ś	н	2	n
Date	8/18		8/19			8/20			8/21			.8/22			8/23		
Wet Soil	176.6	185.7	204.3	192.7	180.3	207.6	205.3	192.2	197.6	205.9	200.8	206.2	178.3	206.8	202.4	209.2	203.1
Samp].e	10	e	Ч	~	m	-1	01	m	Ч	2	m	Ч	N	3	н	2	n
Date	4T/8		8/J8			8/19			8/20			8/21			8/22		

TABLE XXXI (Continued) SOIL WOISTURE IN QUADRAT 3 IN 1968

i

•

- 1

1 1					
3					
Åverage	7.61	8.53	6456	6.06	5.85
& Moisture	7.19 8.32 7.32	8.61 10.62 6.35	5.24	6.63 5.47	6.33 6.31 861
Sample Moisture	1.51	16.0 19.7 12.1	10.8.91	12.7	252
Sample	H Q M	Han	nam	Han	n n n
Dry Soil	182.3	187. 287. 29. 29. 29. 29.	196 <b>.3</b> 187.4 190.2	191.5 182.4 186.6	195.9 195.9 187.1
Samole	Ham	L N M	HNM	H 04 M	1 N N
Date	8/24	8/25	8/26	8/27	8/28
Vet Soil	195.4	201.9	206.6 198.2 209.0	204.5 192.5 8	208.4 207.2 154.1
Sample	Han	- an	Ham	HNM	4 N M
Date	8/23	8/24	8/25	8/26	8/27

## TABLE XXXII

AVERAGE PER CENT MOISTURE IN ALL QUADRATS IN 1968

	C]		Q2		Q3	a se	
Date	Daily I	weekly	Daily	Weekly	Daily	Weekly	
7/29 7/31 8/1 8/2 8/3 8/4 8/5	7.61 12.17 12.93 10.15 6.63 10.66 9.54	9.96	7.13 12.01 9.18 7.29 6.63 9.94 7.09	8.47	10.28 16.61 15.52 11.61 10.76 13.64 11.36	12.83	
8/6 8/7 8/8 8/9 8/10 8/11 8/12	5.60 8.58 6.56 6.61 21.93 12.11 14.48	10.84	6.04 4.24 4.49 7.52 19.46 15.21 22.57	11.36	10.58 8.28 5.35 8.97 20.60 20.59 21.67	13.72	
8/13 8/14 8/15 8/16 8/17 8/18 8/19	17.73 16.85 15.76 13.94 13.27 14.13 11.84	14.79	15.17 14.78 17.66 15.00 13.41 12.41 10.30	14.10	19.41 17.09 19.23 17.23 15.39 12.89 12.25	16.21	
8/20 8/21 8/22 8/23 8/24 8/25 8/26 8/27	10.22 11.53 9.40 10.95 13.65 10.90 6.76	10.49	10.05 9.45 8.33 8.26 6.85 6.88 6.88 6.88	8 <b>.10</b>	10.94 9.58 8.38 7.61 8.53 6.96 6.06	8.29	
0121							

;

## TABLE XXXIII

# ANT COLONY SIZE BY ACTUAL COUNT

Colony	Size	Average
Aphaenogaster (A.) subterranea valida Wheeler	136	
Aphaenogaster (A.) subterranea valida Wheeler	58	
Aphaenogaster (A.) subterranea valida Wheeler	1221	
Aphaenogaster (A.) subterranea valida Wheeler	697	
Aphaenogaster (A.) subterranea valida Wheeler	243	477
Camponotus (T.) vicinus Mayr	7	4/2
Camponotus (T.) vicinus Mayr	67	
Camponotus (T.) vicinus Mayr	230	
Camponotus (T.) vicinus Mavr	60	
Camponotus (T.) vicinus Mayr	1/12	200
Crematogaster lineolata (Sav)	742	100
Crematogaster lineolata (Sav)	6029	01.07
Formica fusca Linnaeus	0720	3401
Formica fusca Linnaens	20	(0
Formica (P.) lasioides Emerry	200	62
Formica (P.) lasicidas Emarg	200	
Formica obscurines Forel	309	221
Formica (N.) nallidefulua Latmaille	3330	3330
Formica (N.) pallidefulre Laterille		
Formice (N.) pollidefulre Latrollie	92	
Formica (N.) pallidetulva latrellie	58	
Forming (N) pollidefulre (atrellig	- 90	
Forming (N) pollidefulva Latrellie	194	8
Tridomumor manine anglis (D. 1)	15	76
Logina alienne meniorus (E. Andre)	305	305
Lastus allenus anericanus mery	43	43
Lastus (C.) brevicornis microps wheeler	70	
Lastus (C.) brevicornis microps wheeler	141	
Lasius (C.) previcornis microps wheeler	141	
Lasius (C.) previcornis microps Wheeler	82	
Lasus (C.) previcornis microps wheeler	56	
Lasius (C.) previcornis microps Wheeler	117	
Lasius (C.) brevicornis microps wheeler	710	1.0
Lasius (C.) brevicornis microps Wheeler	524	
Lasius (C.) brevicornis microps Wheeler	230	230
Lasius (A.) claviger coloradensis Wheeler	422	422
Lasius (A.) latipes (Walsh)	43	43
Lasius niger neoniger Emery	1293	
Lasius niger neoniger Emery	87	
Lasius niger neoniger Emery	364	
Lasius niger neoniger Emery	383	
Lasius niger neoniger Emery	138	
Lasius niger neoniger Emery	129	300
		111

## TABLE XXXIII (Continued)

## ANT COLONY SIZE BY ACTUAL COUNT

Colony	Size	Average
Lasius (C.) umbratus aphidicola (Walsh)	63	
Lasius (C.) umbratus aphidicola (Walsh)	429	
Lasius (C.) umbratus aphidicola (Walsh)	493	
Lasius (C.) umbratus aphidicola (Walsh)	6423	1852
Leptothorax rugatulus Emery	273	
Leptothorax rugatulus Emery	189	231
Liometopum occidentale luctuosum wheeler	1215	1215
Solenopsis (D.) molesta validiuscula Emery	4444	
Solenopsis (D.) molesta validiuscula Emery	281	
Solenopsis (D.) molesta validiuscula Emery	133	
Solenopsis (D.) molesta validiuscula Emery	. 26	221
Tapinoma sessile (Say)	80	
Tapinoma sessile (Say)	211	147

10