

Alterations in the VO_2 -Power Relationship above the Lactate Threshold during a Graded Bicycling Exercise Protocol

By

Eric Contini

Department of Integrative Physiology

Primary Thesis Advisor

William C. Byrnes, Ph.D.

Department of Integrative Physiology

Committee Members

David E. Sherwood, Ph.D.

Department of Integrative Physiology

Darna L. Dufour, Ph.D.

Department of Anthropology

University of Colorado, Boulder

April 2013

ABSTRACT

Purpose: The purpose of this study was to examine the changes in VO₂-power and energy expenditure-power relationships below (B_{LT}) and above (A_{LT}) the lactate threshold (LT). **Methods:** Thirty active subjects (15 males, 15 females) performed a graded exercise protocol on a bicycle ergometer. Initial power outputs were determined based upon subject size and training status. Workloads were increased by 30-Watt increments every 4 minutes until exhaustion. Indirect calorimetry and blood lactate measures were performed during each stage. Linear regressions were developed B_{LT} and A_{LT} for VO₂-power and energy expenditure-power (EE-power) relationships. Only 18 subjects (13 males, 5 females) had a sufficient number of data points B_{LT} and A_{LT} to adequately describe the linear regressions. **Results:** The individual R² values for VO₂-power and EE-power relationships at both B_{LT} and A_{LT} were 0.93 or higher. The mean (±SD) VO₂-power slope (mlO₂/min/Watt) and intercept (mlO₂/min) for A_{LT} were significantly (p<0.01) steeper and lower than the mean B_{LT} slope and intercept (Slope: A_{LT} = 12.7±2.9 vs. B_{LT} = 8.8±1.7; Intercept: A_{LT} = -146.2±617.1 vs. B_{LT} = 656.2±186.8). The mean EE-power slope (kcal/min/Watt) and intercept (kcal/min) for A_{LT} were significantly (p<0.01) steeper and lower than the mean B_{LT} slope and intercept (Slope: A_{LT} = 0.067±0.015 vs. B_{LT} = 0.045±0.009; Intercept: A_{LT} = -1.624±3.248 vs. B_{LT} = 2.934±0.917). No sex effect was observed on the changes in slope or intercept for either VO₂-power or EE-power. **Conclusion:** The relationship between power and either VO₂ or energy expenditure during a graded bicycle test suggests an increased oxidative energy requirement relative to power above the lactate threshold.

INTRODUCTION

Aerobic metabolism can be determined indirectly by measuring an individual's oxygen consumption. Aerobic metabolism occurs primarily during endurance exercise and is the primary energy pathway for physical activities lasting longer than 90 seconds. As such oxygen consumption has been directly linked to the energy requirements for many physical activities. In the laboratory it is common to measure oxygen consumption during incremental exercise on a stationary bicycle or a treadmill. For cycling, oxygen consumption has been shown to increase as work rate (power) is increased. Traditionally, this relationship has been described as being linear (4, 6, 10). Based upon this assumption, predictive relationships have been developed for a variety of parameters such as energy expenditure (13), VO_{2Peak} (5) and oxygen demands at supramaximal intensities (6, 14). However, some have questioned the linearity of this relationship (2, 5, 8, 9, 11, 12). Some researchers have found that the VO_2 -power relationship remains linear for all submaximal workloads (4, 6, 10), while others have found either a disproportionate decrease (2,8,9) or increase (5,11,12) from linearity occurring at higher workloads approaching VO_{2Peak} . It is commonly accepted that oxygen consumption and power are linearly related at light to moderate workloads, however some have found an alteration in the relationship as one transitions from moderate to heavy workloads (2, 5, 8, 9, 11, 12). The alteration from linearity at higher workloads has been associated with a dramatic increase in blood lactate that is a parameter known as lactic threshold (LT) and hydrogen ion concentrations (5,11). Although LT has been documented as a threshold beyond which the relationship between VO_2 and power are altered, causation and the

mechanisms underlying disproportionate increases or decreases in VO_2 are not fully known.

In 1924 a linear relationship between VO_2 and work was first observed (4) and has since been confirmed by subsequent investigations (6,10). More recently Medbo et al. (6) concluded that a linear relationship exists using multiple 10-minute stages of increasing treadmill speeds. In some studies non-linear effects showing higher than expected VO_2 values were found (6, 11), however to fit the linear model disproportionately high oxygen uptakes were attributed to hyperventilation ($>20 \text{ ml}_{\text{STAPD}}/\text{s}/\text{kg}$) and excluded from the data set (6). In these studies it was determined that the oxygen demand during high-intensity exercise could be underestimated for subjects showing the disproportionately high oxygen uptakes since severe hyperventilation is clearly present near exhaustion (7). It is also argued that using indirect calorimetry and finding a linear VO_2 -power relationship is a valid representation of total energy expenditure because it has been found that any energy from anaerobic ATP formation is negligible even when blood and muscle lactate concentrations of several millimolars are present (6).

Conversely, there is evidence that supports a nonlinear VO_2 -power relationship with a decrease in slope at higher exercise intensities (2,8,9). Bickham et al. (2) found a linear VO_2 -speed response in endurance-trained runners below lactic threshold (B_{LT}) but found a significant decrease in the VO_2 -speed slope and increase in the y-intercept above lactic threshold (A_{LT}). It has been proposed that the decrease in the slope of the VO_2 -speed regression could be attributed to an increase in anaerobic contribution A_{LT} (2,8), increased overall efficiency (2,9), and fast twitch fiber recruitment at higher intensities

(2,9). Researchers speculate that an increased overall mechanical efficiency may occur at higher submaximal intensities that can be potentially attributed to training at high cadences and speeds (2). It is suggested that the increase overall mechanical efficiency is due to fast twitch fiber recruitment at the higher intensities combined with a higher contractile efficiency for these fast twitch fibers at the faster pedal rate or speed (9). A disproportionate increase in anaerobic contribution to energy expenditure A_{LT} will result in a decreased slope at higher exercise intensities when using indirect calorimetry. Because indirect calorimetry measures aerobic energy expenditure, a decreasing VO_2 -work slope will be seen if the aerobic portion of total energy expenditure is becoming disproportionately less with increasing energy demand. The reduced slope at higher workloads, and thus, the deviation from linearity, in these studies suggest that a linear regression of VO_2 -speed data B_{LT} may potentially overestimate actual VO_{2Peak} .

Countering findings that show a decreased slope at higher intensities, several studies have suggested that there is a disproportionate increase in VO_2 A_{LT} resulting in a linear relationship that is steeper at higher workloads (1,3,5,11,12). Zoladz et al. (11) discovered a disproportionate increase in VO_2 at higher workloads independent of pedaling rate. Factors that may play a role in the increased VO_2 at high exercise intensities include increases in lactate and/or hydrogen ion concentrations, the presence of a slow component during exercise, and muscle fiber type recruitment (5, 11, 12). It was found that the magnitudes of the change in VO_2 from the expected initial linear relationship at lower workloads followed the degree of lactate accumulation (11). Jones et al. discusses the possibility that the catabolism of lactate and its role as an energy substrate during glyconeogenesis may partially explain the additional energy cost of high

intensity exercises. Zoladz et al. (11) has suggested that the decrease in pH associated with the increase in muscle lactate concentrations may play an important role in shifting the oxyhemoglobin dissociation curve to the right to facilitate the exchange of oxygen at the level of the tissue. The increase in lactate and/or hydrogen ion concentrations may lead to a disproportionate increase in VO_2 by accelerating the rate of mitochondrial respiration by increasing free creatine concentrations (5).

During continuous constant load protocols at workloads A_{LT} there is evidence that VO_2 does not plateau or reach a steady state but continues to rise until $\text{VO}_{2\text{peak}}$ is reached, which is known as the slow component (5,11). During an incremental treadmill test with 7 minute stages, Jones et al. (5) analyzed ventilation between 3.0 and 3.75 minutes and again between 6.0 and 6.75 minutes to assess if VO_2 attains a steady state within 3 minutes. A significant difference in the change in VO_2 was found at running speeds above but not below LT. This alludes to the possibility that stages A_{LT} may have a disproportionately higher VO_2 due to slow component contribution. The slow component may be caused by an increase in additional accessory muscles such as those involved in stabilization, posture and respiration (11). An increase in core body temperature and recruitment of type II muscle fibers have also been linked to the contribution of the slow component (5,12). Current thinking suggests that the energetics of contraction of the different muscle fiber types may be the main driver of the slow component (5). Increases in core body temperature increase the metabolic activity of the body and therefore require a larger O_2 demand. When increasing exercise intensity, the hierarchical recruitment of different fiber types may contribute to the increase in VO_2 (11). The mitochondria of type II muscle fibers have a lower P to O ratio than type I fibers, which is a measure of the

efficiency of coupling phosphorylation to oxidation (5). This would suggest that type II fibers would have a greater VO_2 for any given rate of ATP resynthesis and that type II fibers are less efficient than type I fibers. It has been shown that type II muscle fibers are active during the exercise intensity domain that is associated with the slow component, and that both gross and net efficiency have been shown to be positively correlated with the percentage of type I fibers (5).

My research examined the changes in VO_2 -power and energy expenditure (EE)-power relationships below and above LT. It is hypothesized that a positive deviation from linearity will be observed A_{LT} primarily due to cumulative physiological effects such as an increase in core and muscle temperature, lactate and hydrogen ion concentrations, and slow component effects. This area of research has implications in both research and applied settings as currently linear relationships are still typically used. If there is a difference in the VO_2 -power and EE-power relationships B_{LT} and A_{LT} , the physiological measure to be predicted must be considered carefully. If deviations from linearity exist, predicted physiological measures near maximal or at supramaximal intensities may deviate significantly from actual values. Using linear models to make predictions where a non-linear relationship exist would result in over or underestimations of the actual value.

METHODS

Subjects

Thirty healthy and physically active (15 male, 15 female) (mean \pm SD; age 23.4 ± 4.6 years, weight 69.5 ± 10.5 kg, $\text{VO}_{2\text{Peak}}$ 46.6 ± 8.9 ml/kg/min) volunteered for the

study. Only 18 subjects (13 males, 5 females) had a sufficient number of data points B_{LT} and A_{LT} to adequately describe the linear regressions. This was due to the use of an existing data set obtained from a study that was not originally designed to examine relationships B_{LT} and A_{LT} . Subject characteristics are reported in Table 1. Subjects filled out a Physically Activity Readiness Questionnaire (PAR-Q) to ensure that they were healthy and free of medical contraindications and were classified as physically active by meeting the American College of Sports Medicine (ACSM) and the American Heart Association's (AHA) updated physical activity recommendations. Participants in the study were required to exercise at least three times per week for at least 45 minutes per session for a total of at least 150 minutes per week. All subjects gave a written informed consent, which was obtained by the lead investigator prior to participation. This project was approved by the Internal Review Board of the University of Colorado at Boulder (Protocol Number: 11-0342).

Table 1. Subject Characteristics. Norm, normalized; LT, lactate threshold; Mech Eff, mechanical efficiency; Econ, economy. * Significant difference between M and F, $p < 0.05$.

Subject	Sex	Age (yrs)	BMI (kg/m ²)	VO _{2peak} (l•min ⁻¹)	Norm VO _{2peak} (ml•kg ⁻¹ •min ⁻¹)	LT Power (W)	LT (%VO _{2peak})	Econ at LT (W•l ⁻¹ •min ⁻¹)	Lab HR _{peak} (bpm)
1	F	18	18.7	2.07	41.87	124	62%	97.0	195
2	F	19	22.6	3.00	54.93	146	64%	76.0	183
3	F	21	28.4	2.72	32.56	156	69%	83.1	190
4	F	27	23.6	2.97	44.73	171	68%	84.9	188
5	F	34	23.7	3.07	45.86	195	72%	87.8	179
6	M	19	20.8	3.19	47.12	187	72%	81.2	183
7	M	19	22.2	3.97	54.23	197	63%	78.9	180
8	M	19	24.5	5.01	69.43	230	75%	61.2	190
9	M	20	23.3	4.32	56.88	220	64%	79.5	197
10	M	20	23.3	4.28	47.38	168	73%	53.8	184
11	M	20	25.3	3.99	51.97	225	70%	80.9	194
12	M	22	23.2	3.17	41.45	171	72%	74.7	176
13	M	22	23.2	3.68	51.65	183	67%	74.0	201
14	M	22	22.0	3.82	53.50	230	72%	84.1	167
15	M	22	21.0	2.91	43.83	182	71%	88.5	173
16	M	22	26.0	4.18	57.86	218	68%	76.7	192
17	M	22	30.9	4.27	41.38	218	65%	78.7	171
18	M	28	23.0	2.46	34.89	148	70%	85.5	176
Female	Mean	24	23.4	*2.77	43.99	*158.4	67%	85.8	187.0
	SD	7	3.4	0.41	8.04	26.7	4%	7.6	6.2
Male	Mean	21	23.7	3.79	50.12	198.2	69%	76.7	183.4
	SD	2	2.6	0.69	8.91	27.1	4%	9.6	10.7

* Significant difference between M and F.

Test Methods

This study was part of a larger investigation that aimed to compare actual energy expenditure to predicted energy expenditures from heart rate-power relationships during bicycling and non-bicycling exercise tasks under steady state and non-steady state conditions. Only data obtained from the first session of the larger study consisting of a graded exercise test (GXT) was used for the current protocol.

All testing was done on either a Lode bicycle ergometer or the subject's own bicycle equipped with a CycleOps power meter mounted on a CompuTrainer® electronically braked bicycle trainer. Energy expenditure and VO_2 were measured using open circuit indirect calorimetry requiring subjects to be fitted with a head harness to properly secure a rubber mouthpiece in their mouth and a plastic nose clip to prevent airflow in and out of the nose during exercise. The mouthpiece was attached to a respiratory valve connected to hosing that leads to a gas analyzer and a pneumotach. A Parvo metabolic cart (Parvo Medics TrueOne® 2400) analyzed the expired gases (reported as 15 second averages) during the GXT. The metabolic cart was manually calibrated by performing gas and volume calibrations before each exercise test to ensure accurate results. To gather blood lactate data for all subjects, those with low finger blood flow wore a nitrile glove with a hand-warmer inserted under their palm to increase blood flow to their fingers for lactate sampling (explained below). Because subjects were unable to speak while oxygen consumption is measured, subjects pointed to a Borg's Rate of Perceived Exertion scale (RPE) to evaluate their perceived exertion. Subjects were given strong verbal encouragement to go as long as possible during the GXT to elicit their best performance.

Graded Exercise Test

Prior to the GXT, subjects warmed up for 2-5 min at a lower power output than the initial stage. The subject began the test at a workload between 50 and 105 watts for women and 60 and 100 watts for men. This was done to allow the subject to complete approximately 6 to 8 stages. With more stages completed, there was a greater chance that there would be a sufficient number of data points B_{LT} and A_{LT} to adequately describe the linear relationships B_{LT} and A_{LT} . Subjects were instructed to choose a comfortable cadence between 60 and 110 rpm that they would maintain throughout the test (7). Each stage lasted 4 minutes and the workload was increased by 30 watts with each new stage. Between the 3rd and 4th minutes of each stage, the subjects RPE was recorded and a blood lactate sample was taken. Once the subject reached volitional fatigue and could not maintain their cadence or stopped on their own volition, the test ended.

Blood lactate

Capillary blood samples were obtained prior to exercise, over the last 60 seconds of each workload during the GXT, and two minutes after the end of the test for blood lactate assessment. Blood was drawn into a heparinized capillary tube from blood drops on the pricked finger and 25 μ l of whole blood was mixed with 50 μ l of a buffer solution, lysing agent (Octylphenoxethanol), and glycolytic inhibitor (NaF anhydrous) to prevent further metabolism from occurring. The blood-buffer solution was vortexed and analyzed in duplicate with a YSI 2300 Stat Plus lactate analyzer that was calibrated every 5 samples. Individual subjects' blood lactate concentrations for each workload were plotted to determine LT (15). LT is the point of inflection (breakpoint) where the lactate

concentration begins to increase exponentially during intense exercise and the LT was defined as 1mmol/l above the LT breakpoint. LT was used to separate the high intensity workloads from the lower intensity workloads, more specifically if a change from linearity were to occur we would expect to see it A_{LT} not B_{LT} .

Statistics


Linear regressions and correlations were developed using Microsoft Office Excel. Paired samples t-test and independent t-tests were used where appropriate using either Microsoft Office Excel or IBM SPSS Statistic's version 20 and 21 software. Slopes and y-intercepts for the linear regressions of VO_2 -power and EE-power were compared using paired t-test at a significance level of 0.05.

RESULTS

The individuals R^2 values for VO_2 -power relationship at both B_{LT} and A_{LT} were 0.93 or higher. The mean (\pm SD) VO_2 -power slope (mlO₂/min/Watt) and intercept (mlO₂/min) for A_{LT} were significantly ($p < 0.01$) steeper and lower than the mean B_{LT} slope and intercept (Slope: $A_{LT} = 12.7 \pm 2.9$, $B_{LT} = 8.8 \pm 1.7$; Intercept: $A_{LT} = -146.2 \pm 617.1$, $B_{LT} = 656.2 \pm 186.8$) (Figure 1, Figure2). Additionally, by utilizing the respiratory quotient the subjects' metabolic substrate utilization could be accounted for as a possible reason for a change in the VO_2 -power relationship. A regression of energy expenditure (EE)-power was made and the same trends were observed. The mean EE-power slope (kcal/min/Watt) and intercept (kcal/min) for A_{LT} were significantly ($p < 0.01$) steeper and lower than the mean B_{LT} slope and intercept (Slope: $A_{LT} = 0.067 \pm 0.015$, $B_{LT} =$

0.045±0.009, Intercept: $A_{LT} = -1.624 \pm 3.248$, $B_{LT} = 2.934 \pm 0.917$) (Figure 3, Figure 4). No sex effect was observed on the changes in slope or intercept for either VO_2 -power or EE-power.

Figure 1. The mean slope and y-intercept for VO_2 -power. The slope and y-intercept B_{LT} is significantly different from A_{LT} .



* Significantly steeper slope ($P < 0.000$) and significantly decreased y-intercept ($p < 0.000$) compared to B_{LT} .

Figure 2. The mean \pm SD for slope and y-intercept changes for VO_2 -power relationships. A. The slope change B_{LT} and A_{LT} ; B. The y-intercept change B_{LT} and A_{LT} .




Figure 3. The mean slope and y-intercept for EE-power. The slope and y-intercept B_{LT} is significantly different from A_{LT} .





Figure 4. The mean \pm SD for slope and y-intercept changes for EE-power relationships. A. The slope change B_{LT} and A_{LT} ; B. The y-intercept change B_{LT} and A_{LT} .



DISCUSSION


A significant increase in the slope of the VO_2 -power and EE-power regressions A_{LT} was demonstrated. The average increase in the slope of the VO_2 -power regression A_{LT} was 44.3% when compared to B_{LT} resulting in a significantly different predicted $\text{VO}_{2\text{Peak}}$ between these regressions. The average increase in the slope of the EE-power regression A_{LT} was 44.9% when compared to B_{LT} . This agrees with the data that shows a significant decrease in the y-intercept of the VO_2 -power and EE-power regressions A_{LT} was demonstrated. Using the power at $\text{VO}_{2\text{Peak}}$, the VO_2 predicted using the A_{LT} regression was not significantly different than the actual $\text{VO}_{2\text{Peak}}$. However, VO_2 predicted using the B_{LT} regression was significantly different from $\text{VO}_{2\text{Peak}}$, deviated by -7.7%. This data supports the hypothesis that a positive deviation from linearity occurs above A_{LT} and can provide a better understanding of what may contribute to an increased VO_2 A_{LT} and the possible physiological mechanisms responsible for the observed trend.

These findings are consistent with previous reports from Jones et al. (5) and Zoladz et al. (1995) (11) who reported that the deviations in VO_2 from the expected initial linear relationship at lower workloads followed the degree of blood lactate accumulation. Nevertheless, the finding of the present study and those of earlier studies (5,11,12) are in contradiction to the more traditional views that the VO_2 -power relationship remains linear for all submaximal workloads up to $\text{VO}_{2\text{max}}$ (4,6,10). It has been suggested that the non-linearity of the VO_2 -power relationship has not been recognized more often due to the type of protocol used and that any isolated data points which deviate from linearity may be dismissed either as experimental noise or the consequence of a disproportionate increase in ventilatory, cardiac or postural costs (11).

In many multi-stage tests it is possible that very few data points are recorded at high workloads A_{LT} preventing the identification of non-linearity due to the inability of subjects to maintain exercise past 2-3 stages A_{LT} . Future considerations to account for a lack of data points A_{LT} include using smaller incremental workload increases as well as first determining lactic threshold and VO_{2Peak} and then having the subject at exercise intensities that lie within those two parameters.


To address some of the differences between protocols and to better understand reasons for disproportional VO_2 uptake, a within group pilot study was performed in order to examine some of the differences between using a continuous (CONT) versus discontinuous (DISCONT) GXT protocol. For the pilot study, 4 endurance-trained subjects performed the CONT GXT protocol beginning at 90 W. To maintain consistency with the original study, the workload increased by 30 W every 4 minutes until volitional exhaustion. The following two days after CONT, the subjects returned at the same time during the day and performed the same power outputs achieved during the CONT for 4 minutes, but each stage was separated by 10 minutes of seated rest. The power outputs were paired together and randomized across the two days. The pilot data supported the original findings providing evidence that there is a disproportionate increase in $VO_2 A_{LT}$ during the CONT, while data from the DISCONT shows a linear VO_2 -power relationship (Figure 3, 4, 5). These findings show that the VO_2 -power relationship may be protocol dependent and may allude to possible reasons for the non-linear relationship during a CONT.

Figure 3. The mean VO₂-power relationship for CONT and DISCONT. Significant differences in VO₂ are found at 240 watts and 270 watts.




* Significant difference between CON and DISCON (P<0.05)

Figure 4. The mean \pm SD slope changes for VO₂-power relationships for CONT and DISCONT. Pre, A_{LT}; Post, B_{LT}. The slope change B_{LT} and A_{LT} For both CONT and DISCONT are significantly different.



*Significantly different than Pre-CONT; ** Sign different than Post-CONT

Figure 5. The mean \pm SD y-intercept changes for VO_2 -power relationships for CONT and DISCONT. Pre, A_{LT} ; Post, B_{LT} . The slope change B_{LT} and A_{LT} For both CONT and DISCONT are significantly different.



*Significantly different than Pre-CONT; ** Sign different than Post-CONT

The reasons for a disproportionate increase in $\text{VO}_2 A_{LT}$ is not clear, however, it is suggested that the change in the slope and intercept may be related to the same factors that attribute to the slow component (2,5,11,12). An increase in VO_2 contributing to the slow component could be partly due to the increased cardio and respiratory work as a result of an increase in overall energy demand. Data from the pilot work shows that there was significant difference for the average VE (l/min) between CONT and DISCONT at the higher workloads. Using data reported by Dempsey et al. (16) and Kitamura et al. (17) it was calculated that the additional O_2 cost of high pulmonary ventilation and heart rates would account for 10-12% of the total additional VO_2 at higher workloads in the pilot study. It was determined that 12% of the increase in oxygen consumption between

CONT and DISCONT we see at 240 watts can be attributed to the increase in energy demand of the respiratory and cardiac systems.

Additionally, the energetics of contraction and the recruitment of type II muscle fibers provide a likely explanation for the slow component and disproportionate increase in $\text{VO}_2 A_{LT}$ (5, 11). The mitochondria of type II muscle fibers have an 18% lower P:O ratio than type I fibers demonstrating that type II muscle fibers are less efficient at producing ATP than type I fibers (5). This suggests that type II fibers would have a greater VO_2 for any given rate of ATP resynthesis. A decreased efficiency of working muscle at higher workloads would then require more overall fibers to be recruited which in turn increases VO_2 demand.

In support of a disproportionate increase in O_2 uptake, proposed mechanisms for the decrease in efficiency include the accumulations of lactate and/or hydrogen ion concentrations and temperature (2, 5, 11, 12). The correlation between lactate and/or hydrogen ion concentrations and the disproportional increase $\text{VO}_2 A_{LT}$ may hold some answers in regard to efficiency. The catabolism of lactate and its role as an energy substrate during glyconeogenesis and the decrease in pH associated with the increase in muscle lactate concentrations may partially explain the additional energy cost of high intensity exercises (5,11). The increase in lactate and/or hydrogen ion concentrations may accelerate the rate of mitochondrial respiration causing a disproportionate increase in VO_2 (5). An increase in core and local muscle temperature may also have affected muscle efficiency and $\text{VO}_2 A_{LT}$. Core and muscle temperature were not measured in this present study, however, there is evidence to suggest that muscle temperature can affect muscle fiber recruitment patterns as well as the power-velocity relationship (5). An

increase in core temperature can increase the body's metabolism leading to an overall increase in VO_2 , known as the Q10 effect on enzyme activity.

The progressive increase in VO_2 seen over longer duration constant workload exercise A_{LT} known as the slow component was not evident in the present study. Although the slow component was not evident, a progressive increase in VO_2 during prolonged exercise at intensities A_{LT} is well documented and widely accepted (5, 11, 12). With that said, it is acknowledged that the length of each stage may not have been long enough to see a significant O_2 drift. The slow component could cause the total VO_2 to be greater than the predicted VO_2 from the extrapolation of the VO_2 -power relationship for steady state exercise B_{LT} .

Based on our finding and the data provided from our more recent pilot work, the energy demands and thus VO_2 appear to be at least in part protocol dependent. A disproportionate increase in VO_2 A_{LT} is present during CONT but not DISCONT. This may be contributed to, but is not exclusive to, differences in body temperature, hydrogen ion concentration, and the reliance on oxidative and non-oxidative pathways. The continuous protocol may result in an overall increased core and/or muscle temperature as well as an increase in hydrogen ion concentration compared to the discontinuous protocol, effectively decrease the efficiency of working muscle. Furthermore, the discontinuous protocol may rely more on non-oxidative energy sources for acute 4-minute bouts than the prolonged exercise session of the continuous protocol through the recruitment of type II muscle fibers.

In conclusion, there was a disproportionate increase in VO_2 with increasing power at workloads A_{LT} . Since there is a difference in the VO_2 -power and EE-power slopes B_{LT}

and A_{LT} , the physiological measure to be predicted must be considered carefully. Using one relationship over the other can result in significant differences in VO_{2Peak} or supramaximal VO_2 . With that said, the protocol chosen may play a major role in the different VO_2 -power and EE-power trends, and thus consideration of the continuous and discontinuous protocol must also be considered.

ACKNOWLEDGEMENTS

This project would not have been possible without the great mentorship and support from Dr. William Byrnes and Eric Homestead. The author wishes to thank all subjects participating in the study. The author would also like to thank the members of the honors committee, Dr. William Byrnes, Dr. David Sherwood, and Dr. Darna Dufour.

REFERENCES

1. Bangsbo J., L. Michalsik, and A. Petersen. Accumulated O₂ deficit during intense exercise and muscle characteristics of elite athletes. *Int. J. Sport Med.* 14:207–213, 1993.
2. Bickham D.C., C. Gibbons, and P. F. Le Rossignol. VO_2 is Attenuated above the Lactate Threshold in Endurance-Trained Runners. *Med Sci Sports Exerc.* 2004;36(2):297–301.
3. Boje O. Energy production, pulmonary ventilation, and length of steps in well-trained runners working on a treadmill. *Acta Physiol. Scand.* 7:362, 1944.
4. Hill A. V., and H. Lupton. Muscular exercise, lactic acid, and the supply and utilization of oxygen. *Q. J. Med.* 16:135–171, 1924.
5. Jones A. M., and J. H. Doust. A 1% treadmill grade most accurately reflects the energetic cost of outdoor running. *J. Sports Sci.* 14:321–327, 1996.
6. Medbo J. I. Is the maximal accumulated oxygen deficit an adequate measure of the anaerobic capacity? *Can. J. Appl. Physiol.* 21:370–383, 1996.

7. Medbo J. I., A. C. Mohn, I. Tabata, R. Bahr, O. Vaage, and O. M. Sejersted. Anaerobic capacity determined by maximal accumulated O₂ deficit. *J. Appl. Physiol.* 64:50–60, 1988.
8. Noakes T. D. Challenging beliefs: ex Africa semper aliquid novi. *Med. Sci. Sports Exerc.* 29:571–590, 1997.
9. Russell A. P., P. F. Le Rossignol, R. J. Snow, and S. K. Lo. Cycling at 120 when compared to 80 rev/min increases the accumulated oxygen deficit but does not affect the precision of its calculation. *J. Exerc. Physiol.* 5:32–38, 2002.
10. Skinner J. S., and T. H. McLellan. The transition from aerobic to anaerobic metabolism. *Res. Q. Exerc. Sport* 51:234–248, 1980.
11. Zoladz J. A., A. C. Rademaker, A. J. Sargeant. Non-linear relationship between O₂ uptake and power output at high intensities of exercise in humans. *J Physiol.* 1995;488:211–217.
12. Zoladz J. A., K. Duda, and J. Majerezak. Oxygen uptake does not increase linearly at high power outputs during incremental exercise test in humans. *Eur. J. Appl. Physiol.* 77:445–451, 1998.
13. Ainslie P., T. Reilly, K. Westerterp. Estimating human energy expenditure: a review of techniques with particular reference to doubly labelled water. *Sports Med.* 33: 683–698, (2003)
14. Chatagnon M., T. Busso, Modelling of aerobic and anaerobic energy production during exhaustive exercise on a cycle ergometer. *Eur. J. Appl. Physiol.* 97:755–60, 2006.
15. Coyle F. E., Integration of the Physiological Factors Determining Endurance Performance Ability. *Exercise & Sport Sciences Reviews.* 23(1):25-64, 1995.
16. Dempsey J. A., C. A. Harms, D. M. Ainsworth. Respiratory muscle perfusion and energetics during exercise. *Med Sci Sports Exerc.* 1996 September; 28(9): 1123–1128.
17. Kitamura K., C. R. Jorgensen, F.L. Gobel, H. L. Taylor, Y. Wang. Hemodynamic correlates of myocardial oxygen consumption during upright exercise. *J Appl Physiol.* 1972;32:516–522.