


RESEARCH ARTICLE

Identification of the primary peptide contaminant that inhibits fibrillation and toxicity in synthetic amyloid- β 42

Daniel J. Adams¹, Travis G. Nemkov^{1,2}, John P. Mayer³, William M. Old^{1*}, Michael H. B. Stowell^{1,4*}

1 The Department of Molecular, Cellular, and Developmental Biology, University of Colorado, Boulder, Colorado, United States of America, **2** The Department of Biochemistry and Molecular Genetics, University of Colorado Anschutz Medical Campus, Denver, Colorado, United States of America, **3** Novo Nordisk Research Center, Indianapolis, Indiana, United States of America, **4** The Department of Mechanical Engineering, University of Colorado, Boulder, Colorado, United States of America

* william.old@colorado.edu (WMO); stowellm@colorado.edu (MHBS)



OPEN ACCESS

Citation: Adams DJ, Nemkov TG, Mayer JP, Old WM, Stowell MHB (2017) Identification of the primary peptide contaminant that inhibits fibrillation and toxicity in synthetic amyloid- β 42. PLoS ONE 12(8): e0182804. <https://doi.org/10.1371/journal.pone.0182804>

Editor: Damian Christopher Crowther, MedImmune Ltd Research and Development, UNITED KINGDOM

Received: December 5, 2016

Accepted: July 25, 2017

Published: August 9, 2017

Copyright: © 2017 Adams et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: NIH EUREKA award (M.H.B.S.), an HHMI CIA award (M.H.B.S.), a Corden Pharma Fellowship (D.J.A.) and a NIH (T32 GM065103)/CU Molecular Biophysics Predoctoral Traineeship (D.J.A.).

Competing interests: The University of Colorado has been issued a patent for the manufacturing of

Abstract

Understanding the pathophysiology of Alzheimer disease has relied upon the use of amyloid peptides from a variety of sources, but most predominantly synthetic peptides produced using t-butyloxycarbonyl (Boc) or 9-fluorenylmethoxycarbonyl (Fmoc) chemistry. These synthetic methods can lead to minor impurities which can have profound effects on the biological activity of amyloid peptides. Here we used a combination of cytotoxicity assays, fibrillation assays and high resolution mass spectrometry (MS) to identify impurities in synthetic amyloid preparations that inhibit both cytotoxicity and aggregation. We identify the A β 42 Δ 39 species as the major peptide contaminant responsible for limiting both cytotoxicity and fibrillation of the amyloid peptide. In addition, we demonstrate that the presence of this minor impurity inhibits the formation of a stable A β 42 dimer observable by MS in very pure peptide samples. These results highlight the critical importance of purity and provenance of amyloid peptides in Alzheimer's research in particular, and biological research in general.

Introduction

Since introduction of the amyloid hypothesis of AD over 20 years ago[1], an overwhelmingly large literature has accumulated, cementing the central importance of A β 42 in the mechanism of the disease [2–7]. Due to its causal role in AD, the A β 42 peptide is a favored reagent to model the disease mechanisms and phenotypes. Synthetically derived A β 42 (sA β 42) has been widely used in animal models and cell culture systems to aid in understanding the biological targets and pathological mechanisms of AD. The s A β 42 is also heavily used for *in vitro* characterization of the basic biophysical properties which imbue A β 42 with such unique and toxic potency. While the synthetic peptide has unquestionably proven to faithfully recapitulate much of the disease-relevant molecular and physiological pathology, data generated in these models notably suffer from poor reproducibility[8]. Because of its extreme propensity to

ultrapure amyloid peptides and TN and MS are inventors.

aggregate in solution, harsh conditions are typically employed in the handling of A β 42 to keep it soluble for use in various assays. These include concentrated urea, strong base, guanidine and even organic solvents such as hexafluoro-2-isopropanol (HFIP). However, no single method is standard in the field. Seeding further variability, numerous conflicting methods are used in quantifying A β 42 concentration, including ELISA, Absorbance 280, and BCA assays. Beyond these complications in handling A β , it is now becoming evident that the source and purity of the peptide can have major impacts on its performance in functional assays [8, 9]. While sA β 42 is widely available from many manufacturers and has been used ubiquitously for many years, peptides produced via recombinant methods in bacteria are now emerging as an interesting alternative. Previous studies indicated that sA β peptide contains impurities that alter its neurotoxicity and ability to aggregate[9]. Here we have validated those findings and extended the line of questioning to determine the identity of the contaminants which appear to inhibit the toxic activities of A β 42. Through both discovery-oriented and candidate-based approaches, we have found that a failed valine-valine coupling at position 39–40 in A β 42 produces a truncated peptide that co-purifies with full-length A β 42 and is a potent inhibitor of its aggregation and cytotoxic activity. These results mandate that future assays using sA β 42 should ensure that this impurity is removed and we propose that this truncated A β derivative, and analogues thereof, merits further investigation in bioassays to characterize its potentially therapeutic properties for treating Alzheimer's disease.

Results

Synthetic A β 42 exhibits reduced toxicity

Motivated by reports of divergent functional properties between recombinant and synthetic A β 42[9], we sought to directly compare their cytotoxic potency. Samples of synthetic and recombinant A β 42 were prepared as a monomer by solubilizing in hexafluoroisopropanol (HFIP) and re-drying to film under inert gas flow. The films were re-solubilized in 10 mM NaOH to prevent aggregation, assayed for peptide content and diluted into culture medium immediately prior to application. The concentration of each peptide sample was determined by BCA assay in a method that has been independently verified by SDS-PAGE, UV-Vis spectroscopy and amino acid analysis. The popular pheochromocytoma-derived cell line PC12 was used in a toxicity assay because it is known to express many neuronal proteins and is easily amenable to high throughput bioassays. A concentration series of recombinant and synthetic peptide as well as the NaOH vehicle was prepared and applied to cells for 24 hours. At the end-point, a standard MTT viability assay was performed (Fig 1). We found that recombinant A β 42 was potentially toxic to the neuron-like PC12 cells, inducing measurable toxicity as low as 25 nM with an apparent LD₅₀ of 190 nM. In contrast, the toxicity of the synthetic peptide was significantly lower, with an apparent LD₅₀ of 280 nM.

Synthetic A β 42 exhibits qualitatively different fibrillation characteristics

The aggregation behavior of A β 42 is well documented and highly investigated, but the relationship between this behavior and the protein's acute cytotoxicity is not entirely clear. Therefore, negative stain electron microscopy was used to determine if there were structural correlates of the reduced toxicity displayed by sA β 42. Accordingly, rA β 42 and sA β 42 stocks at 10 μ M were diluted to 1 μ M in phosphate buffered saline (PBS pH 7.4) to induce aggregation. Samples were removed for imaging after 60 minutes and analyzed by negative stain electron microscopy (Fig 2). Both samples show aggregation into fibrils however the rA β 42 shows longer fibrils with little branching. In contrast the sA β 42 displayed shorter more branched clumps of fibrils.




Fig 1. Recombinant Aβ42 is more cytotoxic than synthetic Aβ42. MTT survival assay after 24h of peptide treatment at indicated concentrations. Averages from two experiments and six replicates. Error bars represent SEM. A pairwise T-test across all dose series gave a $p = 0.004$ comparing the synthetic (SYN) and recombinant (REC), a $p = 0.001$ comparing REC and vehicle (VEH), and a $p = 0.06$ comparing SYN to VEH.

<https://doi.org/10.1371/journal.pone.0182804.g001>

Synthetic Aβ42 exhibits quantitatively different fibrillation dynamics

After observing qualitative differences in the morphology of aggregating fibrils of rAβ42 and sAβ42, we sought to determine if differences might also exist in the quantitative dynamics of fibrillation. To this end we performed a standard fluorescence-based assay to track the kinetics of aggregation in real-time. Thioflavin T (ThT) is a benzothiazole dye that upon binding β-sheet aggregate structures, such as amyloid fibrils, undergoes a measurable fluorescence emission shift and has been used for many years as a marker in histological identification of Aβ plaques in brain tissue. Thus an increase in ThT signal represents increased binding to fibrils and is therefore a direct readout for aggregation. Using the ThT dye assay, we observed distinct aggregation dynamics between rAβ42 and sAβ42 (Fig 3). The kinetics of aggregation is set apart by two features of the fluorescent traces. Firstly, the lag time to reach the half-maximal signal, $t_{1/2}$, is about 10% longer in the sAβ42 reaction indicating alterations to the early aggregation steps. Secondly, the fast-phase aggregation trajectory is greater for the rAβ42 (0.33/min) versus the sAβ42 (0.17/min).

Mutations within the Aβ42 glycine zipper motif alter fibrillation and toxicity

We have observed that the ability of Aβ42 to aggregate correlates with its toxicity. This correlation has been noted before [9–11], however it is not clear by what mechanism the tendency




Fig 2. Recombinant Aβ42 displays more uniform fibril formation than synthetic Aβ42. Negative stain EM analysis of recombinant Aβ42 (left) versus synthetic Aβ42 (right).

<https://doi.org/10.1371/journal.pone.0182804.g002>

toward fibrillation would drive toxicity. To further explore this connection, we assayed if manipulations which impact aggregation behavior could also alter toxicity. For this experiment we utilized the G37L mutant of the Aβ42 peptide. This mutation disrupts the glycine zipper motif (Fig 4) that is important for the normal homo-oligomerization of Aβ42 [10, 12]. It has been observed that the G37L mutant peptide can act as a dominant negative in an aggregation assay [9, 10, 13]. Thus, we performed a 1:1 mixing experiment between wild-type and G37L Aβ42 and applied this peptide to PC12 cells in the same toxicity experiment as above (Fig 1). We found that the G37L peptide exhibited a slight protective effect from Aβ42 toxicity (Fig 5). This result supports the claim that features of Aβ42 involved in oligomerization are also important in toxicity.

A typical lot of commercial Aβ42 peptide is stated as >95% pure from multiple manufacturers. Assuming these numbers are accurate, two distinct scenarios could explain the disparity in aggregation dynamics and toxicity between recombinant and synthetic Aβ42. One possibility is that some contaminant in the recombinant peptide is responsible for its comparatively enhanced toxicity and aggregation behavior. If it were possible to identify this contaminant and add it to the synthetic Aβ42, increased performance of the synthetic peptide would be expected. The other possibility is that a minority product present in the synthetic peptide is inhibiting its natural aggregation and toxicity. To distinguish between these possibilities, a doping experiment was performed in which a small amount (5% mole fraction) of synthetic peptide was added to the recombinant peptide and subjected to aggregation analysis by ThT fluorescence (Fig 6). If a contaminant of the recombinant peptide were responsible for its enhanced aggregation behavior compared to the synthetic peptide, then no change in dynamics would be expected; but if an inhibitor of aggregation were present in the synthetic peptide, it should still be able to produce a measurable effect even diluted 20 fold. Strikingly, we observed that only 5% wt/wt of the synthetic peptide was sufficient to perturb the normal dynamics of Aβ42 aggregation (Fig 6). Both the time to $t_{1/2}$ and slope of rapid aggregation




Fig 3. Aggregation of recombinant, and synthetic and Aβ42. ThT fluorescence trace of Aβ42 aggregation, see [methods](#). REC is recombinant and SYN is synthetic Aβ42. The recombinant materials shows a shorter lag time and faster aggregation rate (0.33/min) than synthetic material (0.17/min).

<https://doi.org/10.1371/journal.pone.0182804.g003>

phase indicate that the dynamics of aggregation have indeed been altered by the minority species present in the synthetic peptide. In contrast, 5% wt/wt doping of recombinant Aβ42 into the synthetic peptide did not alter aggregation behavior, again suggesting the sAβ42 contains an inhibitor of aggregation. We next used this doping strategy to assess toxicity of Aβ42 mixtures in PC12 cells. In agreement with our ThT data, just a small dose of synthetic peptide was capable of reducing the toxicity of the recombinant peptide by a measurable amount ([Fig 7](#)), further confirming the potency of the inhibitory species present in the synthetic Aβ42 preparation.

Identification of Aβ inhibitors

Our data indicate that a potent inhibitor of both toxicity ([Fig 5](#)) and aggregation ([Fig 6](#)) of Aβ42 exists as a minority product present in the commercially obtained synthetic peptide. This finding has wide implications due to the ubiquitous use of synthetic Aβ42 in AD research.




Fig 4. Sequence and features of human A β 42. Sequence of human A β 42 peptide. Histidine residues highlighted in yellow. Glycine 37 highlighted in blue. Valine 39 highlighted in red. Purple bar indicates sequence derived from the transmembrane domain of APP. Blue curved lines indicated glycine zipper motif.

<https://doi.org/10.1371/journal.pone.0182804.g004>




Fig 5. A β 42 G37L protects from A β 42 toxicity as dominant negative. A β 42 G37L Protects from Toxicity as Dominant Negative at high concentrations. Wild-type recombinant A β 42 was mixed in an equimolar ratio with A β 42 G37L. PC12 cells were treated for 24 hours and viability was assayed by MTT. Bars represent 6 biological replicates. Error bars represent SEM. A pairwise T-test across all dose series gave an insignificant $p = 0.06$ comparing the recombinant (REC) and recombinant doped with the G37L mutant (REC + G37L). At the highest concentrations, however a significant protective affect was observed, 25nM $p = 0.01$, 65nm $p = 0.0001$.

<https://doi.org/10.1371/journal.pone.0182804.g005>




Fig 6. Small quantities of synthetic A β 42 inhibit recombinant aggregation. ThT fluorescence trace of A β 42 aggregation, see [methods](#). REC is recombinant and SYN is synthetic A β 42. The recombinant materials was doped with 5% (wt/wt) prior to initiation of the aggregation assay. A dramatic increase in lag time and fast phase was observed.

<https://doi.org/10.1371/journal.pone.0182804.g006>

However, due to the recent commercial availability of recombinant A β 42, it may not be necessary to improve upon commercial synthesis and purification to avoid these inhibitory contaminants. Instead, these unknown inhibitors may prove valuable as a starting point for the rational design of molecules for therapeutic intervention in AD. Furthermore, these inhibitors serve as a unique and novel reagent for probing the relationship between A β 42 aggregation and toxicity.

We began our study of the contaminants of synthetic A β 42 by optimizing a chromatographic method for both purification and analysis. We found conditions under which we could elute peaks of recombinant A β 42 from a C8 reverse-phase column in a mobile phase of acetonitrile with TFA as a counter ion. We then used this method to analyze the synthetic A β 42 sample for the presence of contaminants. The same sharp main peak existed, however a number of additional minor peaks were observed. There was a distinct peak that eluted just




Fig 7. Synthetic Aβ42 contains a potent inhibitor of toxicity. MTT toxicity assay of PC12 cells treated with recombinant Aβ42 ± 5% (wt/wt) synthetic Aβ42. Assay performed in triplicate, error bars represent SEM. A pairwise T-test across all dose series gave a $p = 0.007$ comparing the recombinant (REC) and recombinant doped with 5% synthetic (REC + SYN).

<https://doi.org/10.1371/journal.pone.0182804.g007>

before the main peak, as well a significant shoulder on both the leading and trailing edges of the main peak (Fig 8).

We also analyzed both recombinant and synthetic peptide samples by MALDI-TOF mass spectrometry (Fig 9). This analysis revealed that the synthetic peptide sample contains a diverse collection of contaminants; however due to their relatively low abundance no specific identifications from this MALDI-TOF cocktail were possible. Most of the contaminants appear within approximately 400 mass units of the main peak, corroborating previous suggestions [6] that they may be byproducts of synthesis related to the majority product. A loss or addition of one to three residues would result in truncated or augmented peptides consistent with the observed sizes of contaminants.

It has been reported that synthetic Aβ42 contains a significant fraction (>1%) of D-histidine that is suggested to be the functionally relevant contaminant in the synthetic material[6]. The traditional reverse-phase chromatographic methods used to purify Aβ42 would be incapable of resolving peptides epimerized at any of the three histidine residues in Aβ42. Racemized peptide would also be indistinguishable by mass spectrometry, making it very difficult to assay for this contaminant with commonly used methods. To directly test if an Aβ42 histidine racemate was capable of recapitulating the aggregation inhibition imparted by the synthetic peptide, we obtained a sample of racemized peptide for ThT assay. The peptide was synthesized using a standard solid phase Fmoc-protected method, but at positions 6, 13 and 14 a 50:50 mixture of L- and D-histidine enantiomers was applied. Thus, a mixture of Aβ42 molecules




Fig 8. HPLC analysis of recombinant and synthetic A β 42. RP-HPLC of recombinant and synthetic A β 42 reveals contaminants in synthetic preparation. 10 μ g of each peptide sample was loaded. Fractions collected from synthetic sample are indicated 1, 2 and 3 by dashed lines.

<https://doi.org/10.1371/journal.pone.0182804.g008>

representing every possible combination of histidine stereochemistry was yielded (noted below as A β 42-HIS). The A β 42-HIS was doped into recombinant A β 42 at 5% mole fraction, a much higher concentration than might normally appear in the synthetic peptide, and aggregation was monitored by ThT fluorescence (Fig 10). We found that even this large dose of A β 42-HIS produced a minor alteration in the recombinant peptide's velocity of aggregation. Therefore, we conclude that although A β 42-HIS may be capable of inhibiting A β 42




Fig 9. MALDI mass spectra of Aβ42 samples. MALDI-TOF spectra reveal major contaminants in synthetic Aβ42 sample.

<https://doi.org/10.1371/journal.pone.0182804.g009>

aggregation, it alone is not sufficient to illicit effects of the magnitude observed. Therefore, we believe that other as yet undiscovered inhibitors in synthetic Aβ42 must exist.

Using RP-HPLC (Fig 8) and mass spectrometry (Fig 9), we observed many contaminating minority products in synthetic Aβ42. To determine if any of these possessed inhibitory activity, synthetic Aβ42 samples were fractionated by RP-HPLC and then assayed for inhibition of aggregation using the ThT assay. Specifically, the main peak of synthetic Aβ42 was isolated from the contaminating material that eluted on either side of it, yielding fractions labeled 1, 2 and 3 (Fig 8). As before, fractions were doped into recombinant Aβ42 at 5% wt/wt for the aggregation assay. Significant changes in aggregation dynamics were not observed in recombinant peptide doped with fractions 2 and 3, but we found that fraction 1 contained extremely potent inhibitory activity (Fig 11). It is important to note that this fraction was still a complex mixture; therefore, it remains unclear whether its inhibitory activity is caused by a single or multiple species. Without further analysis by more sensitive methods such as tandem MS-MS, it is impossible to ascertain the identity of the active agents isolated in fraction 1, although based on the MALDI-TOF experimental results (Fig 9) we suspect them to be related peptides. With the quantity of relevant contaminants available for study being extremely limiting we were prompted us to adopt a candidate-based approach towards identification of Aβ42 activity inhibitors.

There are several sequence features which are known to pose challenges to standard solid phase peptide synthesis methods, and therefore can inform rational prediction of peptide synthesis byproducts for a given target. Highly hydrophobic aggregation-prone sequences such as




Fig 10. Aβ42 mutation aggregation kinetics. ThT fluorescence was used to assay aggregation dynamics of 5% Aβ42-HIS mixed into recombinant Aβ42 (REC + HIS) or 5% Aβ42Δ39 (REC + Δ39).

<https://doi.org/10.1371/journal.pone.0182804.g010>

found in Aβ42 are known to be particularly difficult to generate[14]. Traditional synthesis methods proceed from the carboxy-terminus towards the amino-terminus coupling a single amino acid at a time each followed by a round of deprotection. Given that the carboxy-terminal $\frac{1}{3}$ of Aβ42 is derived from the transmembrane helix of APP (Fig 4), early steps in the synthesis of Aβ42 are particularly susceptible to aggregation of the nascent peptide chain[14]. Peptide aggregation competes with binding of synthesis reagents including deprotection agents and subsequent amino acid residues. Because of the iterative nature of the synthesis process, if either deprotection or coupling fails for a particular molecule in one round, that molecule may well participate in future rounds of synthesis, yielding a final peptide lacking only a single amino acid. In addition to these challenges, valine to valine coupling reactions, which are used in Aβ42 synthesis, are known to be of lower efficiency[15] than other peptide couplings in general. These facts led us to predict that omission of valine 39 (hereinafter Aβ42Δ39) generated a putative byproduct of synthesis that could be responsible for the observed inhibitory activity. This hypothesis is supported by the presence of a peak mass observed by MALDI mass spectrometry approximately 100 Da below the main peak. Additionally, the HPLC fraction containing the inhibitory activity eluted slightly before Aβ42, indicating slightly reduced hydrophobicity that is also consistent with deletion of a valine residue.

Absolute mass resolution by MALDI-TOF is inversely correlated with analyte mass[16], and therefore the 3% difference in mass of Aβ42Δ39 from the total peptide would prove




Fig 11. Contaminant is potent inhibitor of aggregation. HPLC fraction 1 and 3 was doped into recombinant A β 42 at 5% wt/wt and aggregation was assessed by ThT fluorescence. Fraction 1 contained a potent inhibitor of A β 42 aggregation.

<https://doi.org/10.1371/journal.pone.0182804.g011>

difficult to resolve using solely this method. Digestion of the sample with trypsin would generate a C-terminal fragment A β 42 29–42 (Fig 4). A β 42 29–42 has an expected mass of approximately 1270 Da which would shift to 1171 Da upon deletion of Val 39, a change of nearly 10% which would be more readily resolved. Therefore, this combinatorial method of enzymatic digestion and subsequent MALDI-TOF analysis was performed in order to reveal putative A β 42 Δ 39 contaminants. Digestion was carried out on synthetic A β 42 overnight at 37°C. The resultant peptides were desalted/concentrated with a C₄ reversed-phase ZipTip and eluted in acetonitrile and analyzed by MALDI-TOF.

After significant optimization of desalting conditions and matrix/spotting parameters we were able to identify the expected four trypsin digestion peaks for fragment AA1–5, observed mass 638.21 (Calc. 637.29); fragment AA6–16, observed mass 1337.23 (Calc 1337.60); fragment AA17–28, observed mass 1324.79 (Calc. 1325.67); and fragment AA29–42, observed mass 1271.55 (Calc. 1269.76). We did not observe any peaks corresponding to partial digestion indicating that the reaction had gone to completion. Disappointingly, the 29–42 peptide of interest, containing A β 42 Δ 39, was found to ionize very poorly by MALDI, leading to very weak signal. We did not see evidence of a peak near 1153Da although we considered this uninformative as the A β 42 Δ 39 would represent only a very small fraction of the already weak signal from the 29–42 fragment.




Fig 12. ESI-Orbitrap MS of synthetic and recombinant A β 42. Top panel, ESI-Orbitrap MS analysis of recombinant A β 42 show the presence of a dimeric species at 1290.52 M/Z and the lack of the A β 42 Δ 39 peptide at 1472.42 M/Z. Bottom panel, ESI-Orbitrap MS analysis of synthetic A β 42 showing the clear presence of the A β 42 Δ 39 peptide at 1472.42 M/Z and many other contaminating species but undetectable levels of the 1290.52 M/Z dimer species.

<https://doi.org/10.1371/journal.pone.0182804.g012>

To overcome the sensitivity hurdles of MALDI-TOF in the identification of potential contaminants, we turned to ESI-Orbitrap mass spectrometry analysis on both recombinant and synthetic A β 42 (Fig 12). This data confirmed the presence of the many contaminating species in synthetic A β 42 observed by MALDI MS and HPLC. The ESI-Orbitrap data allowed us to resolve individual peaks out of the shoulder present on the A β 42 peak seen by MALDI MS from approximately 4200 Da to 4500 Da. Importantly, a peak of 4417.25 Da was observed, consistent with the presence of A β 42 Δ 39. Based on this finding, we proceeded with direct evaluation of A β 42 Δ 39 as an inhibitor of A β 42 aggregation. Also of note was the presence of a peak in the recombinant sample of A β 42 dimer, M/Z 1290.52. This indicates not only that small oligomers can form under the inhibitory conditions used (1% NH₄OH), but that these oligomers can remain intact through electrospray ionization, trapping and detection. The absence of this peak in the synthetic sample appears to serve as further evidence of its reduced ability to aggregate.

Finally, to directly test if A β 42 Δ 39 was capable of inhibiting A β 42 aggregation kinetics, we obtained the modified peptide, mixed it into recombinant A β 42 at 5% mole fraction, and assayed aggregation by ThT fluorescence. We observed a remarkable inhibition with a near

two-fold increase of $t_{1/2}$ from 17.9 minutes (A β 42) to 30.5 minutes (A β 42 Δ 39) (Fig 10). This shift was reminiscent of the effect imparted by the HPLC fraction 1 (Fig 11) further corroborating that A β 42 Δ 39 is the major active component of that fraction.

Discussion

The lack of reproducibility in various studies using amyloid peptides has been well documented[8] and in this study we sought to determine a potential cause of the lack of reproducibility. Prior studies demonstrated the additional purification of synthetic A β 42 could restore both aggregation kinetics and toxicity yet the nature of the inhibitory material in synthetic A β 42 was not investigated[9]. We have found that synthetic manufacturing of A β 42 generates contaminating byproducts that inhibit A β 42 aggregation and we have identified A β 42 Δ 39 as a potent agent of this inhibition that is observed in synthetic A β 42 preparations. The A β 42 Δ 39 material mostly likely arises due to the relative inefficiency of β -branched amino acid coupling during solid phase peptide synthesis. To a lesser extent, another suggested contaminant, A β 42-HIS, also exhibited some inhibition of aggregation. Previously observed variability in assays utilizing A β 42 can most readily be ascribed to the primary contaminants observed in synthetically prepared A β 42. Although A β 42 is particularly prone to synthetic contaminants arising from coupling inefficiencies, the potential for minor impurities to alter both the biophysical and biological properties of synthetically prepared peptides should be born in mind regardless of the peptide investigated. Finally, and most intriguingly, the observed correlation between A β 42 toxicity and aggregation kinetics, suggests that A β 42 Δ 39 and perhaps A β 42-HIS may exhibit neuroprotective activity in AD models. Indeed, c-terminal fragments of A β 42 have been shown to inhibit toxicity[17] and aggregation[18]. Future studies will reveal whether treatment with various analogs of A β 42 Δ 39 or A β 42-HIS can serve to attenuate A β 42 toxicity, possibly yielding an important new class of compounds for therapeutic intervention in Alzheimer's disease.

Materials and methods

Materials

Thioflavin T was and MTT assay were purchased from Sigma-Aldrich. Recombinant A β 42 was obtained from either AmideBio (Lot No. 20130515) or rPeptide (Lot No. 9131142H). Synthetic A β 42 was purchased from Bachem (Lot No. 0572194) and AnaSpec (Lot No. 31475). A β 42 Δ 39 was custom synthesized by AnaSpec. All peptide concentrations were determined using the BCA Protein Assay Kit, CAT# 23225, from Pierce using the provided Albumin standard. All other chemicals were of analytical grade.

ThT assay

This assay was performed as previously described. Briefly Stock solutions of ThT were freshly prepared a 2 mM in 50 mM glycine buffer pH 8.5 and aliquoted and stored at -80°C prior to use. For assays, an aliquot was thawed and diluted 50 fold in 50mM glycine pH 8.5 buffer to a 40 μM concentration. A β (1–42) aggregation was measured by adding 50 μL of 10 μM stock A β (1–42) followed by 50 μL of a 40 μM ThT solution and then monitoring fluorescence intensity using an excitation wavelength of 440nm and an emission wavelength of 468nm using a Molecular Devices SpectraMax M5 plate reader at 42 sec intervals (S1 Table).

MTT assay

This assay was performed according to the Sigma-Aldrich guide. Briefly, MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) was dissolved in phenol free culture media at 5 mg/ml and sterile filtered. To cells grown in 96 well plates, stock MTT solution was added at 10% of the total per well and plates incubated at 37°C for 4 hours. MTT solubilization solution was then added to all wells and mixed thoroughly and the plate was read at 570nm using a Molecular Devices SpectraMax M5 plate reader using a reference wavelength of 630 nm (S2 Table).

HPLC analysis

Samples were analyzed on an Agilent 1100 and fractions isolated using a 4.6 × 250 mm reversed phase Vydac MS C8 column (Grace) using a two phase elution comprised of 1% ACN + 0.1% trifluoroacetic acid (Buffer A) and 100% ACN + 0.1% trifluoroacetic acid (Buffer B) at 70°C. The elution profile was, 0–10 min; 0% B, 10–30 min linear increase to 80% B, with a constant flow rate of 0.8 ml/min and protein was detected at 215 nm.

Electron microscopy

Samples for EM were prepared according similar to the methods of Komatsu[19]. Briefly, a 2% w/v aqueous solution of uranyl acetate at pH 4.2 was filtered (0.2 µm) to remove small precipitates and stored in foil covered container prior to use. Approximately 10 ng (3 µL) of amyloid sample was placed on a freshly glow discharged carbon film on a 300 mesh copper grid for 1 minute. The sample was blotted and then the uranyl acetate solution was applied for 30 seconds and then blotted and air dried. Images were recorded on side mounted CCD camera using a Phillips CM100 at 80 kV.

ESI-Orbitrap mass spectrometry

Amyloid Beta samples were reconstituted in pH 10 1% NH₄OH at 1 mg/mL. Samples were infused at 300 nl/min for nano-electrospray ionization and mass spectrometry analysis performed on a LTQ-Orbitrap (ThermoFisher). Survey scans were collected in the Orbitrap at 60,000 resolution (at m/z = 300). The maximum injection time for MS survey scans was 500 ms with 1 microscan and AGC = 1 × 10⁶. For LTQ MS/MS scans, maximum injection time for survey scans was 250 ms with 1 microscan and AGC = 1 × 10⁶. Peptides were fragmented by CAD for 30 ms in 1 mTorr of N₂ with a normalized collision energy of 35% and activation Q = 0.25.

Supporting information

S1 Table.
(CSV)

S2 Table.
(CSV)

Author Contributions

Conceptualization: Daniel J. Adams, William M. Old, Michael H. B. Stowell.

Data curation: Michael H. B. Stowell.

Formal analysis: Daniel J. Adams, John P. Mayer, William M. Old, Michael H. B. Stowell.

Funding acquisition: Michael H. B. Stowell.

Investigation: Daniel J. Adams, Travis G. Nemkov, William M. Old.

Supervision: William M. Old, Michael H. B. Stowell.

Validation: William M. Old, Michael H. B. Stowell.

Writing – original draft: Daniel J. Adams, John P. Mayer, William M. Old, Michael H. B. Stowell.

Writing – review & editing: Daniel J. Adams, Michael H. B. Stowell.

References

- Hardy J, Allsop D. Amyloid deposition as the central event in the aetiology of Alzheimer's disease. *Trends in pharmacological sciences*. 1991; 12(10):383–8. Epub 1991/10/01. PMID: [1763432](#).
- Masters CL, Selkoe DJ. Biochemistry of amyloid beta-protein and amyloid deposits in Alzheimer disease. *Cold Spring Harb Perspect Med*. 2012; 2(6):a006262. Epub 2012/06/08. <https://doi.org/10.1101/cshperspect.a006262> PMID: [22675658](#);
- Karran E, Mercken M, De Strooper B. The amyloid cascade hypothesis for Alzheimer's disease: an appraisal for the development of therapeutics. *Nature reviews Drug discovery*. 2011; 10(9):698–712. Epub 2011/08/20. <https://doi.org/10.1038/nrd3505> PMID: [21852788](#).
- Walsh DM, Selkoe DJ. A beta oligomers—a decade of discovery. *Journal of neurochemistry*. 2007; 101(5):1172–84. Epub 2007/02/09. <https://doi.org/10.1111/j.1471-4159.2006.04426.x> PMID: [17286590](#).
- Seeman P, Seeman N. Alzheimer's disease: beta-amyloid plaque formation in human brain. *Synapse*. 2011; 65(12):1289–97. Epub 2011/06/03. <https://doi.org/10.1002/syn.20957> PMID: [21633975](#).
- Giuffrida ML, Caraci F, De Bona P, Pappalardo G, Nicoletti F, Rizzarelli E, et al. The monomer state of beta-amyloid: where the Alzheimer's disease protein meets physiology. *Reviews in the neurosciences*. 2010; 21(2):83–93. Epub 2010/07/10. PMID: [20614800](#).
- Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science*. 2002; 297(5580):353–6. Epub 2002/07/20. <https://doi.org/10.1126/science.1072994> PMID: [12130773](#).
- Editor. State of aggregation. *Nature neuroscience*. 2011; 14(4):399. Epub 2011/03/30. <https://doi.org/10.1038/nn0411-399> PMID: [21445061](#).
- Finder VH, Vodopivec I, Nitsch RM, Glockshuber R. The recombinant amyloid-beta peptide Abeta1–42 aggregates faster and is more neurotoxic than synthetic Abeta1–42. *Journal of molecular biology*. 2010; 396(1):9–18. Epub 2009/12/23. <https://doi.org/10.1016/j.jmb.2009.12.016> PMID: [20026079](#).
- Fonte V, Dostal V, Roberts CM, Gonzales P, Lacor PN, Velasco PT, et al. A glycine zipper motif mediates the formation of toxic beta-amyloid oligomers in vitro and in vivo. *Mol Neurodegener*. 2011; 6(1):61. Epub 2011/08/25. <https://doi.org/10.1186/1750-1326-6-61> PMID: [21861874](#);
- Hung LW, Ciccotosto GD, Giannakis E, Tew DJ, Perez K, Masters CL, et al. Amyloid-beta peptide (Abeta) neurotoxicity is modulated by the rate of peptide aggregation: Abeta dimers and trimers correlate with neurotoxicity. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2008; 28(46):11950–8. Epub 2008/11/14. <https://doi.org/10.1523/JNEUROSCI.3916-08.2008> PMID: [19005060](#).
- Kim S, Jeon TJ, Oberai A, Yang D, Schmidt JJ, Bowie JU. Transmembrane glycine zippers: physiological and pathological roles in membrane proteins. *P Natl Acad Sci USA*. 2005; 102(40):14278–83. Epub 2005/09/24. <https://doi.org/10.1073/pnas.0501234102> PMID: [16179394](#);
- Harmer A, Wozny C, Rost BR, Munter LM, Hua H, Georgiev O, et al. Role of amyloid-beta glycine 33 in oligomerization, toxicity, and neuronal plasticity. *The Journal of neuroscience: the official journal of the Society for Neuroscience*. 2009; 29(23):7582–90. Epub 2009/06/12. <https://doi.org/10.1523/JNEUROSCI.1336-09.2009> PMID: [19515926](#).
- Tickler AK, Clippingdale AB, Wade JD. Amyloid-beta as a "difficult sequence" in solid phase peptide synthesis. *Protein Pept Lett*. 2004; 11(4):377–84. Epub 2004/08/26. PMID: [15327371](#).
- Young JD, Huang AS, Ariel N, Bruins JB, Ng D, Stevens RL. Coupling efficiencies of amino acids in the solid phase synthesis of peptides. *Pept Res*. 1990; 3(4):194–200. Epub 1990/07/01. PMID: [2134063](#).
- Reiber DC, Grover TA, Brown RS. Identifying proteins using matrix-assisted laser desorption/ionization in-source fragmentation data combined with database searching. *Analytical chemistry*. 1998; 70(4):673–83. Epub 1998/03/10. PMID: [9491752](#).

17. Li H, Du Z, Lopes DH, Fradinger EA, Wang C, Bitan G. C-terminal tetrapeptides inhibit Abeta42-induced neurotoxicity primarily through specific interaction at the N-terminus of Abeta42. *J Med Chem.* 2011; 54 (24):8451–60. <https://doi.org/10.1021/jm200982p> PMID: 22087474;
18. Gessel MM, Wu C, Li H, Bitan G, Shea JE, Bowers MT. Abeta(39–42) modulates Abeta oligomerization but not fibril formation. *Biochemistry.* 2012; 51(1):108–17. <https://doi.org/10.1021/bi201520b> PMID: 22129303;
19. Komatsu H, Feingold-Link E, Sharp KA, Rastogi T, Axelsen PH. Intrinsic linear heterogeneity of amyloid beta protein fibrils revealed by higher resolution mass-per-length determinations. *The Journal of biological chemistry.* 2010; 285(53):41843–51. Epub 2010/10/14. <https://doi.org/10.1074/jbc.M110.165068> PMID: 20940298;